

Facile Syntheses of Pseudo- α -D-glucopyranose and Pseudo- α -D-mannopyranose

Tony K. M. Shing,^{*a} Yu-xin Cui^a and Ying Tang^b

^a Department of Chemistry, The Chinese University of Hong Kong, Shatin, Hong Kong

^b Department of Chemistry, University of Manchester, Manchester M13 9PL, UK


Pseudo- α -D-glucopyranose **1** and pseudo- α -D-mannopyranose **2** are obtained *via* a stereoselective *cis*- and *trans*-hydroxylation respectively of the alkene **8**, which is readily derived from (–)-quinic acid.

The potential use of pseudo-sugars¹ in biochemical studies of specific enzyme inhibition^{2,3} and as non-nutritive sweeteners⁴ has led to their recent syntheses in enantiomerically pure forms.^{3–6} Optically active and crystalline pseudo- α -D-glucopyranose **1** has been synthesised from D-glucose in 14 stages;⁵ the racemate⁷ has been shown to inhibit glucokinase activity and glucose-stimulated insulin release.² However, the synthesis of enantiomerically pure pseudo- α -D-mannopyranose **2** has not been reported. Recently, we described the enantiospecific syntheses of 2-crotonyloxymethyl-(4*R*,5*R*,6*R*)-4,5,6-trihydroxycyclohex-2-enone (COTC),⁸ pseudo- β -D-mannopyranose and pseudo- β -D-fructopyranose⁹ from quinic acid; this paper further demonstrates the flexibility of this approach by short, facile and stereocontrolled syntheses of **1** and **2**.


The route to pseudo- α -D-glucopyranose **1** and pseudo- α -D-mannopyranose **2** is shown in Scheme 1. The known diol **3**,⁹ readily available from quinic acid **4** in four steps, was protected as the benzyl ether **5**,[†] [α]_D + 5.5° (c 0.29, CH₂Cl₂). The double bond in **5** was subjected to a stereocontrolled hydroboration–oxidation sequence at the less hindered β -face, forming exclusively the cyclohexane derivative **6** which was esterified to the acetate **7**, m.p. 78–79 °C; [α]_D + 3.1° (c 0.26, CH₂Cl₂). The stereochemistry of the 4-OAc group was evident from the ¹H NMR spectrum ($J_{4,3} \approx J_{4,5}$ 9.0 Hz). Acidic removal of the acetal in **7** followed by Corey–Winter deoxygenation¹⁰ of the resulting diol gave the alkene **8** m.p. 57–58 °C; [α]_D – 78.4° (c 0.25, CH₂Cl₂). A stereocontrolled *cis*-hydroxylation of the double bond in **8** occurred smoothly at the less hindered β -face, providing exclusively **9**, m.p. 106–107 °C; [α]_D + 82.4° (c 0.33, CH₂Cl₂). Deprotection then gave crystalline pseudo- α -D-glucopyranose **1**, m.p. 146–147 °C; [α]_D + 63.0° (c 0.6, H₂O) {lit.⁵ m.p. 151–152°; [α]_D + 68.4° (MeOH)}.

trans-Hydroxylation¹¹ of the double bond in **8** with HCO₂H–H₂O₂ proceeded *via* a *trans*-diaxial opening¹² of an

intermediate epoxide, giving the corresponding hydroxy formate which was hydrolysed to the triol **10**, m.p. 123–124 °C; [α]_D + 10.5° (c 0.2, CH₂Cl₂), as the sole product.



Scheme 1 Reagents and conditions: i, see ref. 9; ii, NaH, benzyl bromide, Bu₄N⁺I[–], tetrahydrofuran (THF) (72%); iii, 9-borabicyclo[3.3.1]nonane, THF, room temp., then NaOH (3 mol dm^{–3}), H₂O₂ (94%); iv, acetic anhydride, pyridine (py), *N,N*-dimethylaminopyridine, CH₂Cl₂ (97%); v, CF₃CO₂H, CH₂Cl₂ (97%); vi, 1,1'-thiocarbonyldiimidazole, toluene, reflux, then (OMe)₃P, reflux (85%); vii, OsO₄, trimethylamine-*N*-oxide, py, water, Bu^tOH (90%); viii, NaOMe, MeOH, then Rh/C, H₂, EtOH (81%); ix, HCO₂H, H₂O₂, reflux, then NaOH (5 mol dm^{–3}), THF, reflux (45%); x, Pd(OH)₂, H₂, EtOH (100%)



[†] All new compounds gave satisfactory analytical and spectral data. All optical rotations were measured at ambient temperature.

Hydrogenolysis of this material afforded for the first time pseudo- α -D-mannopyranose **2**, $[\alpha]_D + 1.9^\circ$ (c 1.0, MeOH). The structure of **2** was confirmed by ^1H NMR analysis in D_2O at 300 MHz: δ 1.60–1.72 (m, 2H, 7-H), 1.74–1.84 (m, 1H, 5-H), 3.57 (t, 1H, $J_{4,3} \approx J_{4,5}$ 9.8 Hz, 4-H), 3.61 (dd, 1H, $J_{6',5}$ 5.5, $J_{6',6}$ 11.3 Hz, 6'-H), 3.67 (dd, 1H, $J_{3,2}$ 3.4 Hz, 3-H), 3.69 (dd, 1H, $J_{6,5}$ 3.8 Hz, 6-H), 3.90 (bt, 1H, $J_{2,1}$ 3.2 Hz, 2-H), 3.98 (q, 1H, $J_{1,7a} \approx J_{1,7e}$ 3.2 Hz, 1-H).

It is noteworthy that none of the original chiral centres in quinic acid are retained in the target molecules. This feature indicates the degree of steric control which can be achieved at each transformation.

We thank the Chinese University of Hong Kong for financial support (to Y. C.) and the University of Manchester for a University Award (to Y. T.).

Received, 13th February 1991; Com. 1/00697E

References

- 1 S. Ogawa, *J. Synth. Org. Chem. Jpn.*, 1985, **43**, 26.
- 2 I. Miwa, H. Hara, J. Okuda, T. Suami and S. Ogawa, *Biochem. Int.*, 1985, **11**, 809.
- 3 C. S. Wilcox and J. J. Gaudino, *J. Am. Chem. Soc.*, 1986, **108**, 3102.
- 4 S. Ogawa, Y. Uematsu, S. Yoshida, N. Sasaki and T. Suami, *J. Carbohydr. Chem.*, 1987, **6**, 471.
- 5 R. Blattner and R. J. Ferrier, *J. Chem. Soc., Chem. Commun.*, 1987, 1008.
- 6 H. Paulsen and W. Deyn, *Liebigs. Ann. Chem.*, 1987, 125; K. Tadano, H. Maeda, M. Hoshino, Y. Iimura and T. Suami, *J. Org. Chem.*, 1987, **52**, 1946; H. Paulsen, W. Deyn, and W. Roben, *Liebigs Ann. Chem.*, 1984, 433.
- 7 S. Ogawa, T. Toyokuni, T. Konoh, Y. Hattori, S. Iwasaki, M. Suetsugu and T. Suami, *Bull. Chem. Soc. Jpn.*, 1981, **54**, 1739.
- 8 T. K. M. Shing and Y. Tang, *J. Chem. Soc., Chem. Commun.*, 1990, 312; *Tetrahedron*, 1990, 6575.
- 9 T. K. M. Shing and Y. Tang, *J. Chem. Soc., Chem. Commun.*, 1990, 748.
- 10 E. J. Corey, F. A. Carey and R. A. E. Winter, *J. Am. Chem. Soc.*, 1965, **87**, 934.
- 11 H. Adkins and A. K. Roebuck, *J. Am. Chem. Soc.*, 1948, **70**, 4041.
- 12 D. Swern, *Org. React.*, 1952, **7**, 378.