

CHEMISTRY

AN **ASIAN** JOURNAL

www.chemasianj.org

Accepted Article

Title: A Photostable AIEgen for Specific and Real-time Monitoring of Lysosomal Processes

Authors: Zhiming Wang, Weijie Zhang, Fan Zhou, Jian Du, Zujin Zhao, Anjun Qin, and Ben Zhong Tang

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Chem. Asian J.* 10.1002/asia.201801676

Link to VoR: <http://dx.doi.org/10.1002/asia.201801676>

A Journal of



A sister journal of *Angewandte Chemie*
and *Chemistry – A European Journal*

WILEY-VCH

COMMUNICATION

We have previously reported on lipid droplets-targeting DPAS (2-(((diphenylmethylene)hydrazono)-methyl)-phenol), which has excellent biocompatibility and monitoring dynamics.^[16] In this work, we designed and synthesized a novel AIE fluorogen, namely 2M-DPAS, based on DPAS (Figure 1A), and its synthetic route and analysis data are shown in Scheme S1 and Figure S1, S2. The chain length of six carbons ensure probe flexibility and the two basic morpholine groups allow 2M-DPAS to easily target lysosomes which contain many acid hydrolases. As discussed further below, in our investigation of the potential of 2M-DPAS in monitoring the dynamic changes of lysosomes with autophagy, mitophagy and phagocytosis of macromolecules, 2M-DPAS consistently showed excellent performance.




Figure 2. (A) Bright field and (B) fluorescent images of HeLa cells stained with 2M-DPAS (10 μ M, 30 min) or (C) Lysotracker Red (75 nM, 30 min). (D) A superimposed image from (B) and (C). Scale bar = 30 μ m.

The photophysical properties of 2M-DPAS were studied first. The absorption band of 2M-DPAS peaked at 415 nm in THF solution (Figure S3) while its emission peak location was observed at 550 nm (Figure 1B). Thanks for its classical ESIPT emission process, the Stokes-shift of 2M-DPAS reaches to 135 nm, which is benefit to high quality imaging. In fact, 2M-DPAS emitted faintly in pure THF solutions, and its emission remained low even the solution contained a large amount of water (50 vol%). However, with further increase of the water fraction, 2M-DPAS fluoresced more strongly at 550 nm with a light yellow color, a clear property of AIE, thus avoiding the ACQ problem with most of the conventional fluorescent probes of lysosomes. Obviously, the weak emission of 2M-DPAS in pure THF solutions was due to non-radiative decay caused by dynamic intramolecular motion, while in the aggregated state, the ESIPT process was activated and the strong emission from keto-form was shown due to the restricted intramolecular hydrogen bond motion.

With DPAS specifically targeting lipid droplets, the specificity of 2M-DPAS in cells was explored as well. We speculated that the basicity of the two morpholine groups would enable 2M-DPAS to accumulate in the acidic environments of lysosomes (pH 4.5–5.0). Thus, the well-known commercial lysosome-targeting dye Lysotracker Red (LTR) was co-stained with 2M-DPAS. As shown in Figure 2, green fluorescence from 2M-DPAS and red

fluorescence from LTR are clearly visualized, respectively. The merged image shows that the distribution of 2M-DPAS in cells is totally consistent with that of LTR (Figure 2D). The Pearson's correlation coefficient, which describes the degree of linear dependence between two variables, is 0.96, suggesting that 2M-DPAS specifically targets lysosomes. Additionally, 2M-DPAS showed a higher contrast than LTR under the same incubation time, which can be attributed to the bright emission and high selectivity of 2M-DPAS. Furthermore, the PL intensity of 2M-DPAS was tested in different pH buffered solutions. Besides, there was not a larger shift in emission peak location in aqueous solution and in lysosomes of HeLa cells, implying its pH-independent properties (Figure S4).




Figure 3. (A) Signal loss (%) of fluorescent emission of 2M-DPAS (red circles) and LTR (black squares) with increasing number of scans. (B) Cell viability of 2M-DPAS evaluated on HeLa cells by MTT assay.




Figure 4. (A–C) Bright field and (D–F) fluorescent images of HeLa cells stained with 10 μ M 2M-DPAS: (A, D) before treatment, (B, E) after treatment of rapamycin and (C, F) after treatment of rapamycin and 3-MA. Scale bar = 30 μ m.

Besides selectivity, photostability is considered one of the key factors in evaluating the performance of fluorescent imaging probes. The photostability of HeLa cells co-stained with 2M-DPAS and LTR was investigated by monitoring the fluorescence intensity before and after 30 consecutive scans by a confocal microscopy. As shown in Figure 3A, the signal loss of 2M-DPAS was less than 15% of the original intensity after 30 scans. In contrast, the fluorescence signal loss of LTR reached up to 95%. There were no significant differences observed between the first and the 30th fluorescence images of 2M-DPAS, whereas no fluorescence was observed in the 30th fluorescence image of LTR (Figure S5). The main reason was the difference in working concentrations of different fluorescent probes for lysosomal imaging due to their own properties. Truly, the working

COMMUNICATION

concentration of 2M-DPAS is up to 133 times that of LTR, resulting in higher brightness and photostability of 2M-DPAS in lysosomal imaging. If the LTR reached this concentration, the fluorescence intensity would decrease greatly because its ACQ effect was activated. Thanks for the AIE nature of 2M-DPAS, the excellent photostability was realized, and the long-term lysosomal imaging could become possible.

In addition, the cytotoxicity of 2M-DPAS on HeLa cells was assessed using MTT assay to confirm the safety of 2M-DPAS in visualizing lysosomes. As illustrated in Figure 3B, when the cells were incubated with up to 20 mM of 2M-DPAS for 24 h, the cell viability remained above 80%. Even at concentration as high as 30 mM, the cell growth was not greatly impaired, indicating a weak effect of 2M-DPAS on cell growth. The excellent biocompatibility and photostability of 2M-DPAS therefore renders it a useful fluorescent probe for dynamically monitoring lysosomes during different physiological processes.




Figure 5. Fluorescent images of HeLa cells at different layers stained with 10 μM 2M-DPAS and 50 nM MTR (A-F) before treatment of rapamycin; Fluorescent images of HeLa cells at different layers stained with 10 μM 2M-DPAS and 50 nM MTR (G-L) after treatment of rapamycin. Scale bar = 20 μm .

Autophagy is a process that transports damaged cells, degenerated or aging proteins, and organelles to the lysosomes for digestion and degradation. Visualizing and tracking activities of lysosomes enrich the insight into the process of autophagy given the close association of lysosomes and autophagy. Considering that 2M-DPAS has high specificity for lysosomes and strong photostability, the ability of real-time monitoring of dynamic changes of autophagy was tested. Since mammalian target of

rapamycin (mTOR) is an important suppressor of autophagy functioning upstream of the autophagy protein,^[17] inhibition of mTOR potentially up-regulates autophagy during the cellular response to the specific mTOR inhibitor, rapamycin.^[18] As shown in Figure 4 (D–F), the lysosomes of HeLa cells can be clearly visualized, with each green spot representing a lysosome under the fluorescence microscope. First, we found that the lysosomes increased in quantity and were widely distributed intracellularly after HeLa cells were treated with rapamycin (50 $\mu\text{g}/\text{mL}$), suggesting that the process of autophagy was monitored in real time (Figure 4E). Meanwhile, we also used 3-MA, a well characterized inhibitor of the early stage of autophagy.^[19] As illustrated in Figure 4F, no obvious changes were observed in the quantity and spatial distribution of lysosomes 30 min after simultaneously adding 3-MA (10 mM) and rapamycin (50 $\mu\text{g}/\text{mL}$) into the culture medium of 2M-DPAS stained HeLa cells. This resulted from the addition of 3-MA to the HeLa cells, inhibiting the initial stage of autophagy despite the presence of rapamycin. Therefore, 2M-DPAS can accurately monitor the process of autophagy in real time, which, along with its high selectivity for lysosomes and excellent photostability, make it an excellent choice for investigating the process of autophagy.




Figure 6. Fluorescent images of HMCs after treatment of 2 μM FAM-A β for 1 h (A) or 24 h (D), followed by 10 μM 2M-DPAS for 30 min (B, E). Merged fluorescent images of HMCs treatment and 2M-DPAS (C, F); Fluorescent images of HMCs after treatment of 2 μM FAM-A β for 1 h (G) or 24 h (J), followed by 50 nM LTR for 30 min (H, K) Merged fluorescent images of HMCs treatment and LTR (I, L). The yellow arrows indicate the locations of FAM-A β . Scale bar = 10 μm .

Autophagy is a non-selective process whereby mitochondria are the targets of autophagy. Therefore, mitophagy can also be triggered by rapamycin.^[20] As shown in Figure 5, different layers of mitochondria could be clearly captured by confocal microscopy

COMMUNICATION

in HeLa cells co-stained with 2M-DPAS and MitoTracker Red (MTR) after treatment without or with rapamycin. The long filamentous structures of mitochondria were clearly shown in red from MTR while the punctate structures of lysosomes were well portrayed as green spots from 2M-DPAS. Compared with HeLa cells cultured without rapamycin (Figure 5A-F) and with rapamycin (Figure 5G-L) (50 $\mu\text{g}/\text{mL}$) to induce mitophagy, pronounced changes had taken place in the quantity and spatial distribution of lysosomes and the morphology of mitochondria (Fig. 5). The network structures of mitochondria appeared broken and punctate, and numerous lysosomes could be seen distributed widely in the three-dimensional layer of one mitochondrion. In addition, the coincidence degree between lysosomes and mitochondria increased markedly. Therefore, the process of mitophagy was effectively monitored in the presence of 2M-DPAS and MTR, further demonstrating the superiority of 2M-DPAS in monitoring the dynamics of lysosomes.

Apart from digesting metabolic organelles such as mitochondria that have been sequestered by autophagy, lysosomes also serve to digest extracellular macromolecules that have been internalized by endocytosis. We have demonstrated that 2M-DPAS can monitor the process of autophagy and mitophagy, but can 2M-DPAS monitor the process of digesting extracellular macromolecules in the lysosomes? One of the pathological hallmarks of Alzheimer's disease (AD) is the formation of senile plaques composed primarily of aggregated amyloid β peptides ($\text{A}\beta$) in the brain. The $\text{A}\beta$ aggregation is widely considered to have a causal role in the development of AD.¹⁸ Therefore, the ability of 2M-DPAS to monitor lysosomal phagocytosis of macromolecules was tested using HMCs (human meningeal cells) and FAM- $\text{A}\beta$ (FAM-labelled $\text{A}\beta$). As shown in Figure 6 (A-C, G-I), 1 h following FAM- $\text{A}\beta$ treatment, there was no overlap between the green light emitted by FAM- $\text{A}\beta$ and the red light from lysosomes labeled with 2M-DPAS or LTR, indicating that the passage of FAM- $\text{A}\beta$ into lysosomes via endocytosis had not occurred. However, in Figure 6 (D-F, J-L), 24 h after HMCs were treated with FAM- $\text{A}\beta$, several yellow spots were observed where lysosomes were labeled with 2M-DPAS or LTR, suggesting that FAM- $\text{A}\beta$ had been phagocytosed by lysosomes labeled with 2M-DPAS or LTR. Therefore, 2M-DPAS can also monitor the lysosomal phagocytosis of macromolecules, while showing an advantage over LTR which requires a long time of laser irradiation. 2M-DPAS may be used to study the lysosomal processes in AD given the phagocytotic role of lysosomes in AD.

In summary, we have reported a novel fluorescent probe 2M-DPAS with AIE characteristics for specific lysosomal imaging. Its good biocompatibility, high selectivity, bright emission, pH-independence and excellent photostability enable continuous monitoring of the dynamic changes of lysosomes in real time. This probe has been demonstrated to effectively monitor and track the processes of autophagy, mitophagy and phagocytosis of macromolecules in lysosomes in real time. Importantly, 2M-DPAS could be a promising tool in the study of lysosome-related processes in Alzheimer's disease.

Acknowledgements

This work is financially supported by National Natural Science Foundation of China (21788102, 51673118, 81472401 and 51603127), Science & Technology Program of Guangzhou

(201804010218), Science and Technology Plan of Shenzhen (JCYJ20160428150429072 and JCYJ20160229205601482), the Innovation and Technology Commission of Hong Kong (ITC-CNERC14S01), and the Fundamental Research Funds for the Central Universities (2015ZY013, 2017JQ013 and 2017B0036).

Conflicts of interests

There are no conflicts to declare.

Keywords: Lysosomes • Aggregation-induced emission • Monitoring • Autophagy • Phagocytosis of macromolecules

- [1] a) F. Appelmans, R. Wattiaux, C. De Duve, *Biochem. J.*, **1955**, *59*, 438-447; b) C. de Duve, *Amer. Physiol. Soc.*, **1959**.
- [2] P. Saftig, B. Schroder, J. Blanz, *Biochem. Soc. Trans.*, **2010**, *38*, 1420-1423.
- [3] T. Lübke, P. Lobel and D. E. Sleat, *Biochimica et Biophysica (BBA)-Mol Cell Res.*, **2009**, *1793*, 625-635.
- [4] a) D. F. Bainton, *J. Cell Biol.*, **1981**, *91*, 66s-76s; b) K. Okamoto, *J. Cell Biol.*, **2014**, *205*, 435-445.
- [5] a) V. I. Korolchuk, S. Saiki, M. Lichtenberg, F. H. Siddiqi, E. A. Roberts, S. Imarisio, L. Jahreiss, S. Sarkar, M. Futter, F. M. Menzies, C. J. O'Kane, V. Deretic, D. C. Rubinsztein. *Nat Cell Biol.*, **2011**, *13*, 453-460; b) T. Kirkegaard, M. Jäättelä, *BBA-Mol Cell Res.*, **2009**, *1793*, 746-754.
- [6] a) Q. Wan, S. Chen, W. Shi, L. Li, H. Ma, *Angew Chem. Int. Ed.*, **2014**, *53*, 11096-11100; b) X. F. Zhang, T. Zhang, S. L. Shen, J. Y. Miao, B. X. Zhao, *J Mater. Chem. B*, **2015**, *3*, 3260-3266; c) W. J. Wang, P. Ning, Q. Wang, W. Zhang, J. Jiang, Y. Feng, X. M. Meng, *J. Mater. Chem. B*, **2018**, *6*, 1764-1770.
- [7] N. B. Finter, *J. Gen. Virol.*, **1969**, *5*, 419-427.
- [8] A. C. Allison, M. R. Young, *Lysosomes in biology and pathology*, Vol. 2, Elsevier, p. 600-628.
- [9] a) C. Svendsen, D. J. Spurgeon, P. K. Hankard, *Ecotoxicol. Environ. Saf.*, **2004**, *57*, 20-29; b) A. Pierzyńska-Mach, P. A. Janowski, J. W. Dobrucki, *Cytom. Part A*, **2014**, *85*, 729-737.
- [10] Z. Li, Y. Song, Y. Yang, L. Yang, X. Huang, J. Han, S. Han, *Chem. Sci.*, **2012**, *3*, 2941-2948.
- [11] a) J. C. Jiang, X. H. Tian, C. Z. Xu, S. X. Wang, Y. Feng, M. Chen, H. Z. Yu, M. Z. Zhu, X. M. Meng, *Chem. Commun.*, **2017**, *53*, 3645-3648; b) L. L. Hou, P. Ning, Y. Feng, Y. Q. Ding, L. Bai, L. Li, H. Z. Yu, X. M. Meng, *Anal. Chem.*, **2018**, *90*, 7122-7126; c) W. J. Wang, P. Ning, Q. Wang, W. Zhang, J. Jiang, Y. Feng, X. M. Meng, *J. Mater. Chem. B*, **2018**, *6*, 1764-1770.
- [12] a) J. Luo, Z. Xie, J. W. Y. Lam, L. Cheng, H. Chen, C. Qiu, H. S. Kwok, X. Zhan, Y. Liu, D. Zhu, B. Z. Tang, *Chem. Commun.*, **2001**, *18*, 1740-1741; b) Y. Hong, J. W. Y. Lam, B. Z. Tang, *Chem. Soc. Rev.*, **2011**, *40*, 5361-5388; c) J. Mei, Y. Hong, J. W. Y. Lam, A. Qin, Y. Tang and B. Z. Tang, *Adv. Mater.*, **2014**, *26*, 5429-5479; d) J. Mei, N. L. C. Leung, R. T. K. Kwok, J. Y. W. Lam and B. Z. Tang, *Chem. Rev.*, **2015**, *115*, 11718-11940.
- [13] a) W. Qin, P. Zhang, H. Li, J. W. Y. Lam, Y. Cai, R. T. K. Kwok, J. Qian, W. Zheng, B. Z. Tang, *Chem. Sci.*, **2018**, *9*, 2705-2710; b) M. T. Gabr, F. C. Pigge, *Dalton T.*, **2018**, *47*, 2079-2085.
- [14] Y. Hong, J. W. Lam, B. Z. Tang, *Aggregation-induced emission*, John Wiley & Sons Inc, 2014.
- [15] a) W. Chen, C. Gao, X. Liu, F. Liu, F. Wang, L. J. Tang, J. H. Jiang, *anal. Chem.*, **2018**, *90*, 8736-8741; b) M. Grossi, M. Morgunova, S. Cheung, D. Scholz, E. Conroy, M. Terrile, A. Panarella, J. C. Simpson, W. M. Gallagher, D. F. O'Shea. *Nat. Commun.*, **2016**, *7*, 10855-10868; c) W. C. Zhu, X. J. Zheng, Y. Huang, Z. Y. Lu, H. Ai, *Sci. China Chem.*, **2018**, *61*, 483-489; d) Y. B. Zhou, H. X. Liu, N. Zhao, Z. M. Wang, M. Z. Michael, N. Xie, B. Z. Tang, Y. H. Tang, *Sci. China Chem.*, **2018**, *61*, 892-897; e) K. P. Prasad, A. Than, N. Li, M. A. SK, H. W. Duan, K. Y. Pu, X. T. Zheng, P. Chen, *Mater. Chem. Front.*, **2017**, *1*, 152-157; f) C. W. T. Leung, Z. M. Wang, E. G. Zhao, Y. N. Hong, S. J. Chen, R. T. K. Kwok, A. C. S. Leung, R. We, B. S. Li, J. W. Y. Lam, B. Z. Tang, *Adv. Healthcare Mater.*, **2016**, *5*, 427-431; g) M. Gao, Q. L. Hu, G. X. Feng, B. Z. Tang, B. Liu, *J. Mater. Chem. B*, **2014**, *2*, 3438-3442; h) F. Hu, B. Liu, *Org. Biomol. Chem.*, **2016**, *14*, 9931-9944.
- [16] Z. Wang, C. Gui, E. Zhao, J. Wang, X. Li, A. Qin, Z. Zhao, Z. Yu and B. Tang, *ACS Appl. Mater. Interfaces*, **2016**, *8*, 10193-10200.
- [17] S. Wullschlegler, R. Loewith, M. N. Hall, *Cell*, **2006**, *124*, 471-484

COMMUNICATION

-
- [18] E. F. C. Blommaart, J. J. F. P. Luiken, P. J. E. Blommaart, G. M. van Woerkom, A. J. Meijer, *J. Biol. Chem.*, **1995**, *270*, 2320-2326.
- [19] P. O. Seglen, P. B. Gordon, *Proceedings of the National Academy of Sciences*, **1982**, *79*, 1889-1892.
- [20] J. Hardy and D. J. Selkoe, *Science*, **2002**, *297*, 353-356.


WILEY-VCH

Accepted Manuscript

COMMUNICATION

COMMUNICATION

2M-DPAS, as a specific probe for lysosome, can be used to continuously monitor the dynamic changes of lysosomes including autophagy and mitophagy, as well as to track the process of endocytosis of macromolecules in lysosomes with significant advantages of good biocompatibility, high selectivity, bright emission and excellent photostability.



Weijie Zhang,^{#[a, b, d]} Fan Zhou,^{#[c]}
Zhiming Wang,^[a, c] Zujin Zhao,^[c] Anjun
Qin,^[c] and Ben Zhong Tang^{#[a, c, d]}

Page No. – Page No.

Photostable AIEgen for Specific and
Real-time Monitoring of Lysosomal
Processes