


## Atom-transfer Reaction of Difluoroiodoacetate

Osamu KITAGAWA, Akihiko MIURA, Yoshiro KOBAYASHI, and Takeo TAGUCHI\*  
Tokyo College of Pharmacy, 1432-1 Horinouchi, Hachioji, Tokyo 192-03

Radical chain reaction of difluoroiodoacetate with olefins catalyzed by copper provided atom-transfer products in good yields.

Fluorinated organic compounds are attracting interest due to the characteristic features of fluorine atom, particularly in the field of medicinal chemistry and material science,<sup>1)</sup> and exploration of efficient synthetic methods has been widely required. We reported a new method for the preparation of 2,2-difluoroesters through a coupling reaction of methoxycarbonyldifluoromethylcopper (4), formed from methyl difluoroiodoacetate 1 and copper in aprotic polar solvent (DMF, DMSO or HMPA), with organic halides.<sup>2)</sup> As a part of our ongoing search for new synthetic reactions using the iodide 1 in organo-fluorine chemistry, we became interested to know whether the expected intermediacy radical can be synthetically usable in atom-transfer reaction with olefins. In this paper, we report atom-transfer reaction of methyl difluoroiodoacetate (1) with olefins 2 catalyzed by copper giving good yields of the adduct 3, which provides a useful process for C-C bond formation of difluoroacetate.<sup>3)</sup>



First, we examined the reaction of the iodide 1 with the terminal olefin, methyl 9-dodecenoate (2a) in the presence of copper powder<sup>4)</sup> in various solvent monitoring the reaction mixture by  $^{19}\text{F}$ -NMR (Table 1). In DMF, reaction proceeded at room temperature

and only addition product **3a** was identified and isolated in high yield (entry 1), while in HMPA a competitive formation of the copper reagent **4** to some extent was observed (entry 2).<sup>5)</sup> In nonpolar solvents such as diethyl ether or benzene, **3a** was obtained under refluxing condition in slightly lower yields than that in DMF solvent. The reaction proceeded by a catalytic amount of copper to give **3a** in 88% yield (entry 3), while the use of a slightly excess amount of the iodide **1** and copper resulted in nearly quantitative formation of **3a** (entry 4). From these results, it is concluded that single electron transfer (SET) from copper to the iodide **1** occurs under mild condition and resultant radical reacts with olefin to give the atom-transfer product in high yield, when DMF is used as the solvent.<sup>6)</sup>




Table 1. Solvent Effect on Cu-Catalyzed Atom-Transfer Reaction

| Entry | Solvent | <b>1</b> : <b>2a</b> : Cu | Temp    | Time/h | <b>3</b> Yield/% <sup>a)</sup> | <b>4</b> Yield/% <sup>b)</sup> |
|-------|---------|---------------------------|---------|--------|--------------------------------|--------------------------------|
| 1     | DMF     | 1 : 1 : 1                 | rt,     | 6      | 85                             | — <sup>c)</sup>                |
| 2     | HMPA    | 1 : 1 : 1                 | rt,     | 4      | 75                             | 13 <sup>d)</sup>               |
| 3     | DMF     | 1.3 : 1 : 0.3             | rt,     | 4      | 88                             | —                              |
| 4     | DMF     | 1.5 : 1 : 1.5             | rt,     | 6      | 96                             | —                              |
| 5     | benzene | 1.5 : 1 : 1.5             | reflux, | 24     | 79                             | —                              |
| 6     | ether   | 1.5 : 1 : 1.5             | reflux, | 20     | 87                             | —                              |












a) Isolated yield.      b) Determined by <sup>19</sup>F-NMR based on internal benzotrifluoride.

c) Recovery of **2a**, 7%.    d) Recovery of **2a**, 18%.

Reaction of the iodide **1** with various olefins catalyzed by copper in DMF are summarized in Table 2. It is noteworthy that 1) reaction proceeds under mild conditions; 2) atom-transfer products are obtained in good yield not only with terminal olefins (entries 1-3), but also with internal ones (entries 4, 5); 3) olefins having functional group such as acetal, ester or amide can be applicable; 4) formation of the tetrahydrofuran derivative by the reaction of the iodide **1** with diallyl ether also supports that the present reaction involves a radical chain mechanism (entry 2).<sup>7)</sup>


Since the atom-transfer product obtained by the reaction of **1** with vinylglycine derivative<sup>8)</sup> was labile on silica gel, the crude adduct was further treated with Zn in AcOH-Et<sub>2</sub>O to give 5,5-difluorohomoglutamic acid derivative (entry 6).

Table 2. Cu-Catalyzed Reaction of Difluoroiodoacetate **1** with Olefins<sup>a)</sup>

| Entry | Olefin <b>2</b>                                                                    | Time /h | Product <b>3</b>                                                                    | Yield/% <sup>b)</sup> |
|-------|------------------------------------------------------------------------------------|---------|-------------------------------------------------------------------------------------|-----------------------|
| 1     |   | 13      |   | 58                    |
| 2     |   | 15      |   | 72                    |
| 3     |   | 5       |   | 72                    |
| 4     |   | 6       |   | 79 <sup>c)</sup>      |
| 5     |   | 7.5     |   | 69                    |
| 6     |  | 20      |  | 42 <sup>d)</sup>      |

a) Solvent, DMF. b) Isolated Yield. c) trans:cis = 5:3 d) see Text.

Copper catalyzed reaction proceeded favorably with electron-rich olefins as described above, but with electron-deficient olefin such as acrylamide we obtained the iodine-free adduct **6** in 5% yield. When the reaction was conducted using Sn in THF, the adduct **6** was obtained in 40% yield.<sup>9)</sup>




In conclusion, atom-transfer reaction of difluoroiodoacetate **1** with olefins described here provides an efficient method for the preparation of functionalized molecules containing the difluoromethylene group.

We are grateful to Asahi Glass Co., for providing the iodide **1**.<sup>2)</sup>

#### References

- 1) R. Filler and Y. Kobayashi, "Biomedical Aspects of Fluorine Chemistry," Elsevier Biomedical Press and Kodansha Ltd., (1982); J. T. Welch, *Tetrahedron*, **43**, 3123 (1987).

- 2) T. Taguchi, O. Kitagawa, T. Morikawa, T. Nishiwaki, H. Uehara, H. Endo, and Y. Kobayashi, *Tetrahedron Lett.*, **27**, 6103 (1986); O. Kitagawa, T. Taguchi, and Y. Kobayashi, *Chem. Lett.*, **1989**, 389.
- 3) A part of this work was reported at the 109th Annual Meeting of the Pharmaceutical Society of Japan, Nagoya, Japan, April 1989.
- 4) R. Q. Brewster and T. Groening, *Org. Synth.*, Coll. Vol. II, 455(1948).
- 5) Q-Y. Chen, Z-Y. Yang, and Y-B. He, *J. Fluorine Chem.*, **37**, 171 (1987).
- 6) It is noteworthy that the reaction of the copper reagent **4**<sup>2)</sup> formed in DMF with **2a** proceeded very slowly to give the iodine-free olefinic compound **5** in 26% yield along with the recovery of the starting olefin **2a**. Moreover, the copper reagent **4** did not react with  $\alpha,\beta$ -unsaturated carbonyl compounds.



- 7) Atom-transfer reactions of fluoroalkyl iodide with olefins were reported. For example: N. O. Brace, *J. Org. Chem.*, **31**, 2879 (1966); M. Kuroboshi and T. Ishihara, *J. Fluorine Chem.*, **39**, 299 (1988); Q-Y. Chen, Z-Y. Yang, C-X. Zhao, and Z-M. Qui, *J. Chem. Soc., Perkin Trans. 1*, **1988**, 563 and references cited therein; Z-Y. Yang and D. J. Burton, *J. Fluorine Chem.*, **45**, 4817 (1989).
- 8) A. Afzali-Ardakani and H. Rapoport, *J. Org. Chem.*, **45**, 4817 (1980).
- 9) It was found that reaction of the iodide **1** with Sn in THF produced a tin compounds as a major product (80% yield), whose structure was tentatively assigned as  $\text{I}_2\text{Sn}(\text{CF}_2\text{COOMe})_2$  from its NMR spectra and MS spectrum, [ $^{19}\text{F}$ -NMR ( $\text{CDCl}_3$ , benzotrifluoride as an internal standard) -37.3ppm ( $^2J_{\text{F-117Sn}}=426$ ,  $^2J_{\text{F-119Sn}}=448\text{Hz}$ ),  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ )  $\delta$  164.5 (t,  $^2J_{\text{C-F}}=24\text{Hz}$ , CO), 124.2 (t,  $^1J_{\text{C-F}}=308\text{Hz}$ ,  $\text{CF}_2$ ), 54.3 (s,  $\text{CH}_3$ )] along with  $\text{Sn}(\text{CF}_2\text{COOMe})_4$ . These tin compounds did not react with acrylamide.
- 10) Aldol and related reactions of zinc reagent and 2,2-difluoroketene silyl acetal generated from the iodide **1** were reported by us. O. Kitagawa, T. Taguchi, and Y. Kobayashi, *Tetrahedron Lett.*, **29**, 1803 (1988); T. Taguchi, O. Kitagawa, Y. Suda, S. Ohkawa, A. Hashimoto, Y. Iitaka, and Y. Kobayashi, *ibid.*, **29**, 5291 (1988).

(Received April 4, 1990)