

CONTRIBUTION FROM THE NATIONAL CHEMICAL RESEARCH LABORATORY,
COUNCIL FOR SCIENTIFIC AND INDUSTRIAL RESEARCH, PRETORIA, REPUBLIC OF SOUTH AFRICA

Some Reactions of Compounds of the Type [(diene)RhX]₂ (diene = cycloocta-1,5-diene, bicyclo[2.2.1]hepta-2,5-diene; X = Cl, Br) with Tertiary Phosphines and Phosphites¹

By LINDA M. HAINES

Received November 20, 1969

The reactions of various monodentate ligands with the dimeric compounds [(diene)RhX]₂ (diene = C₈H₁₂, X = Cl, Br; diene = C₇H₈, X = Cl) are described. Whereas it has been previously established that amines and tertiary phosphines, L, cleave the halogen bridges of [(diene)RhX]₂ to form derivatives of the type (diene)RhLX, it is now found that P(OC₆H₅)₃ preferentially displaces the diene groups. Thus stepwise addition of P(OC₆H₅)₃ to [(diene)RhX]₂ in dichloromethane leads to the formation of the complexes [P(OC₆H₅)₃]₂(diene)X₂Rh₂, {Rh[P(OC₆H₅)₃]₂X}₂, and Rh[P(OC₆H₅)₃]₃X. It is also shown that halogen displacement in (diene)RhLCl by the ligands L can be effected provided that alcohols are used as solvents. Thus the reactions of [(diene)RhCl]₂ with the ligands L in methanol at room temperature and in the presence of NaB(C₆H₅)₄ afford derivatives of the type [(diene)RhL₂]B(C₆H₅)₄ [diene = C₈H₁₂, L = P(C₆H₅)₃, P(*n*-C₄H₉)₃, C₆H₁₁NH₂; diene = C₇H₈, L = P(C₆H₅)₃]. The novel ionic derivative {Rh[P(OC₆H₅)₃]₂}B(C₆H₅)₄ is similarly obtained by allowing excess P(OC₆H₅)₃ to react with [C₈H₁₂RhCl]₂ in methanol at room temperature. The preparation of the analog {Rh[P(*n*-C₄H₉)₃]₂}B(C₆H₅)₄ is also described.

Introduction

It is well-established that monodentate ligands, L, containing group V donor atoms readily react with compounds of the type [(diene)RhX]₂ [diene = C₈H₁₂ (cycloocta-1,5-diene), C₇H₈ (bicyclo[2.2.1]hepta-2,5-diene); X = Cl, Br, I] with cleavage of the halogen bridges to form the neutral derivatives (diene)RhLX.²⁻⁶ Until very recently^{7,8} further reaction of these ligands, L, with the compounds (diene)RhLX involving displacement of the halogen to yield derivatives of the type [(diene)RhL₂]X had not been observed. For instance the addition of ligand L to the compound C₈H₁₂RhLCl [L = P(C₆H₅)₃, As(C₆H₅)₃] in a number of solvents had been monitored conductometrically, but no evidence was found for the presence of any ionic species in solution.⁵

A kinetic study on the reaction of mono- and bidentate ligands with the compounds C₈H₁₂RhLCl [L = amine, P(C₆H₅)₃] using methanol as solvent gave results that were significantly different from those obtained using other solvents.⁹ In view of this anomaly it was decided to reinvestigate the reactions of the compounds [(diene)RhX]₂ with triphenylphosphine and related ligands in alcohols and other solvents. The results of this preparative study are reported here.

Experimental Section

The compounds [C₈H₁₂RhX]₂ (X = Cl, Br) and [C₇H₈RhCl]₂

were synthesized by established methods.^{2,10} All ligands were obtained commercially. The nmr spectra were recorded using a Varian A-60 instrument. Conductivities were determined by conventional methods. The elemental analyses were performed by the Bernhardt Microanalytical Laboratory, Elbach über Engelskirchen, West Germany, and by Mr. K. P. Kunz, National Chemical Research Laboratory, CSIR, Pretoria, Republic of South Africa. All yields were good varying between 60 and 80%.

Cycloocta-1,5-dienebis(triphenylphosphine)rhodium(I) Tetraphenylborate-Dichloromethane Solvate (I).—Triphenylphosphine (0.5 g, 1.9 mmol) was added to a suspension of [C₈H₁₂RhCl]₂ (0.5 g, 1 mmol) in methanol (25 ml) and the solution was stirred until complete dissolution. Addition of sodium tetraphenylboron (1.0 g, 2.9 mmol) in methanol (5 ml) resulted in the precipitation of the complex, which was crystallized from dichloromethane-pentane.

Cycloocta-1,5-dienebis(tri-*n*-butylphosphine)rhodium(I) Tetraphenylborate (II).—The precipitate obtained from the reaction of [C₈H₁₂RhCl]₂ (0.3 g, 0.6 mmol), tri-*n*-butylphosphine (0.8 ml, 4 mmol), and sodium tetraphenylboron (0.5 g, 1.5 mmol) by the method described for I was crystallized from methanol.

Bis(cyclohexylamine)cycloocta-1,5-dienerrhodium(I) Tetraphenylborate (III).—Cyclohexylamine (0.2 g, 1.8 mmol) was added to a suspension of [C₈H₁₂RhCl]₂ (0.1 g, 0.2 mmol) in methanol (20 ml) at 50° and the solution was stirred until complete dissolution. Addition of sodium tetraphenylboron (0.5 g, 1.5 mmol) in methanol (5 ml) to this solution led to the formation of yellow crystals of the required complex.

Bicyclo[2.2.1]hepta-2,5-dienebis(triphenylphosphine)rhodium(I) Tetraphenylborate (IV).—The precipitate obtained from the reaction of [C₇H₈RhCl]₂ (0.25 g, 0.5 mmol), triphenylphosphine (0.6 g, 2.3 mmol), and sodium tetraphenylboron (0.5 g, 1.5 mmol) by the method described for I was crystallized from dichloromethane-methanol.

Di-μ-chloro-cycloocta-1,5-dienebis(triphenyl phosphite)dirhodium(I) (V).—Triphenyl phosphite (0.37 g, 1.1 mmol) was added to [C₈H₁₂RhCl]₂ (0.3 g, 0.6 mmol) in dichloromethane (10 ml) and the solution was stirred for 5 min. The solvent was removed under reduced pressure and the oily residue was crystallized from dichloromethane-pentane.

Di-μ-bromo-cycloocta-1,5-dienebis(triphenyl phosphite)dirhodium(I) (VI).—The product was obtained from the reaction of

(1) A preliminary communication of this work has been presented: L. M. Haines, *Inorg. Nucl. Chem. Letters*, **5**, 399 (1969).

(2) J. Chatt and L. M. Venanzi, *J. Chem. Soc.*, 4735 (1957).

(3) M. A. Bennett and G. Wilkinson, *ibid.*, 1418 (1961).

(4) L. Cattalini, R. Ugo, and A. Orto, *J. Am. Chem. Soc.*, **90**, 4800 (1968).

(5) K. Vrieze, H. C. Volger, and A. P. Praat, *J. Organometal. Chem. (Amsterdam)*, **13**, 185 (1968).

(6) K. Vrieze, H. C. Volger, and A. P. Praat, *ibid.*, **15**, 195 (1968).

(7) At the same time as the preliminary account of this work appeared, Osborn and coworkers reported the preparation of the cation C₈H₁₂Rh[P(C₆H₅)₃]₂⁺, by the same method.⁸

(8) J. R. Shapley, R. R. Schrock, and J. A. Osborn, *J. Am. Chem. Soc.*, **91**, 2816 (1969).

(9) W. Robb and L. M. Haines, unpublished results.

(10) E. W. Abel, M. A. Bennett, and G. Wilkinson, *J. Chem. Soc.*, 3178 (1959).

TABLE I
 COLOR, CONDUCTIVITY, MOLECULAR WEIGHT, AND ANALYTICAL DATA

No.	Compound	Color	Conductivity, ^a ohm ⁻¹ cm ² mol ⁻¹	Mol wt ^b		Calcd				Anal, %				Found			
				Calcd	Found	C	H	P or N	Cl or Br	C	H	P or N	Cl or Br	C	H	P or N	Cl or Br
I	{Rh(C ₆ H ₅) ₃ [P(C ₆ H ₅) ₃] ₂ }B(C ₆ H ₅) ₄ ·CH ₂ Cl ₂	Orange	89	...	...	72.7	5.7	...	6.2	72.6	5.6	...	6.0	...	...	...	...
II	{Rh(C ₆ H ₅) ₃ [P(<i>n</i> -C ₄ H ₉) ₃] ₂ }B(C ₆ H ₅) ₄	Brown	85	...	...	71.9	9.3	...	...	71.2	9.2	...	...	...	...	...	...
III	{Rh(C ₆ H ₅) ₃ (C ₆ H ₁₁ NH ₂) ₂ }B(C ₆ H ₅) ₄	Yellow	81	...	...	72.5	8.0	3.8	...	72.3	7.8	3.8	...	...	...	...	...
IV	{Rh(C ₇ H ₉) ₃ [P(C ₆ H ₅) ₃] ₂ }B(C ₆ H ₅) ₄	Red	87	...	...	77.5	5.6	...	...	77.7	5.5	...	...	...	...	...	...
V	Rh ₂ Cl ₂ (C ₆ H ₅) ₂ [P(OC ₆ H ₅) ₃] ₂	Orange	0.3	1006	976	52.6	4.2	6.2	7.1	52.5	4.2	6.1	6.9	...	...	...	...
VI	Rh ₂ Br ₂ (C ₆ H ₅) ₂ [P(OC ₆ H ₅) ₃] ₂	Orange	0.5	1095	1058	48.2	3.9	...	14.6	48.4	3.8	...	14.9	...	...	...	...
VII	Rh ₂ Cl ₂ (C ₇ H ₉) ₂ [P(OC ₆ H ₅) ₃] ₂	Yellow	0.3	...	<i>c</i>	52.2	3.9	...	...	52.1	3.9	...	...	...	...	...	...
VIII	{RhCl[P(OC ₆ H ₅) ₃] ₂ } ₂	Yellow	0.2	1518	1368 ^d	57.0	4.0	...	4.8	56.9	3.9	...	5.0	...	...	...	...
IX	{RhBr[P(OC ₆ H ₅) ₃] ₂ } ₂	Yellow	0.4	1607	1367 ^d	53.8	3.8	...	10.0	53.8	3.9	...	10.1	...	...	...	...
X	RhCl[P(OC ₆ H ₅) ₃] ₃	Yellow	0.2	1069	1070	60.7	4.3	...	3.3	60.2	4.1	...	4.6	...	...	...	...
XI	RhBr[P(OC ₆ H ₅) ₃] ₃	Yellow	0.3	1114	1104	58.2	4.1	...	7.2	58.0	4.2	...	7.3	...	...	...	...
XII	Rh[P(OC ₆ H ₅) ₃] ₂ (CH ₃ COCHCOCH ₃)	Pale yellow	0.4	...	<i>c</i>	59.9	4.5	7.5	...	59.8	4.5	7.6	...	...	...	...	...
XIII	{Rh[P(OC ₆ H ₅) ₃] ₄ }B(C ₆ H ₅) ₄	Yellow	74	...	...	69.3	4.9	...	...	68.8	4.8	...	...	...	...	...	...
XIV	{Rh[P(OC ₆ H ₅) ₃] ₄ }ClO ₄	Yellow	80	...	...	59.9	4.2	...	2.5	59.7	4.2	...	2.3	...	...	...	...
XV	{Rh[P(<i>n</i> -C ₄ H ₉) ₃] ₄ }B(C ₆ H ₅) ₄	Brown	94	...	...	70.2	10.5	10.1	...	70.0	10.4	10.0	...	...	...	...	...

^a Solutions are ~10⁻³ M in acetone. ^b Solutions are ~2 × 10⁻² M in benzene. ^c Not measured. ^d Decomposition.

triphenyl phosphite (0.32 g, 1.1 mmol) and [C₆H₅RhBr]₂ (0.3 g, 0.5 mmol) by the method described for V.

Di-μ-chloro-bicyclo[2.2.1]hepta-2,5-dienebis(triphenyl phosphite)dirhodium(I) (VII).—The product was obtained from the reaction of triphenyl phosphite (0.33 g, 1.1 mmol) and [C₇H₉RhCl]₂ (0.25 g, 0.5 mmol) by the method described for V.

Di-μ-chloro-tetrakis(triphenyl phosphite)dirhodium(I) (VIII).—Triphenyl phosphite (0.76 g, 2.5 mmol) was added to [C₆H₅RhCl]₂ (0.3 g, 0.6 mmol) in dichloromethane (10 ml) and the solution was stirred for 5 min. The solvent was removed under reduced pressure, and the resultant oil was washed with methanol and then crystallized from dichloromethane-methanol.

Di-μ-bromo-tetrakis(triphenyl phosphite)dirhodium(I) (IX).—The complex was obtained from the reaction of triphenyl phosphite (0.64 g, 2.1 mmol) and [C₆H₅RhBr]₂ (0.3 g, 0.5 mmol) by the method described for VIII.

Chlorotris(triphenyl phosphite)rhodium(I) (X).—Triphenyl phosphite (1.15 g, 3.7 mmol) was added to [C₆H₅RhCl]₂ (0.3 g, 0.6 mmol) in dichloromethane (10 ml) and the solution was stirred for 5 min. The solvent was removed under reduced pressure, and the oily residue was washed with methanol and then crystallized from dichloromethane-methanol.

Bromotris(triphenyl phosphite)rhodium(I) (XI).—The complex was obtained from the reaction of triphenyl phosphite (0.96 g, 3.1 mmol) and [C₆H₅RhBr]₂ (0.3 g, 0.5 mmol) by the method described for X.

2,4-Pentanedionatobis(triphenyl phosphite)rhodium(I) (XII).—A mixture of {Rh[P(OC₆H₅)₃]₂Cl}₂ (0.25 g, 0.2 mmol), anhydrous Na₂CO₃ (0.1 g, 0.9 mmol), and 2,4-pentanedione (0.1 g, 1 mmol) in dry acetone (20 ml) was stirred at room temperature for 2 hr. The acetone was removed under reduced pressure, the residue was extracted with dichloromethane, and the extract was filtered. Addition of pentane to the concentrated filtrate afforded yellow crystals of the complex.

Tetrakis(triphenyl phosphite)rhodium(I) Tetraphenylborate (XIII).—A methanol solution (20 ml) containing triphenyl phosphite (1.5 g, 4.8 mmol) and suspended [C₆H₅RhCl]₂ (0.3 g, 0.6 mmol) was stirred until complete dissolution. Addition of sodium tetraphenylboron (0.5 g, 1.5 mmol) in methanol (5 ml) to this solution led to the precipitation of the complex, which was crystallized from acetone-water.


Tetrakis(triphenyl phosphite)rhodium(I) Perchlorate (XIV).—The complex was obtained from the reaction of triphenyl phosphite (1.5 g, 4.8 mmol), [C₆H₅RhCl]₂ (0.3 g, 0.6 mmol), and lithium perchlorate trihydrate (0.4 g, 2.5 mmol) by the method described for XIII.

Tetrakis(tri-*n*-butylphosphine)rhodium(I) Tetraphenylborate

(XV).—An ethanol solution (20 ml) containing tri-*n*-butylphosphine (1.3 g, 6.4 mmol) and suspended [C₆H₅RhCl]₂ (0.2 g, 0.4 mmol) was refluxed for 3 hr. Addition of sodium tetraphenylboron (0.5 g, 0.9 mmol) in ethanol (5 ml) led to the precipitation of the complex, which was crystallized from ethanol.

Results and Discussion

The uv-visible spectra of solutions of the compound C₆H₅RhP(C₆H₅)₃Cl alone and in the presence of triphenylphosphine and lithium chloride were investigated as a preliminary to kinetic studies. When methanol was used as solvent, spectral changes were observed which could only be explained in terms of the equilibrium



No such spectral changes were found when other solvents were employed. In view of these observations, an attempt was made to isolate the derivative {C₆H₅Rh[P(C₆H₅)₃]₂}Cl from a methanol solution of the compound [C₆H₅RhCl]₂ and excess triphenylphosphine, but this was unsuccessful. However, addition of sodium tetraphenylboron to this methanol solution led to the immediate precipitation of the ionic derivative {C₆H₅Rh[P(C₆H₅)₃]₂}B(C₆H₅)₄. The complexes [C₆H₅RhL₂]B(C₆H₅)₄ [L = P(*n*-C₄H₉)₃, C₆H₁₁NH₂] were similarly prepared. Replacement of cycloocta-1,5-diene by bicyclo[2.2.1]hepta-2,5-diene had no effect on these reactions, as evidenced by the preparation of the derivative {C₇H₉Rh[P(C₆H₅)₃]₂}B(C₆H₅)₄. The chemical composition of the complexes [(diene)RhL₂]B(C₆H₅)₄ [diene = C₈H₁₂, L = P(C₆H₅)₃, P(*n*-C₄H₉)₃, C₆H₁₁NH₂; diene = C₇H₈, L = P(C₆H₅)₃] was established by elemental analysis (Table I) and from a comparison of the integrated intensities of the diene and ligand proton resonances in the nmr spectra. All of the compounds are 1:1 electrolytes in acetone and were assumed to be diamagnetic from the sharpness

TABLE II
NMR DATA AND ASSIGNMENTS^a


No.	Compound	Aromatic protons	Diene		Other protons	Assignment
			Olefinic protons	Aliphatic protons		
I	$\{\text{Rh}(\text{C}_6\text{H}_{12})[\text{P}(\text{C}_6\text{H}_5)_3]_2\}\text{B}(\text{C}_6\text{H}_5)_4 \cdot \text{CH}_2\text{Cl}_2$	2.72 m	5.51 b, s	7.78 b, s	4.77 s	CH_2Cl_2
II	$\{\text{Rh}(\text{C}_8\text{H}_{12})[\text{P}(n\text{-C}_4\text{H}_9)_3]_2\}\text{B}(\text{C}_6\text{H}_5)_4$	2.83 m	5.24 b, s	7.85 b, s	8.70 m	$\text{P}(n\text{-C}_4\text{H}_9)_3$
III	$\{\text{Rh}(\text{C}_8\text{H}_{12})(\text{C}_6\text{H}_{11}\text{NH}_2)_2\}\text{B}(\text{C}_6\text{H}_5)_4$	2.70 m	5.89 b, s		4.23 m	$-\text{NH}_2$
IV	$\{\text{Rh}(\text{C}_7\text{H}_8)[\text{P}(\text{C}_6\text{H}_5)_3]_2\}\text{B}(\text{C}_6\text{H}_5)_4$	2.70 m	5.64 b, s	6.23 m, 8.60 m	8.30 m	All aliphatic protons
V	$\text{Rh}_2\text{Cl}_2(\text{C}_8\text{H}_{12})[\text{P}(\text{OC}_6\text{H}_5)_3]_2$	2.77	6.26	8.2 m	...	
VI	$\text{Rh}_2\text{Br}_2(\text{C}_8\text{H}_{12})[\text{P}(\text{OC}_6\text{H}_5)_3]_2$	2.72 s	6.10	8.2	...	
VII	$\text{Rh}_2\text{Cl}_2(\text{C}_7\text{H}_8)[\text{P}(\text{OC}_6\text{H}_5)_3]_2$	2.73 s	6.48	8.94	...	
XII	$\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{CH}_3\text{COCHCOCH}_3)$	2.81 s			4.96 s	$\left. \begin{matrix} \text{H} \\ \text{CH}_3 \end{matrix} \right\} \text{of } \text{CH}_3\text{COCHCOCH}_3$
					8.53 s	
					8.60 m	
XV	$\{\text{Rh}[\text{P}(n\text{-C}_4\text{H}_9)_3]_4\}\text{B}(\text{C}_6\text{H}_5)_4$	2.73 m				$\text{P}(n\text{-C}_4\text{H}_9)_3$

^a τ scale; measured in CDCl_3 at 38°. Abbreviations: s, singlet; m, multiplet; b, broad.

of the nmr spectral peaks. The assignments of the nmr spectra are given in Table II.

The halogen displacement reactions by the ligands $\text{P}(\text{C}_6\text{H}_5)_3$, $\text{P}(n\text{-C}_4\text{H}_9)_3$, and $\text{C}_6\text{H}_{11}\text{NH}_2$ only occurred in methanol and ethanol and could not be effected in tetrahydrofuran, acetone, dimethylformamide, chloroform, dichloromethane, and benzene. It has been recently noted that the halogen displacement reactions involved in the preparation of some cationic carbon disulfide derivatives of rhodium and iridium, *e.g.*, $\{\text{M}(\pi\text{-CS}_2)[\text{P}(\text{C}_6\text{H}_5)_3]_3\}\text{B}(\text{C}_6\text{H}_5)_4$ ($\text{M} = \text{Rh}, \text{Ir}$),¹¹ and in the isomerism of the compound $\{\text{Ir}[\text{P}(\text{CH}_3)_2\text{C}_6\text{H}_5]_2(\text{CO})\text{Cl}_2(\sigma\text{-C}_6\text{H}_5)\}$, *via* the cation $\text{Ir}[\text{P}(\text{CH}_3)_2\text{C}_6\text{H}_5]_2(\text{CO})\text{Cl}(\pi\text{-C}_6\text{H}_5)^+$,¹² are specific to alcohols.

When tertiary phosphite ligands were used as an extension to these studies, unexpected results were obtained. Thus the reaction sequence



(diene = C_7H_8 , C_8H_{12}) was not observed for the ligand $\text{L} = \text{P}(\text{OC}_6\text{H}_5)_3$. Instead it was found that the reaction of 2 mol of triphenyl phosphite with 1 mol of the compound $[\text{C}_8\text{H}_{12}\text{RhCl}]_2$ at room temperature in dichloromethane gave a product which was characterized as the complex $[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{C}_8\text{H}_{12})\text{Cl}_2\text{Rh}_2$. The nmr spectrum of this compound showed the presence of both cycloocta-1,5-diene and triphenyl phosphite ligands, while the dimeric nature was established from molecular weight measurements. Based on this physical and spectral evidence the structure shown in Figure 1 was proposed. This has now been confirmed






Figure 1.

by an X-ray crystal analysis.¹³ The structure determination has also shown that the



moiety in the molecule $[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{C}_8\text{H}_{12})\text{Cl}_2\text{Rh}_2$ is bent about the Cl-Cl axis. A similar type of bending has been observed for the molecule $[\text{Rh}(\text{CO})_2\text{Cl}]_2$,¹⁴ but the



group in the dimer $[\text{C}_8\text{H}_{12}\text{RhCl}]_2$ ¹⁵ is planar.

One mole of the compound $[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{C}_8\text{H}_{12})\text{Cl}_2\text{Rh}_2$ was found to react with 2 mol of triphenyl phosphite to yield a product in which the sole cycloocta-1,5-diene group has been replaced by the triphenyl phosphite ligands. This product was characterized as the compound $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2\text{Cl}\}_2$. Although molecular weight measurements were unreliable, the dimeric nature of this complex was demonstrated by its reaction with various reagents. Thus with acetylacetone in basic medium the compound $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2\text{Cl}\}_2$ yielded the derivative $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{CH}_3\text{COCHCOCH}_3)$. It should be noted that the thiocyanato complex $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2\text{SCN}\}_2$ has been synthesized by treating the compound $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{CO})\text{Cl}$ with excess potassium thiocyanate.^{16,17} The addition of 2 mol of triphenyl phosphite to 1 mol of the complex $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2\text{Cl}\}_2$ resulted in the cleavage of the halogen bridges and the formation of the known compound $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_3\text{Cl}$ previously synthesized from the derivative $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{CO})\text{Cl}$.^{16,18} Similarly the stepwise addition of triphenyl phosphite to the compound $[\text{C}_8\text{H}_{12}\text{RhBr}]_2$ afforded the derivatives $[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{C}_8\text{H}_{12})\text{Br}_2\text{Rh}_2$, $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2\text{Br}\}_2$, and $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_3\text{Br}$. The reaction of the compound $[\text{C}_7\text{H}_8\text{RhCl}]_2$ with triphenyl phosphite gave a similar series of derivatives. However, addition of triphenyl

(14) L. F. Dahl, C. Martell, and D. L. Wampler, *J. Am. Chem. Soc.*, **83**, 1761 (1961).

(15) J. A. Ibers and R. G. Snyder, *Acta Cryst.*, **15**, 923 (1962).

(16) L. Vailarino, *J. Chem. Soc.*, 2473 (1957).

(17) M. A. Jennings and A. Wojcicki, *Inorg. Chem.*, **6**, 1854 (1967).

(18) L. Vailarino, *J. Chem. Soc.*, 2287 (1957).

(11) M. P. Yagupsky and G. Wilkinson, *J. Chem. Soc., A*, 2813 (1968).

(12) A. J. Deeming and B. L. Shaw, *ibid.*, **A**, 1562 (1969).

(13) J. Coetzer and G. Gafner, *Acta Cryst.*, in press.

phosphite to the compound $[\text{C}_8\text{H}_{12}\text{RhI}]_2$ gave unstable products and these were not further studied.

All of the compounds $[\text{P}(\text{OC}_6\text{H}_5)_3]_2(\text{diene})\text{X}_2\text{Rh}_2$, $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_2\text{X}\}_2$, and $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_3\text{X}$ (diene = C_8H_{12} , $\text{X} = \text{Cl}, \text{Br}$; diene = C_7H_8 , $\text{X} = \text{Cl}$) were characterized by elemental analysis (Table I) and, where appropriate, from the nmr spectral data. The nature of the species was established from molecular weight measurements. The compounds are nonelectrolytes in acetone and are diamagnetic. The nmr spectral data are given in Table II.

As all previous reports have shown that the action of monodentate ligands L on the dimeric compound $[\text{C}_8\text{H}_{12}\text{RhCl}]_2$ results in the formation of derivatives of the type $\text{C}_8\text{H}_{12}\text{RhLCl}$, it is surprising that triphenyl phosphite should, under identical experimental conditions, successively replace the cycloocta-1,5-diene groups in preference to cleaving the halogen bridges. This difference in reaction schemes between triphenyl phosphite and the other donor ligands previously studied can be associated with the softer character of triphenyl phosphite.

In view of the solvent specificity in the synthesis of the derivative $\{\text{C}_8\text{H}_{12}\text{Rh}[\text{P}(\text{C}_6\text{H}_5)_3]_2\}\text{B}(\text{C}_6\text{H}_5)_4$ from the compound $\text{C}_8\text{H}_{12}\text{RhP}(\text{C}_6\text{H}_5)_3\text{Cl}$, the possibility of displacing the halogen in the complex $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_3\text{Cl}$ by triphenyl phosphite in alcohol was investigated. It was found that the compound $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_3\text{Cl}$, normally insoluble in methanol, slowly dissolves in this solvent in the presence of excess triphenyl phosphite. A cationic species $\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_4^+$ precipitated from this solution on addition of sodium tetraphenylboron or lithium perchlorate. An alternative structure for this cation, $[\text{P}(\text{OC}_6\text{H}_5)_3]_3(\text{C}_6\text{H}_5\text{O})_2\text{POC}_6\text{H}_5\text{H}_4\text{RhH}^+$, involving a rhodium-phenyl bond similar to that recently proposed¹⁹ for the derivatives "M- $[\text{P}(\text{OC}_6\text{H}_5)_3]_4$ " (M = Rh, Ir)²⁰ was eliminated as there was no nmr or ir evidence for a hydrogen bonded to rhodium. The derivative $\{\text{Rh}[\text{P}(\text{n-C}_4\text{H}_9)_3]_4\}\text{B}(\text{C}_6\text{H}_5)_4$ has also been synthesized by allowing the com-

pound $[\text{C}_8\text{H}_{12}\text{RhCl}]_2$ to react with tri-*n*-butylphosphine in refluxing ethanol and then adding sodium tetraphenylboron. As it has already been established that tri-*n*-butylphosphine reacts at room temperature with the compound $[\text{C}_8\text{H}_{12}\text{RhCl}]_2$ in methanol to form the cation $\text{C}_8\text{H}_{12}\text{Rh}[\text{P}(\text{n-C}_4\text{H}_9)_3]_2^+$, it is apparent that this ion is an intermediate in the formation of the species $\text{Rh}[\text{P}(\text{n-C}_4\text{H}_9)_3]_4^+$. It has been previously shown that the compound $\text{Rh}[\text{P}(\text{C}_6\text{H}_5)_3]_3\text{Cl}$ dissociates in solution with loss of triphenylphosphine.^{21,22} It is, therefore, not surprising that attempts to isolate the cation $\text{Rh}[\text{P}(\text{C}_6\text{H}_5)_3]_4^+$ from the reaction of the compound $\text{Rh}[(\text{C}_6\text{H}_5)_3]_3\text{Cl}$ or $[\text{C}_8\text{H}_{12}\text{RhCl}]_2$ with excess triphenylphosphine in refluxing alcohols were unsuccessful.

The compounds $\{\text{RhL}_4\}\text{B}(\text{C}_6\text{H}_5)_4$ [L = $\text{P}(\text{n-C}_4\text{H}_9)_3$, $\text{P}(\text{OC}_6\text{H}_5)_3$] and $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_4\}\text{ClO}_4$ were characterized by elemental analysis (Table I) and nmr spectral intensity data where appropriate. The complexes are 1:1 electrolytes in acetone and are diamagnetic. The nmr data are presented in Table II. These complexes are the first examples of four-coordinate rhodium(I) derivatives which contain identical monodentate group V donor ligands. The species $\text{Rh}(\text{L}_2)_2^+$ [$\text{L}_2 = \text{R}_2\text{PCH}_2\text{CH}_2\text{PR}_2$ (R = CH_3 ,²³ C_6H_5 ²⁴), *cis*-(C_6H_5)₂AsCH=CHAs(C_6H_5)₂²⁵], which contain bidentate ligands, have been reported recently. The reactivity of the compound $\{\text{RhL}_4\}\text{B}(\text{C}_6\text{H}_5)_4$ toward hydrogen is of interest. The derivative $\{\text{Rh}[\text{P}(\text{n-C}_4\text{H}_9)_3]_4\}\text{B}(\text{C}_6\text{H}_5)_4$ reacts readily with molecular hydrogen to yield a stable complex $\{\text{Rh}[\text{P}(\text{n-C}_4\text{H}_9)_3]_4\text{H}_2\}\text{B}(\text{C}_6\text{H}_5)_4$. In contrast, the compound $\{\text{Rh}[\text{P}(\text{OC}_6\text{H}_5)_3]_4\}\text{B}(\text{C}_6\text{H}_5)_4$ adds hydrogen reversibly, the resulting hydride being stable only under an atmosphere of hydrogen. The details of these and similar oxidative addition reactions will be the subject of a further publication.

Acknowledgments.—The author wishes to thank Dr. K. G. R. Pachler for the measurement of the nmr spectra.

(21) J. A. Osborn, F. H. Jardine, J. F. Young, and G. Wilkinson, *J. Chem. Soc., A*, 1711 (1966).

(22) M. A. Bennett and P. A. Longstaff, *Chem. Ind. (London)*, 846 (1965).

(23) J. Chatt and S. A. Butter, *Chem. Commun.*, 501 (1967).

(24) A. Sacco and R. Ugo, *J. Chem. Soc.*, 3274 (1964).

(25) J. T. Mague and J. P. Mitchener, *Chem. Commun.*, 911 (1968).

(19) W. H. Knoth and R. A. Schunn, *J. Am. Chem. Soc.*, **91**, 2400 (1969).

(20) S. D. Robinson, *Chem. Commun.*, 521 (1968).