Thermal Claisen Rearrangement of Aryl Allenylmethyl Ethers: Synthesis of 2-(*o*-Hydroxyaryl)buta-1,3-dienes and 4-Methyl-2*H*-1-benzopyrans

T. Balasubramanian and K. K. Balasubramanian*

Department of Chemistry, Indian Institute of Technology, Madras 600 036, India

Thermal Claisen rearrangement of aryl allenylmethyl ethers in diethylene glycol afforded 2-(*o*-hydroxyaryl)buta-1,3-dienes and 4-methyl-2*H*-1-benzopyrans in good yield.

Few classical named reactions have been studied so extensively from synthetic, stereochemical, mechanistic and theoretical aspects as the Claisen rearrangement. The many exhaustive reviews surveying the rapid development of this reaction illustrate the attention it continues to receive.1 Though a large number of reports on the Claisen rearrangement of non-cumulated double bond systems are available in the literature,1,2 only a few examples can be found for allenes.³ In particular, the thermal behaviour of aryl allenylmethyl (buta-2,3-dienyl) ethers has remained virtually uninvestigated till recently^{4,5} in contrast to that of the isomeric aryl prop-2-ynyl ethers which has been studied extensively.^{1,6} Herein, we report a simple route to 4-methyl-2H-1-benzopyrans and an expedient synthesis of 2-(o-hydroxyaryl)buta-1,3-dienes by the thermal Claisen rearrangement of aryl allenylmethyl ethers.

Heating the 2-naphthyl allenylmethyl ether 1e (2 mmol) in diethylene glycol (DEG) (5 ml) at 160–165 °C (bath temperature) for 20 min afforded the 2-(2-hydroxy-1-naphthyl)buta-1,3-diene 2e as a colourless liquid in 82% yield.† Other aryl allenylmethyl ethers 1a–d and 1f were also found to undergo this [3,3] sigmatropic rearrangement furnishing the respective 2-(o-hydroxyaryl)buta-1,3-dienes 2a–d and 2f in good yield‡ (Scheme 1, Table 1).



Scheme 1 Conditions (in DEG under N_2): i, heat, shorter time; ii, reflux, longer time

 \dagger The duration of heating was found to be critical: after short times unreacted starting material was present to a considerable extent whereas heating for a long period led to a lower yield of **2e**.

[‡] All the new compounds reported in this communication were thoroughly characterised by spectral and HRMS data.

The 1-naphthyl allenylmethyl ether 1f rearranged faster than the isomeric ether 1e (Table 1, Entries 5 and 6). The lower yield of **2f** in the case of the 1-naphthyl allenylmethyl ether is due to its fast rearrangement and a more rapid cyclisation of the intermediate phenolic product, viz., 2f, to the naphthopyran 3f. This cyclisation can be avoided by trapping the 2-(1-hydroxy-2-naphthyl)buta-1,3-diene as its acetate 4f. Refluxing the 1-naphthyl allenylmethyl ether 1f (2 mmol) in a mixture of o-dichlorobenzene (oDCB) (5 ml) and acetic anhydride (1 ml) in the presence of a catalytic amount of 4-N, N-dimethylamino-pyridine (DMAP) for 8 h afforded the 2-(1-acetoxyl-2-naphthyl)buta-1,3-diene 4f as a colourless liquid in 80% yield§ (Scheme 2). The 2-arylbuta-1,3-dienes are useful synthons in Diels-Alder reactions and there is no good method for their synthesis.⁷ The thermal Claisen rearrangement of aryl allenylmethyl ethers thus provides a convenient method for the preparation of these compounds.

Benzopyran compounds are widespread in a variety of natural products and have interesting properties, *e.g.*, psychotropic (tetrahydrocannabinol), and insecticidal: ageratochromene (precocene), phytoalexins (pterocarpans) *etc.*⁸ Claisen rearrangement of aryl prop-2-ynyl ethers is a good method for the synthesis of benzopyrans.⁹ Surprisingly the utility of thermal Claisen rearrangement of aryl allenylmethyl ethers for the synthesis of benzopyrans has not been explored. In the present study, 4-methyl-2H-1-benzopyrans **3a-f** were



Scheme 2 Conditions: (MeCO)₂O, DMAP, *o*DCB, reflux, 8 h, under N₂

Other aryl allenylmethyl ethers were also trapped as acetates in 62-82% yield.

J. CHEM. SOC., CHEM. COMMUN., 1992

Table 1	Synthesis of	f 2-(o-h	ydroxyary	yl)buta-1,3-dienes	3 2 and 4-meth	yl-2H-1-benzopyrans 3
---------	--------------	----------	-----------	--------------------	----------------	-----------------------

1	Entry	Starting ether	R ¹	R ²	R ³	R ⁴	Conditions for 2	Yield of 2 (%)	Time at reflux for 3	Yield of 3 (%)
	1	1a	н	Н	OMe	н	200–210 °C, 12 min	78	30 min	86
	2	1b	н	Н	Me	Н	200–210 °C, 16 min	72	40 min	82
	3	1c	н	н	н	н	200-210 °C, 20 min	66	40 min	80
	4	1d	н	н	Cl	Н	200–210 °C, 20 min	62	40 min	80
:	5	1e	н	н	Benzo		160–165 °C, 20 min	82	5 min	88
(6	1f	Benzo		Н	Н	140–145 °C, 12 min	52 ^a	5 min	88

^a The rest of the product was 4-methyl-2H-1-naphthopyran and starting material.

Table 2 Comparison of the ease of rearrangement of allenylmethyl, allyl, prop-2-ynyl and but-2-ynyl ethers of 1-naphthol and 2-naphthol A^a

A								
	Entry	Compound	R ¹ in A	R ² in A	t/min	Unreacted (%)	Products	
	1	1e	Н	OCH ₂ CH=C=CH ₂	20	0	2e + 3e	
	2	5	Н	OCH ₂ CH=CH ₂	20	66	1-Allyl-2-naphthol	
	3	6	Н	OCH ₂ C≡CH	20	95	3H-Naphtho[2,1-b]pyran	
	4	7	Н	OCH ₂ C≡CMe	20	100		
	5	1f	OCH ₂ CH=C=CH ₂	Н	8	0	2f + 3f	
	6	8	OCH ₂ CH=CH ₂	Н	8	63	2-Allyl-1-naphthol	
	7	9	OCH ₂ C≡CH	Н	8	94	2H-Naphtho[1,2-b]pyran	
	8	10	OCH ₂ C≡CMe	Н	8	100		

^a In DEG at 160-165 °C.

obtained in good yield by refluxing the aryl allenylmethyl ethers 1a-f in diethylene glycol for 5–40 min (Scheme 1 and Table 1).

Since substituent effects on Claisen rearrangement have been the subject of several theoretical studies in recent years,10 it was of interest to compare the relative ease of rearrangement of allenylmethyl, allyl, prop-2-ynyl and but-2ynyl ethers of 1-naphthol as well as of 2-naphthol (Table 2). The results from our preliminary study indicate that the allenylmethyl ethers rearrange much faster than the other ethers, in both the 1-naphthyl (Table 2, Entries 5-8) and the 2-naphthyl (Table 2, Entries 1-4) systems. This is in good agreement with the calculations of Jorgensen et al.¹¹ from their theoretical studies on [3,3] sigmatropic reactions. They have shown that a terminus which is part of a cumulene lowers the E_a by 25 kJ mol⁻¹. In all these cases 1-naphthyl ethers were observed to rearrange faster than the corresponding 2-naphthyl ethers. Also, the possibility of fast prototropic rearrangement of le to the propynylic derivative 7, prior to the Claisen rearrangement step, is ruled out (Table 2, Entries 1 and 4).

Our successful thermal Claisen rearrangement of aryl allenylmethyl ethers to 2-(o-hydroxyaryl)buta-1,3-dienes after a short time and to 4-methyl-2H-1-benzopyrans after a longer time is in marked contrast to the recent report of Dikshit *et al.*⁴ who observed the thermal rearrangement of *p*-methoxyphenyl allenylmethyl ether in refluxing N, Ndiethylaniline (b.p. 217 °C) or diethylene glycol (b.p. 245 °C) led to a complex mixture. It is evident that the products had decomposed under their experimental conditions.¶

T.B.S. is grateful to CSIR, India for a fellowship. We acknowledge RSIC, IIT, Madras and Dr K. Nagarajan, Searle

(India) Ltd, Bombay for spectral data and M/s. BASF, Germany for a gift of prop-2-ynyl alcohol.

Received, 3rd August 1992; Com. 2/04163D

References

- 1 R. P. Lutz, Chem. Rev., 1984, 84, 205; F. E. Ziegler, Chem. Rev., 1988, 88, 1423; S. Blechert, Synthesis, 1989, 71.
- L. M. Harwood, A. J. Oxford and C. Thomson, J. Chem. Soc., Chem. Commun., 1991, 1303; P. A. Grieco, E. B. Brandes, S. McCann and J. D. Clark, J. Org. Chem., 1989, 54, 5849; K. Nonoshita, H. Banno, K. Maruoka and H. Yamamoto, J. Am. Chem. Soc., 1990, 112, 316.
- M. J. Sleeman and G. V. Meehan, *Tetrahedron Lett.*, 1989, **30**, 3345; J-P. Dulcere and J. K. Crandall, *J. Chem. Soc.*, *Chem. Commun.*, 1990, 561; M. Ochiai, T. Ito, Y. Takaoka and Y. Masaki, *J. Am. Chem. Soc.*, 1991, **113**, 1319.
- 4 D. K. Dikshit, S. Singh and S. K. Panday, J. Chem. Res. (S), 1991, 298; For Hg(OCOCF₃)₂ catalysed transformation of aryl allenic ethers, see-T. Balasubramanian and K. K. Balasubramanian, Tetrahedron Lett., 1991, 32, 6641.
- 5 Part of this work was reported by us at the VIIIth IUPAC Symposium on Organic synthesis held at University of Helsinki, Finland, 23–27th July 1990.
- 6 W. K. Anderson and E. J. LaVoie, J. Org. Chem., 1973, 38, 3832;
 I. Iwai and J. Ide, Chem. Pharm. Bull., 1962, 10, 926.
- 7 C.-T. Lin and W.-J. Hsu, Can. J. Chem., 1989, 67, 2153; S. W. Baldwin, P. Greenspan, C. Alaimo and A. T. McPhail, *Tetrahedron Lett.*, 1991, 32, 5877.
- 8 G. P. Ellis, Chromenes, Chromanones and Chromones, Wiley, New York, 1977; G. P. Ellis and I. M. Lockhart, Chromans and Tocopherols, Wiley, New York, 1981.
- 9 J. D. Hepworth, in *Comprehensive Heterocyclic Chemistry*, ed. A. R. Katritzky, C. W. Rees, A. J. Boulton and A. McKillop, Pergamon Press, Oxford, 1984, vol. 3.
- S. D. Kahn and W. L. Hehre, J. Org. Chem., 1988, 53, 301;
 M. J. S. Dewar and C. Jie, J. Am. Chem. Soc., 1989, 111, 511.
- 11 J. S. Burnier and W. L. Jorgensen, J. Org. Chem., 1984, 49, 3001.

 $[\]P$ Reaction time is not mentioned, and only one aryl allenylmethyl ether was tried by the authors.