


This article was downloaded by: [Korea University]

On: 02 January 2015, At: 17:40

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

BiCl₃ Catalyzed Thiolyzation Reaction of 1,2-Epoxydes with Diaryl Disulfides in the Presence of Zinc Powder and Ionic Liquid

A. R. Khosropour^a, M. M. Khodaei^a & K. Ghazati^a

^a Department of Chemistry, Razi University, Kermanshah, Iran

Published online: 16 Aug 2006.

To cite this article: A. R. Khosropour, M. M. Khodaei & K. Ghazati (2005) BiCl₃ Catalyzed Thiolyzation Reaction of 1,2-Epoxydes with Diaryl Disulfides in the Presence of Zinc Powder and Ionic Liquid, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 180:11, 2525-2531, DOI: [10.1080/104265090929931](https://doi.org/10.1080/104265090929931)

To link to this article: <http://dx.doi.org/10.1080/104265090929931>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

BiCl₃ Catalyzed Thiolyzation Reaction of 1,2-Epoxides with Diaryl Disulfides in the Presence of Zinc Powder and Ionic Liquid

A. R. Khosropour

M. M. Khodaei

K. Ghozati

Department of Chemistry, Razi University, Kermanshah, Iran

A new and efficient method has been developed for the regio- and chemoselective synthesis of β -hydroxysulfides with the ring-opening reaction of 1,2-epoxides by diaryl disulfides and zinc powder in the presence of a catalytic amount of BiCl₃ in molten tetrabutyl phosphonium bromide as an ionic liquid.

Keywords β -hydroxysulfides; aryl disulfides; BiCl₃; epoxides; ionic liquid; ring opening reaction

INTRODUCTION

In recent years, β -hydroxysulfide derivatives have attracted considerable interest due to their wide utilization in organic synthesis for the preparation of benzoxathiepinines,¹ benzotiazepines,² β -hydroxysulfoxides,³ or α -thioketones,⁴ which are versatile in the realm of pharmaceutical, biological, or organic chemistry. Thus, the synthesis of this heterochemical system is of much interest. One of the classical synthetic methods for the preparation of β -hydroxysulfides consists of a ring opening of an epoxide with an excess of thiol.⁵ Because some functional groups may be sensitive to high temperature, some modified methods have been developed.⁶ However, many of these procedures are associated with one or more disadvantages, such as involving toxic and expensive catalysts or hazardous and carcinogenic organic solvents for reactions. In addition, low yields, long reaction times, and the utilization of large amounts of catalyst or solvent limit these synthetic scopes

Received September 14, 2004; accepted December 1, 2004.

The authors are grateful to the Razi University Research Council for partial support of this work.


Address correspondence to A. R. Khosropour, Razi University, Department of Chemistry, Faculty of Science, Kermanshah 67149, Iran. E-mail: arkhosropour@razi.ac.ir

to small scales. Hence, a simple, green, and efficient procedure avoiding these drawbacks that leads to convenient procedures and better yields is recommended.

RESULTS AND DISCUSSION

The toxic and volatile nature of many organic solvents have posed a serious threat to the environment. Consequently, methods that minimize their use are the focus of much attention. In this respect, ionic liquids are attracting a growing interest as an alternative reaction media for various chemicals and biotransformations.⁷ These compounds offer an alternative and ecologically friendly media that, in contrast to organic solvents, are nonvolatile and non-explosive.⁸ On the other hand, the catalysts that have polar or ionic character can be immobilized without additional structural modification, and thus, the ionic solutions containing the catalyst easily can be separated from products. Thus, ionic liquids have proven to be excellent solvents for many organic reactions.⁹

Most bismuth (III) salts are relatively nontoxic, easy to handle, and can tolerate small amounts of moisture.¹⁰ With increasing environmental concerns and the need for green reagents, the interest in bismuth (III) salts has increased tremendously in the last decade.¹¹ In continuation of our studies using Bi(III) salts for various organic transformations,¹² we wish to present a new and convenient method for the synthesis of β -hydroxysulfides toward the ring opening of epoxides with diaryl disulfides and zinc powder catalyzed by BiCl_3 that is immobilized on molten tetrabutylphosphonium bromide (TBPB) as an ionic liquid (Scheme 1).



SCHEME 1

The experimental procedure is very simple. To the mixture of molten TBPB and BiCl_3 , zinc powder, diaryl disulfide and 1,2-epoxide were


TABLE I Ring Opening of 1,2-Epoxydes With ArSSAr in the Presence of Zinc Powder and 10 mol% of BiCl₃ in the Presence of Molten TBPB

Entry	Epoxide	Ar	Yield (%) ^a / Time(min)	Product
1		Ph	92/20	$\text{OHCH}_2\text{CH}(\text{Ph})\text{—SAr}$
2		<i>P</i> -CH ₃ C ₆ H ₄	92/20	$\text{OHCH}_2\text{CH}(\text{Ph})\text{—SAr}$
3		2-Naphthyl	88/25	$\text{OHCH}_2\text{CH}(\text{Ph})\text{—SAr}$
4		Ph	80/60 ^b	
5		<i>P</i> -CH ₃ C ₆ H ₄	80/45 ^b	
6		2-Naphthyl	78/65 ^b	
7		Ph	89/40	$\text{PhOCH}_2\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$
8		<i>P</i> -CH ₃ C ₆ H ₄	90/45	$\text{PhOCH}_2\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$
9		Ph	80/65	$\text{CH}_2=\text{CHCH}_2\text{OCH}_2\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$
10		<i>P</i> -CH ₃ C ₆ H ₄	80/65	$\text{CH}_2=\text{CHCH}_2\text{OCH}_2\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$
11		Ph	86/30	$\text{CH}_3(\text{CH}_2)_5\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$
12		Ph	90/60	$\text{CH}_3(\text{CH}_2)_2\text{CH}_2\text{OCH}_2\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$
13		Ph	86/55	$\text{ClCH}_2\text{CH}(\text{OH})\text{CH}_2\text{—SAr}$

^aIsolated yields.^bIn the presence of 20 mol% of BiCl₃.

added and the system was stirred at 65°C and the product was isolated by usual work-up. The results are summarized in Table I. In the method described here, a variety of epoxides with different functional groups such as PhO-, RO-, ClCH₂-, and allyl ether were converted to the products with high efficiency and chemoselectivity. It is important to note that in the case of terminal epoxides (Table I, entries 7–13), the products were obtained by the attack of nucleophiles on the terminal carbon. This regioselectivity is probably due to the steric hinderance of nucleophile that attacks on the less-hindered carbon of the epoxides.

TABLE II Competitive Ring Opening Reaction of 1,2-Epoxydes With Diaryl Disulfides in the Presence of Zinc Powder and BiCl₃ in the Presence of Molten TBPB



However, in the case of the reaction with styrene oxide, we observed a reverse regioselectivity in which the attack on the benzylic carbon (which may be due to the formation of more stable carbocation on benzylic carbon) was preferred (Table I, entries 1–3). The products that were obtained using cyclohexene oxide possess a *trans*-configuration by ¹H NMR spectroscopy (Table I, entries 4–6). This high reactivity and selectivity is similar to those observed in the case of strong Lewis acids with thiophenols.⁶ On the other hand, we found that aliphatic disulfides such as dibenzyl disulfide or dicyclohexyl disulfide are quite resistant to the reaction conditions and completely recovered.

In order to evaluate the intermolecular chemoselectivity of the reagents, we also examined the competitive reactions (Table II). It is noteworthy that high chemoselectivity was exhibited with styrene oxide and the corresponding β -hydroxysulfide that was obtained in the presence of the other epoxides. In this competitive reaction, other acid-sensitive epoxides such as cyclohexane oxide in the presence of styrene oxide were stable.

CONCLUSIONS

In conclusion, we have demonstrated a new, efficient, and green catalytic procedure for the regio- and chemoselective thiolysation of 1,2-epoxides with diaryl disulfides in the ionic liquid media. In addition,

the ease of availability and low toxicity of the catalyst and the ionic liquid make this procedure environmentally acceptable. Furthermore, the low cost of the catalyst and ionic liquid, the high catalytic nature of the Lewis acid, and fast reaction rates make the present procedure a practical protocol for the β -hydroxysulfide synthesis.

EXPERIMENTAL

All yields refer to isolated products. The products were characterized by comparison with authentic samples.⁶ All ¹H and ¹³C NMR spectra were recorded on a Bruker Avance 200 MHz spectrophotometer in CDCl₃ as a solvent. Diaryl disulfides were prepared according to the literature.¹³ BiCl₃-TBPB was prepared by mixing BiCl₃ (0.1–0.2 mmol) and molten tetrabutylphosphonium bromide (0.5 mmol), which heated to 120°C, then this mixture cooled to 65°C.

General Procedure for the Ring Opening Reaction of 1,2-Epoxides With Diaryl Disulfides in the Presence of Zinc Powder Catalyzed by BiCl₃-TBPB

In a 50 mL round bottom, 1,2-epoxide (1 mmol), diaryl disulfide (1 mmol), and zinc powder (1 mmol) were added to a molten mixture of tetrabutylphosphonium bromide (0.5 mmol) and BiCl₃ (0.1–0.2 mmol), then the reaction mixture stirred at 65°C for the appropriate times (Table I). The progress of the reaction was followed by TLC (n-heptane/ethyl acetate: 3/1). After completion of the reaction, Et₂O (20 mL) was added and the mixture was filtered. The evaporation of the organic layer gave the corresponding crude products, which were purified by chromatography on a silica-gel plate to afford the pure β -hydroxythioethers in 78–92% yields. Spectroscopy data: Product of entry **1**. IR ν_{\max} cm⁻¹: 3370, 3005, 2895, 2860, 1585, 1465, 1050, 740, 695 (neat). ¹H NMR (200 MHz, CDCl₃) δ 7.15–7.55 (m, 10H), 4.35 (t, J 6.8 Hz, 1H), 3.95 (d, J 6.8 Hz, 2H), 2.10 (s, 1H). ¹³C NMR (50 MHz, CDCl₃) δ 139.40, 134.20, 132.90, 129.40, 129.20, 128.50, 128.20, 128.00, 65.70, 56.50. Product of entry **2**. IR ν_{\max} cm⁻¹: 3430, 3015, 2905, 2880, 1590, 1495, 1440, 1060, 850, 735, 690 (neat). ¹H NMR (200 MHz, CDCl₃) δ 7.50–7.90 (m, 9H), 4.75 (t, J 6.9 Hz, 1H), 4.35 (d, J 6.9 Hz, 2H), 2.80 (s, 4H). ¹³C NMR (50 MHz, CDCl₃) δ 139.60, 138.30, 133.70, 130.30, 130.20, 129.10, 128.60, 128.20, 65.50, 56.80, 21.60. Product of entry **3**. IR ν_{\max} cm⁻¹: 3520, 3030, 2905, 2890, 1595, 1490, 1450, 1090, 850, 730, 695 (neat). ¹H NMR (200 MHz, CDCl₃) δ 7.30–7.95 (m, 12H), 4.52 (t, J 8.0 Hz, 1H), 4.05 (d, J 7.0 Hz, 2H), 2.25 (s, 1H). ¹³C NMR (50 MHz, CDCl₃) δ 139.30, 133.90, 132.80, 131.80, 131.50, 130.10, 129.20, 128.90, 128.60,

128.30, 128.10, 127.90, 127.00, 126.80, 65.70, 56.40. Product of entry **4**. IR ν_{\max} cm^{-1} : 3650, 3015, 2905, 2885, 1590, 1495, 1440, 1065, 840, 735, 690 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.25–7.60 (m, 5H), 3.30–3.50 (m, 1H), 2.90 (s, 1H), 2.70–2.85 (m, 1H), 2.05–2.25 (m, 2H), 1.62–1.83 (m, 2H), 1.20–1.50 (m, 4H). ^{13}C NMR (50 MHz, CDCl_3) δ 134.20, 133.00, 129.30, 128.20, 72.40, 56.90, 34.20, 33.10, 26.60, 24.70. Product of entry **5**. IR ν_{\max} cm^{-1} : 3450, 3010, 2900, 2880, 1590, 1495, 1450, 1060, 820 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.85 (d, J 10.0 Hz, 2H), 7.55 (d, J 10.0 Hz, 2H), 3.70–3.85 (m, 1H), 3.55 (s, 1H), 3.10–3.25 (m, 1H), 2.8 (s, 3H), 2.50–2.65 (m, 2H), 2.05–2.20 (m, 2H), 1.10–1.60 (m, 4H). ^{13}C NMR (50 MHz, CDCl_3) δ 138.50, 134.90, 130.10, 128.90, 72.20, 56.90, 34.20, 32.90, 26.60, 24.70, 21.60. Product of entry **6**. IR ν_{\max} cm^{-1} : 3440, 3020, 2905, 2885, 1595, 1490, 1440, 1085, 810 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.50–8.10 (m, 7H), 3.40–3.60 (m, 1H), 2.90–3.05 (m, 2H), 2.10–2.25 (m, 2H), 1.70–1.90 (m, 2H), 1.15–1.60 (m, 4H). ^{13}C NMR (50 MHz, CDCl_3) δ 133.90, 133.30, 132.90, 131.50, 130.30, 128.90, 128.10, 127.90, 127.00, 126.90, 72.50, 57.00, 34.30, 33.20, 26.60, 24.70. Product of entry **7**. IR ν_{\max} cm^{-1} : 3400, 3010, 2900, 2880, 1590, 1480, 1245, 1050, 730, 695 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.85–7.70 (m, 10H), 3.95–4.25 (m, 3H), 3.10–3.40 (m, 2H), 2.75 (s, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ 158.80, 135.60, 130.25, 130.00, 129.60, 127.10, 121.70, 115.00, 70.50, 69.00, 38.00. Product of entry **9**. IR ν_{\max} cm^{-1} : 3350, 3020, 2905, 2880, 1600, 1490, 1380, 1060, 735, 690 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.15–7.50 (m, 5H), 5.80–6.05 (m, 1H), 5.15–5.35 (m, 2H), 4.00 (d, J = 6.8 Hz, 2H), 3.80–3.95 (m, 1H), 3.45–3.65 (m, 2H), 3.00–3.25 (m, 2H), 2.70 (s, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ 135.90, 134.80, 130.10, 129.50, 126.90, 117.80, 72.80, 72.70, 69.40, 37.90. Product of entry **10**. IR ν_{\max} cm^{-1} : 3700, 3015, 2905, 2885, 1590, 1495, 1440, 1080, 790 (neat). ^1H NMR (200 MHz, CDCl_3) 7.75 (d, J = 8.0 Hz, 2H), δ 7.55 (d, J 8.0 Hz, 2H), 6.20–6.45 (m, 1H), 5.60–5.80 (m, 2H), 4.40 (d, 2H), 4.25–4.35 (m, 1H), 3.90–4.10 (m, 2H), 3.35–3.65 (m, 2H), 3.20 (s, 1H), 2.80 (s, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 137.10, 134.80, 132.10, 130.90, 130.20, 117.70, 72.80, 72.70, 69.30, 38.70, 21.40. Product of entry **11**. IR ν_{\max} cm^{-1} : 3405, 3015, 2905, 2880, 1585, 1490, 1435, 735, 690 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.15–7.50 (m, 5H), 3.50–3.80 (m, 1H), 3.10–3.25 (dd, J 4.5 Hz, J 4.5 Hz, 2H), 2.80–2.95 (dd, J 9.5 Hz, J 9.5 Hz, 2H), 2.70 (s, 1H), 1.15–1.70 (m, 8H), 0.95 (t, J 7.0 Hz, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 136.00, 130.30, 130.00, 126.90, 69.90, 42.50, 36.60, 32.20, 29.60, 26.10, 23.00, 14.50. Product of entry **12**. IR ν_{\max} cm^{-1} : 3410, 3025, 2900, 2890, 1590, 1490, 1430, 735, 690 (neat). ^1H NMR (200 MHz, CDCl_3) δ 7.15–7.50 (m, 5H), 3.85–4.00 (m, 1H), 3.40–3.65 (m, 4H), 3.00–3.25 (m, 2H), 2.80 (s, 1H), 1.55–1.70 (m, 2H), 1.30–1.50 (m, 2H), 0.95 (t, J 8.3 Hz, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 136.10, 129.90, 129.40, 126.70, 73.70,

71.70, 69.40, 37.80, 32.10, 19.70, 14.30. Product of entry **13**. IR ν_{\max} cm⁻¹: 3425, 3010, 2920, 2830, 1600, 1490, 1430, 1365, 1110, 740, 690 (neat). ¹H NMR (200 MHz, CDCl₃) δ 7.20–7.50 (m, 5H), 3.90–4.10 (m, 1H), 3.72 (d, J = 8.0 Hz, 2H), 3.05–3.18 (m, 2H), 2.88 (s, 1H). ¹³C NMR (50 MHz, CDCl₃) δ 135.10, 130.50, 129.70, 127.30, 69.90, 48.40, 38.60.

REFERENCES

- [1] (a) C. G. Aifheli and P. T. Kaye, *Synth. Commun.* **26**, 4459 (1996); (b) H. Sugihara, H. Mabuchi, M. Hirata, T. Iamamoto, and Y. Kawamatsu. *Chem. Pharm. Bull.*, **35**, 1930 (1987).
- [2] M. A. Adger, J. B. Barkley, S. Bergeron, M. W. Cappi, B. E. Floweden, M. P. Jackson, R. McCange, T. C. Nugent, and S. M. Roberts. *J. Chem. Soc. Perkin Trans.*, **1**, 3501 (1997).
- [3] (a) C. Gabbi, F. Ghelgi and R. Grandi. *Synth. Commun.*, **27**, 2857 (1997); (b) C. Alvarez-Ibarra, R. Cuervo-Rodriguez, M. C. Fernanedz-Monreal, and M. P. Ruiz. *J. Org. Chem.*, **59**, 7287 (1994).
- [4] J.-P. Bégué, D. Monnet-Delpon, and A. Kornilov. *Synthesis* 529 (1996).
- [5] G. Capozzi and G. Modena, *The Chemistry of the Thiol Group* (John Wiley, New York, 1974).
- [6] (a) F. Fringuelli, F. Pizzo, S. Tortoioli, and L. Vaccaro. *Green Chem.*, **5**, 436 (2003); (b) C. Alan, A. C. Spivey, R. Srikanan, C. M. Diaper, J. David, and D. Turner. *J. Org. Biomol. Chem.*, **1**, 1638 (2003); (c) F. Fringuelli, F. Pizzo, S. Tortoioli, and L. Vaccaro. *Tetrahedron Lett.*, **44**, 6785 (2003); (d) D. Amantini, F. Fringuelli, F. Pizzo, S. Tortoioli, and L. Vaccaro. *Synthesis*, 2292 (2003); (e) J. S. Yadav, B. V. S. Reddy, and G. Baishya. *Chem. Lett.*, 906 (2002).
- [7] (a) T. Welton, *Chem. Rev.*, **99**, 2071 (1999); (b) P. Wasserscheid and W. Keim, *Angew. Chem. Int. Ed.*, **39**, 3772 (2000).
- [8] (a) B. Ganchegui, S. Bouquillon, F. Henin, and J. Muzart, *Tetrahedron Lett.*, **43**, 6641 (2002); (b) B. C. Ranu, A. Das, and S. Samanta, *J. Chem. Soc., Perkin Trans.*, **1**, 1520 (2002); (c) K. Selvakumar, A. Zapt, and M. Beller, *Org. Lett.*, **4**, 3031 (2002).
- [9] (a) J. R. Harjani, S. J. Nara, and M. M. Salunkhe, *Tetrahedron Lett.*, **43**, 1127 (2002); (b) V. V. Namboodiri and R. S. Varma, *Chem. Commun.*, 342 (2002); (c) R. Sheldon, *Chem. Commun.*, 2399 (2001).
- [10] (a) S. Cunha, B. R. Lima, and A. R. Souza. *Tetrahedron Lett.*, **43**, 49 (2002); (b) S. Cunha, B. R. Lima, and A. R. Souza. *Tetrahedron Lett.*, **43**, 49 (2002); (c) G. Sabitha, E. V. Reddy, R. Swapna, N. M. Reddy, and J. S. Yaddav, *Synlett.*, 1276 (2004).
- [11] T. Ollevier and G. Lavie-Compin, *Tetrahedron Lett.*, **45**, 49 (2004).
- [12] (a) I. Mohammadpoor-Baltork and A. R. Khosropour, *Monatsh. Chem.*, **133**, 189 (2002); (b) I. Mohammadpoor-Baltork, A. R. Khosropour, and H. Alian. *J. Chem. Res. (S)*, 280 (2001); (c) I. Mohammadpoor-Baltork, H. Alian, and A. R. Khosropour. *Tetrahedron*, **57**, 5851 (2001); (d) M. M. Khodaei, A. R. Khosropour, and K. Ghozati. *Tetrahedron Lett.*, **45**, 3525 (2004). (e) M. M. Khodaei, A. R. Khosropour, and S. J. Hoseini Jomor. *J. Chem. Res. (S)*, **10**, 638 (2003); (f) A. R. Khosropour, M. M. Khodaei, and K. Ghozati. *Chem. Lett.*, **33**, 304 (2004); (g) A. R. Khosropour, M. M. Khodaei, and M. Khookhazadeh. *Tetrahedron Lett.*, **45**, 1725 (2004).
- [13] (a) H. Firouzabadi, N. Iranpoor, and M. A. Zolfigol, *Synth. Commun.*, **28**, 1179 (1998); (b) A. R. Hajipour and S. E. Mallakpour, *J. Chem. Research (S)*, 32 (2000).