

0040-4039(94)02180-5

A HIGHLY ASYMMETRIC PUMMERER-TYPE CYCLIZATION OF CHIRAL, NON-RACEMIC β-AMIDOSULFOXIDES INDUCED BY O-SILYLATED KETENE ACETALS

Yasuyuki Kita,*a Norio Shibata,^a Noriyuki Kawano,^a Takashi Tohjo,^a Chino Fujimori,^a and Keita Matsumoto^b

^a Faculty of Pharmaceutical Sciences, Osaka University 1-6, Yamada-oka, Suita, Osaka 565, Japan ^b Recearch Center, Taisho Pharmaceutical Co., Ltd., 1-403, Yoshino-cho, Omiya, Saitama 330, Japan

Abstract: The first highly asymmetric Pummerer-type cyclization of chiral, non-racemic β -amidosulfoxides leading to enantiomerically enriched β -lactams (80-85 % ee) is described. S-and R-sulfoxides (S-2a-d and R-2a-c) were treated with O-methyl-O-tert-butyldimethylsilyl ketene acetal (1) in the presence of a catalytic amount of ZnCl₂ in CH₂Cl₂ to predominantly give the corresponding 4R- and 4S- β -lactams (R-4a-d and S-4a-c) in more than 80 % ee. These results show that the stereoinduction is completely influenced by the absolute configuration of the sulfoxides.

The asymmetric Pummerer reaction of chiral, non-racemic sulfoxides,¹⁾ which is one of the selfimmolative-type asymmetric inductions, is of significant interest, because it would provide a means for the synthesis of enantiomerically pure α -substituted sulfides.²⁾ Especially the intramolecular version of the asymmetric Pummerer-type reaction is quite useful for the synthesis of optically active heterocyclic compounds.^{3, 4)} Only a few examples of these types of reactions have been reported, but the enantiomeric excess (ee) yields were low.^{3, 4)} We report here the first highly asymmetric Pummerer-type cyclization of chiral, non-racemic β -amidosulfoxides leading to enantiomerically enriched β -lactams (80-85 % ee).

Several years ago, we reported a novel silicon-induced Pummerer-type reaction of sulfoxides using Omethyl-O-tert-butyldimethylsilyl ketene acetal (1), which gave α -siloxy sulfides under mild conditions ⁵) and then applied it to the intramolecular cyclization of ω -amidosulfoxides to produce α -thio-N-heterocycles involving 4- to 7-membered α -thiolactams.^{4, 6}) Very recentry, we reported the first highly asymmetric Pummerer-type reaction of chiral, non-racemic acyclic sulfoxides which gave enantiomerically enriched α siloxysulfides in high yields.⁷) We have now applied this reaction to the asymmetric intramolecular cyclization of chiral, non-racemic β -amidosulfoxides (2a-d).

Starting materials 2a-d were prepared in good yields from the known carboxylic acid (3)⁸⁾ by condensation with the corresponding amine in the presence of 1,3-dicyclohexylcarbodiimide (DCC) in dimethylformamide (DMF) (Scheme 1).





Under the standard silicon-induced Pummerer conditions ^{4, 6}) (1, cat. ZnI₂, MeCN), treatment of optically pure S-sulfoxides (2a), which have the chiral amido group, gave R- β -lactam (4a) in 60 % de (72 % chemical yield) stereoselectively. The use of ZnCl₂ as a catalyst in CH₂Cl₂ was found to give the best results. Surprisingly high ee's were also obtained from 2b-d having an achiral amido group under the same conditions. S-and R-Sulfoxides (S-2 and R-2) were treated with 1 to predominantly give 4R- and 4S- β -lactams (R-4 and S-4), respectively in more than 80 % ee's. These results show that the stereoinduction is influenced by the absolute configuration of the sulfoxides (Table 1). Optically pure R- and S-4c were readily obtained by simple recrystallization in about 60 % yield {R-4c; $[\alpha]_D^{17} - 47.03^\circ$ (c=0.370, CHCl₃), S-4c; $[\alpha]_D^{20} + 49.4^\circ$ (c=0.599, CHCl₃). The present Pummerer cyclization has the highest optical induction compared with the previously examined method.^{3, 4})





Product [a]						
2a-d	R	Conditions	4	% Ee [b] (% Yield)	[α] _D (<i>c</i> , CHCl ₃)	CD (MeOH) λ ext (Δε)
S-2a	CH(Me)Ph [c]	0°C, 1d [d]	R-4a	60 [e] (72)	- 98.8 (1.96) [f]	253 (+2.9), 242 (0), 222 (-20.4)[f]
<i>S-</i> 2a	CH(Me)Ph [c]	0°C, 3d	R-4a	82 [e] (96)	- 98.8 (1.96) [f]	253 (+2.9), 242 (0), 222 (-20.4)[f]
R-2a	CH(Me)Ph [c]	0°C, 3d	S-48	85 [e] (89)	+116 (0.954) [f]	254 (-5.7), 240 (0), 221 (+24.5)[f]
<i>S</i> -2b	CH ₂ Ph	5°C, 6d	<i>R</i> -4b	80 (54)	-73.2 (1.01)	252 (+3.1), 239 (0), 220 (-16.6)
R-2b	CH ₂ Ph	5°C, 6d	S-4b	82 (54)	+75.2 (0.934)	252 (-2.6), 240 (0), 220 (+19.2)
S-2c	CHPh2	15°C, 2d	R-4c	80 (84)	-37.0 (0.303)	253 (+5.1), 243 (0), 221 (-21.5)[g]
<i>R</i> -2c	CHPh2	15°C, 2d	S-4c	83 (90)	+38.4 (0.522)	255 (-4.3), 243 (0), 221 (+21.4)[g]
<i>S</i> -2 d	CH2-0-BrPh	5°C, 7d	<i>R</i> -4d	83 (5 9)	-59.2 (1.49)	253 (+1.5), 240 (0), 221 (-14.4)

[a] Small amount of α -siloxy- β -amidosulfides were obtained as side-products in all reactions. [b] Ee value was determined by ¹H-NMR (C₆D₆) with Eu(hfc)₃. [c] S-Phenethylamine was used. [d] Znl₂ is used as a catalyst in MeCN. [e] % De. [f] This value was obtained from the diastereomerically pure sample. [g] This value was obtained from the enantiomerically pure sample.

The absolute stereochemistry at the newly generated chiral center of the β -lactams was ascertained by its CD-spectrum, which had a strong positive or negative Cotton effect at 210-220 nm (Octant rule) in agreement with the value from monocyclic β -lactams reported by Reling and Jensen ⁹) and confirmed by the conversion of *R*-4c to the known (+)-PS-5 intermediate {(-)-5; $[\alpha]_D^{22} - 40.9^\circ$ (c=1.02, CHCl₃), lit.⁶¹; $[\alpha]_D^{25} - 39.59^\circ$ (c=2.92, CHCl₃)}.¹⁰ Finally, the absolute stereochemistry was determined by X-ray crystallographic analysis ¹¹) of oxidized derivertive (*R*-6) of *R*-4c. (Fig. 1)



Fig 1. X-Ray Crystallographic Structure of R-6

The following mechanism is proposed to explain the results and the transition-state is representatively analyzed for the reaction of S-2 with 1 (Fig. 2). Silylation of S-2 with 1 affords an intermediate (A). Thus, A may yield a chiral pseudo isothiazolone derivative (B) 12) through axial attack of the amido anion generated by abstraction with ester anion and elimination of the siloxy ligand. Then, the hydrogen neighboring to the sulfur atom was removed by the siloxy anion and the amido ligand was rearranged from the α -face to give the β -lactam (R-4).



Fig. 2

References and Notes

- Asymmetric Pummerer rearrangement; a) McCormick, J. E.; McElhinney, R. S. Chem. Commun. 1969, 171; b) Glue, S.; Kay, I. T.; Kipps, M. R. Chem. Commun. 1970, 1158; c) Numata, T.; Oae, S. Tetrahedron Lett. 1977, 1337; d) Numata, T.; Itoh, O.; Oae, S. Chem. Lett. 1977, 909; e) Mikolajczyk, M.; Zatorski, A.; Grzejszczak, S.; Costisella, B.; Midura, W. J. Org. Chem. 1978, 43, 2518; f) Numata, T.; Itoh, O.; Oae, S. Tetrahedron Lett. 1979, 1869; g) Masuda, T.; Numata, T.; Furukawa, N.; Oae, S. J. Chem. Soc., Perkin Trans. 2 1978, 1302; h) Numata, T.; Itoh, O.; Yoshimura, T.; Oae, S. Bull. Chem. Soc. Jpn. 1983, 56, 257; i) Itoh, O.; Numata, T.; Yoshimura, T.; Oae, S. Bull. Chem. Soc. Jpn. 1983, 56, 266. Asymmetric additive Pummerer reaction (enantioselective 3,3-sigmatropic rearrangement); Marino, J. P.; Perez, A. D. J. Am. Chem. Soc. 1984, 106, 7643.
- Reviews, see: a) Oae, S.; Numata, T. " The Pummerer and Pummerer Type of Reactions," in "Isotopes in Organic Chemistry," ed. by Buncel, E.; Lee, C. C. Elsevier, New York, 1980, Vol. 5. Chapter 2; b) Lucchi, O. D.; Miotti, U.; Modena, G. Organic Reactions 1990, 40, 157.
- a) Stridsberg, B.; Allenmark, S. Acta Chem. Scand. 1974, B28, 591; Stridsberg, B.; Allenmark, S. Acta Chem. Scand. 1976, B30, 219; b) Wolfe, S.; Kazmaier, P. M.; Auksi, H. Can. J. Chem. 1979, 57, 2404; c) Kaneko, T.; Okamoto, Y.; Hatada, K. J. Chem. Soc., Chem. Commun. 1987, 1511.
- 4. Kita, Y.; Tamura, O.; Miki, T.; Tono, H.; Shibata, N.; Tamura, Y. Tetrahedron Lett. 1989, 30, 729.
- a) Kita, Y.; Yasuda, H.; Tamura, O.; Itoh, F.; Tamura, Y. *Tetrahedron Lett.* 1984, 25, 4681; b) Kita, Y; Yasuda, H.; Tamura, O.; Itoh, F.; Tamura, Y. *Chem. Pharm. Bull.* 1985, 33, 4235; c) Kita, Y.; Tamura, O.; Itoh, F.; Yasuda, H.; Miki, T.; Tamura, Y. *Chem. Pharm. Bull.* 1987, 35, 562.
- a) Kita, Y.; Tamura, O.; Miki, T.; Tamura, Y. Tetrahedron Lett. 1987, 28, 6479; b) Kita, Y.; Tamura, O.; Shibata, N.; Miki, T. J. Chem. Soc., Perkin Trans. 1 1989, 1862; c) Kita, Y.; Tamura, O.; Shibata, N.; Miki, T. Chem. Pharm. Bull. 1990, 38, 1473; d) Kita, Y.; Shibata, N.; Miki, T.; Takemura, Y.; Tamura, O. J. Chem. Soc., Chem. Commun. 1990, 727; e) Kita, Y.; Shibata, N.; Tamura, O.; Miki, T. Chem. Pharm. Bull. 1991, 39, 2225; f) Kita, Y.; Shibata, N.; Miki, T.; Takemura, Y.; Tamura, O. Chem. Pharm. Bull. 1992, 40, 12.
- a) Kita, Y.; Shibata, N.; Yoshida, N. Tetrahedron Lett. 1993, 34, 4063; b) Kita, Y.; Shibata, N.; Yoshida, N.; Fukui, S.; Fujimori, C. Tetrahedron Lett. 1994, 35, 2569; c) Kita, Y.; Shibata, N.; Kawano, N.; Fukui, S.; Fujimori, C. Tetrahedron Lett. 1994, 35, 3575.
- 8. Albinati, A.; Bravo, P.; Ganazzoli, F.; Resnati, G.; Viani, F. J. Chem. Soc., Perkin Trans. 1 1986, 1405.
- 9. Rehling, H.; Jensen, H. Tetrahedron Lett. 1972, 2793.
- 10. The full details of this method for the formal total synthesis of (+)-PS-5 will be reported elsewhere.

- The crystal data for R-6 are as follow: orthorhombic; P2₁2₁2₁ with a=8.914(3), b=37.70(1), c=5.928(2) Å, V=1992(1) Å³, Z=4, and μ(Cu Kα)=15.29 cm⁻¹ by Mac Science MXC 18 instrument. Final R value was 0.038 for 1956 reflections. The supplementary materials have been deposited at the Cambridge Crystallographic Data Centre.
- The formation of the acylaminosulfonium ion (isothiazolone B) as an intermediate has been widely accepted under Pummerer-type conditions; a) Uchida, Y.; Kozuka, S. J. Chem. Soc., Chem. Commun. 1981, 510; b) Kaneko, T. J. Am. Chem. Soc. 1985, 107, 5490; c) Uchida, Y.; Oae, S. Cazzetta Chimica Italiana 1987, 117, 640; d) Yamamoto, K.; Yamazaki, S.; Murata, I. J. Org. Chem. 1987, 52, 5239; e) Wright, S. W.; Abelman, M. M.; Bostrom, L. L.; Corbett, R. C. Tetrahedron Lett. 1992, 33, 153.

(Received in Japan 25 August 1994; accepted 6 October 1994)