Selective Access to Electron Donating Group Substituted $(\eta^{5}$ -cyclohexadienyl)Mn(CO)₃ Complexes by Bimetallic **Manganese–Palladium Activation**

Audrey Auffrant, Damien Prim,[†] Françoise Rose-Munch,^{*} and Eric Rose^{*}

Laboratoire de Synthèse Organique et Organométallique, UMR CNRS 7611, Université Pierre et Marie Curie, Tour 44 Ier étage, 4 place Jussieu, BP 181, 75252 Paris Cedex 05, France

Jacqueline Vaissermann

Laboratoire de Chimie Inorganique et Matériaux, UMR CNRS 7071, Université Pierre et Marie Curie, 4 place Jussieu, BP 42, 75252 Paris Cedex 05, France

Received April 8, 2002

Summary: Unprecedented (η⁵-cyclohexadienyl)Mn(CO)₃ complexes bearing an electron-donating group at the end of the π system have been selectively prepared using Pd catalysis and primary, secondary cyclic, and acyclic amines. This catalytic methodology has been extended to O-, S-, and P-based derivatives. The X-ray structure of one coupling product exhibits an unusual conformation in its η^5 system.

In the course of reactions involving electrophilic η^{6} arene metal complexes and nucleophiles,¹ η^5 -cyclohexadienyl transition-metal complexes can be isolated or transformed in situ into highly fuctionalized arenes or cyclohexadienes. Those entities coordinated to chromium have been intensively studied, demonstrating their use in organic synthesis.²

In contrast, relatively few investigations have been undertaken to exploit the potential of (η^5 -cyclohexadienyl)manganese complexes as valuable tools for organic synthesis.³ This is probably due to the fact that, until recently, the only way to obtain the required (η^5 cyclohexadienyl)manganese complexes consisted of a nucleophilic addition to the parent η^6 complex. Such reactions are known to be efficient with carbon nucleo-

philes and hydrides.^{4,5} Nucleophilic attack on the substituted arene takes place at the 6- or 5-position of the aromatic ring, depending on the electronic effect of a substituent.⁶ Electron-withdrawing substituents such as chloride in complex 1 are known to induce mainly an ortho addition, giving complex **2** (Scheme 1, path a).^{6b,7} In contrast, electron-donating groups (4; Y = OMe, NR_2) exclusively direct meta addition, yielding complex 5 (Scheme 1, path b).^{3d} Indeed, electronic effects preclude ortho nucleophilic addition and thereby avoid the formation of the 1-substituted η^5 -complex **6** (Scheme 1, path c). In other words, no synthesis of (η^5 -cyclohexadienyl)manganese complexes substituted with an electron-donating group at the end of the π -system, **6**, had been previously reported. We thought that a possible strategy to prepare these complexes might be found in the reaction of the palladium intermediate 3, arising from the insertion of Pd(0) into the carbon-chloride bond of (η^5 -chlorocyclohexadienyl)manganese complexes 2,8 with nucleophiles such as amine, alcoholate, and thiolate in a Buchwald-Hartwig type methodology to afford complex 6 (Scheme 1, path a).⁹ Therefore, in this communication we describe an efficient synthesis of previously unknown complexes using a palladiumcatalyzed coupling reaction. While elaborate catalytic systems seem to be required and beneficial in Buchwald-Hartwig reactions, the present strategy takes

H. V.; Lotz, S. Organometallics 1990, 9, 293.
(6) (a) Pauson, P. L.; Segal, J. A. J. Chem. Soc., Dalton Trans. 1975, 1677. (b) Pauson, P. L.; Segal, J. A. J. Chem. Soc., Dalton Trans. 1975, 1683. (c) Stephenson, G. R.; Owen, D. A. Tetrahedron Lett. 1991, 32, 1291. (d) Rose-Munch, F.; Rose, E. Curr. Org. Chem. 1999, 3, 445.
(7) Chung, Y. K.; Williard, P. G.; Sweigart, D. A. Organometallics 1982, 1, 1053.

(8) (a) Prim, D.; Auffrant, A.; Rose-Munch, F.; Rose, E.; Vaisser-mann, J. *Organometallics* **2001**, *20*, 1901. (b) Auffrant, A.; Prim, D.; Rose-Munch, F.; Rose, E.; Vaissermann, J. Organometallics 2001, 20, 3214.

(9) See for example (a) Hartwig, J. F. Angew. Chem., Int. Ed. Engl. **1998**, 37, 2046. (b) Buchwald, S. L. J. Organomet. Chem. **1999**, 576, 125. (c) Prim, D.; Campagne, J. M.; Joseph, D.; Andrioletti, B. Tetrahedron **2002**, 58, 2041.

^{*} To whom correspondence should be addressed. E-mail: rose@ ccr.jussieu.fr (E.R.); rosemun@ccr.jussieu.fr (F.R.). [↑] Present address: Laboratoire SIRCOB, UPRESA A 8086, Univer-

sité de Versailles, 45 avenue des Etats-Unis, 78035 Versailles Cedex, France

^{(1) (}a) McQuillin, F.; Parker, D. G.; Stephenson, G. R. Transition Metal Organometallics for Organic Synthesis; Cambridge University Press: Cambridge, U.K., 1991; p 429. (b) Semmelhack, M. F. In *Comphrensive Organometallic Chemistry*, Abel, W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 12, Chapter 9, p 979. (c) Rose-Munch, F.; Gagliardini, V.; Renard, C.; Rose, E. Coord. Chem. Rev. 1998, 249, 198. (d) Pike, R. D.; Sweigart, D. A. Coord. Chem. Rev. 1999, 187, 183. (e) Astruc, D. Chimie organométallique; EDP Sciences: Les Ulis, France, 2000; p 111. (f) Rose-Munch, (2) See for example: (a) Semmelhack, M. F.; Hall, H. T., Jr.; Farina,

 ⁽a) Set tol example: (a) Seminerrack, M. F., Hali, H. L., Shi, Farma, R.; Yoshifuji, M.; Clark, G.; Bargar, T.; Hirotsu, K.; Clardy, J. J. Am. Chem. Soc. 1980, 101, 3535. (b) Pape, A. R.; Krishna, K. P.; Kündig, E. P. Chem. Rev. 2000, 100, 2917. (c) Bernardinelli, G.; Gillet, S.; Kündig, E. P.; Ronggang, L.; Ripa, A.; Saudan, L. Synthesis 2001, 13, 0010 2040

^{(3) (}a) Miles, W. H.; Brinkman, H. R. *Tetrahedron Lett.* **1992**, *33*, 589. (b) Lee, T. Y.; Kang, Y. K.; Chung, Y. K.; Pike, R. D.; Sweigart, D. A. *Inorg. Chem. Acta* **1993**, *214*, 125. (c) Roell, B. C.; McDaniel, K. F.; Vaughan, W. S.; Macy, T. S. *Organometallics* **1993**, *12*, 224. (d) Pearson, A. J.; Gontcharov, A. V.; Zhu, P. Y. *Tetrahedron* **1997**, *53*, 0440 3849 and references herein. (e) Pearson, A. J.; Vickerman, R. J. Tetrahedron Lett. **1998**, *39*, 5931.

⁽⁴⁾ McDaniel, K. F. In *Comprehensive Organometallic Chemistry*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 6, Chapter 4, p 93.

⁽⁵⁾ Addition of heteronucleophiles usually led to equilibria, reversible reactions, or reactions at the metal center; see, for example: (a) Evans, E. J.; Kane-Maguire, L. A. P.; Sweigart, D. A. J. Organomet. Chem. **1981**, 215, C27. (b) Chung, Y. K.; Honig, E. D.; Sweigart, D. A. J. Organomet. Chem. **1983**, 256, 277. (c) Schindehutte, M.; Rooyen, P. H. V.; Lotz, S. Organometallics **1990**, 9, 293.



15-20 (Table 2)

Table 1. Pd-Catalyzed Amination of $(\eta^{5}-1-chlorocyclohexadienyl)Mn(CO)_{3}$ Complexes

| entry | complex | YH | complex | yield (%) |
|-------------------------|------------------------------------|---|---------|-----------|
| 1 <i>a,c</i> | 7 , $R_1 = H$ | HN(CH ₂ CH ₂) ₂ O | 11 | 84 |
| $2^{b,c}$ | 7 , $R_1 = H$ | HN(CH ₂ CH ₂) ₂ O | | 0 |
| $3^{a,c}$ | 8 , R ₁ = 4-OMe | HNEt ₂ | 12 | 84 |
| 4 <i>a,c</i> | 9 , $R_1 = 4$ -Me | H ₂ NPh-4-OMe | 13 | 42 |
| 5 ^{<i>a,c</i>} | 10 , R ₁ = 2-OMe | HN(CH ₂ CH ₂) ₂ O | 14 | 0 |

^a Catalyst: Pd₂dba₃, AsPh₃. ^b Without catalyst. ^c Base: Cs₂CO₃.

advantage of the use of a simple Pd/AsPh $_3$ 10 catalytic combination.

The requisite $(\eta^5-1-\text{chlorocyclohexadienyl})Mn(CO)_3$ complexes 7-10 were readily prepared by hydride addition to the parent (η^6 -chloroarene)Mn(CO)₃⁺ complexes, as already described.¹¹ In a preliminary experiment, morpholine and Cs₂CO₃ were added to premixed complex 7, Pd₂(dba)₃, and AsPh₃ in freshly distilled and degassed THF under a nitrogen atmosphere (Scheme 2). Complete disappearance of the starting material was observed over 6 h at 60 °C, giving rise to the formation of 11 in 84% yield (Table 1, entry 1). A control experiment conducted under the same reaction conditions but without using a catalyst gave no reaction (entry 2). These initial results prompted us to extend this strategy to different substituted starting materials and a variety of amines. Indeed, the (η^{5} -1-chloro-4-substituted cyclohexadienyl)Mn(CO)₃ complexes 8 and 9 underwent amination using secondary and primary amines, respectively (entries 3 and 4). Only complex 10 did not display the expected amination product, probably due to steric hindrance (entry 5).

Next we examined the palladium-catalyzed reactions of various heteronucleophiles such as O-, S-, and Pbased derivatives (Table 2). Preformed sodium isoamylate reacted with complex **8** under the aforementioned

| Table 2. Pd-Catalyzed Reaction of |
|--|
| $(\eta^{5}-1-chlorocyclohexadienyl)Mn(CO)_{3}$ Complexes |
| with Alcohol, Thiols, Phosphine, and Phosphite |
| Oxide |

| entry | complex | YH | complex | yield (%) |
|------------------|---------|--------------------------|---------|-----------|
| 1 <i>a,b</i> | 8 | isoamyl-OH | 15 | 37 |
| $2^{a,b}$ | 8 | p-tolyl-SH | 16 | 93 |
| $3^{a,b}$ | 8 | $C_5H_{11}SH$ | 17 | 73 |
| 4 ^{a,b} | 7 | p-tolyl-SH | 18 | 81 |
| 5 ^{a,c} | 9 | (EtO) ₂ P(O)H | 19 | 77 |
| 6 ^{a,c} | 10 | PPh ₂ H | 20 | 65 |
| | | | | |

^a Catalyst: Pd₂dba₃,AsPh₃. ^b Base: NaH. ^c Base: Et₃N.

Scheme 3



conditions to afford the corresponding ether, **15**, in 37% yield (entry 1). Alkyl and aryl thioethers **16–18** were isolated in better yields (73–93%) using the same methodology (entries 2–4). Extension of this methodology to P-based nucleophiles was also successful. Indeed, diethyl phosphite oxide and diphenylphosphine gave the corresponding complexes **19** and **20** in 77 and 65% yields, respectively (entries 5 and 6). Reactions involving P-based nucleophiles require the use of triethylamine instead of NaH as a base.

Taking advantage of these observations, we have carried out further investigations using complex **21**, in which the sp³ carbon is substituted by a phenyl group.¹² Combinations of morpholine/Cs₂CO₃ and *p*-tolylSH/NaH under palladium catalysis were effective for the introduction of a nucleophile at position 1 (Scheme 3). No problem with steric hindrance was detected. Complexes

^{(10) (}a) Farina, V.; Krishnan, B.; Marshall, D. R.; Roth, G. P. J. Org. Chem. 1993, 58, 5434. (b) For an application with tricarbonylchromium complexes see: Prim, D.; Tranchier, J. P.; Rose-Munch, F.; Rose, E.; Vaissermann, J. Eur. J. Inorg. Chem. 2000, 5, 901.
(11) Balssa, F.; Gagliardini, V.; Rose-Munch, F., Rose, E. Organo-

⁽¹¹⁾ Balssa, F.; Gagliardini, V.; Rose-Munch, F., Rose, E. Organometallics **1996**, *15*, 4373.

⁽¹²⁾ Complex ${\bf 21}$ was prepared according to the procedure described in ref 7.

⁽¹³⁾ A typical procedure for the Pd-catalyzed reaction is given by the preparation of complex 16: Pd2(dba)3 (0.029 g, 0.03 mmol) and AsPh₃ (0.34 g, 0, 11 mmol) were added to a solution of complex **2** (0.089 g, 0.32 mmol) in anhydrous degassed THF (10 mL). After 15 min at room temperature, a THF solution of p-methylthiophenol (0.040 g, 0.32 mmol) and NaH (0.005 g, 0.38 mmol) was introduced. The reaction mixture was refluxed overnight, poured into cold water (100 mL), and extracted with 50 mL of pentane; the organic phase was washed with water (2 \times 20 mL), dried over magnesium sulfate, and filtered. Solvents were evaporated under a nitrogen flush. The residue obtained was purified by silica gel chromatography (pentane) to deliver complex **16** as a yellow solid in 93% yield. IR (CH₂Cl₂): ν (CO) 1917, 1998 cm⁻¹. ¹H NMR (CDCl₃; 200 MHz): δ 2.91 (dd, J = 13 and 6.5 Hz, 1H, H-6 endo), 2.48 (d, J = 13 Hz, 1H, H-6 exo), 2.32 (s, 3H), 3.01 (d, J = 6.5 Hz, 1H, H-5), 3.45 (s, 3H), 4.96 (d, J = 5.5 Hz, 1H, H-2), 5.65 (d, J = 5.5 Hz, 1H, H-3), 7.10 (d, J = 8 Hz), 7.21 (d, J = 8 Hz). ¹³C NMR (CDCl₃; 100 MHz): δ 21.2 (CH₃), 33.0 (C-6), 36.3 (C-5), 54.5, (OMe), 65.0 (C-1), 66.9 (C-3), 96.2 (C-2), 128.8, 130.0, 131.7, 133.8, (C₆H₄), 142.7 (C-4), 221.3 (CO). Anal. Calcd: C, 55.14; H, 4.08. Found: C, 55.12; H, 4.10. Crystal data for 16: C₁₇H₁₅MnO₄S, $M_r = 370.3$, orthorhombic, space group $P2_12_12_1$, a = 8.463(2) Å, b = 13.716(6) Å, c= 14.385(3) Å, $D_c = 1.47$ g cm⁻³, Z = 4, crystal size 0.10 × 0.40 × 0.40 mm, $\mu = 8.94$ cm⁻¹, 2325 data collected at 295 K (θ range 1–28°) on an Enraf-Nonius CAD-4 diffractometer. An absorption correction using DIFABS ($T_{min} = 0.65$, $T_{max} = 1$) was applied. Anomalous dispersion terms and correction of secondary extinction were applied. The structure was solved by direct methods (SHELXS) and refined on F (CRYSTALS) by least-squares analysis using anisotropic thermal parameters for all non-hydrogen atoms. H atoms were introduced in calculated positions and were allocated an overall isotropic thermal parameter. A total of 1356 reflections with $I > 3\sigma(I)$ were used to solve and refine the structure to R = 0.0445 and $R_w = 0.0516$, 210 leastsquares parameters, GOF = 1.02.

22 and **23** were isolated in 55 and 91% yields, respectively.

Well-formed crystals suitable for X-ray analysis were obtained after crystallization of complex 16 from a hexane/diethyl ether mixture.13 The structure undoubtedly confirmed the regioselectivity of the reaction. Interestingly, the η^5 -cyclohexadienyl moiety did not exhibit the expected classical five-coplanar-carbon geometry (C₁, C₂, C₃, C₄, C₅)¹⁴ with one carbon, C₆, lying out of this plane. Rather, we observed that two carbons, C_6 and C_2 , are located out of the plane formed by the other C₁, C₃, C₄, and C₅ sp² carbons and away from the metal center. Indeed, the $Mn-C_1/C_3/C_4/C_5$ bond lengths lie between 2.15 and 2.20 Å, whereas Mn-C₂ and Mn-C₆ bonds manifest lengths of 2.38 and 2.48 Å, respectively. The dihedral angles between the $C_1, C_3, C_4, C_5/$ C1, C2, C3 and C1, C3, C4, C5/C1, C5, C6 planes reach 19.8 and 25.2°, respectively.

This could be better described as a π -allyl C₃, C₄, C₅ σ -alkyl C₁ derivative¹⁵ which might be represented by a "high-heeled-shoe" conformation, as shown in Figure 1. For the moment it is difficult to explain this unexpected structure, and we can only suggest that the thiophenyl residue might play an important role in forcing the complex to adopt this conformation in the solid state.

In conclusion, we have successfully developed a general palladium-catalyzed approach to new (η^5 -cyclohexadienyl)Mn(CO)₃ complexes bearing electron-donating groups at the terminus of the π -system. These complexes cannot be synthesized using classical methods. An X-ray structure of one of the C₁-substituted cyclohexadienyl manganese compounds shows an un-

(15) One of the reviewers mentioned an interesting example of a σ, π -cyclohexenyl complex of Mn (Pike, R. D.; Ryan, W. J.; Lennhoff, N. S.; Van Epp, J.; Sweigart, D. A. *J. Am. Chem. Soc.* **1990**, *112*, 4798) presenting a striking resemblance to that described here. However, in this structure, C2 atom is an sp³ carbon not bonded to the Mn atom.



Figure 1. X-ray structure of complex **16** with the numbering scheme and the "high heeled-shoe" analogy. Thermal ellipsoids are presented at the 30% probability level. Selected bond lengths (Å) and angles (deg): $Mn-C_1 = 2.17$, $Mn-C_2 = 2.38$, $Mn-C_3 = 2.15$, $Mn-C_4 = 2.20$, $Mn-C_5 = 2.17$, $Mn-C_6 = 2.46$; $[C_1C_2C_3]/[C_1C_3C_4C_5] = 19.8$, $[C_1C_5C_6]/[C_1C_3C_4C_5] = 25.5$.

usual conformation which contravenes the "rule" of planarity among the five sp² carbons. Further investigations on the structural and chemical properties of such compounds are currently in progress.

Acknowledgment. We thank the CNRS, EU Training and Mobility of Researchers Program (Contract No. ERBFMRXCT-CT98-0166), and the EU COST D14 Program for financial support and the Ministère de l'Education Nationale et de la Recherche for a MENRT grant to A.A.

Supporting Information Available: Text giving spectral data for the new compounds and tables of crystal data, atomic coordinates, and bond distances and angles for complex **16**. This material is available free of charge via the Internet at http://pubs.acs.org.

OM0202773

⁽¹⁴⁾ For X-ray studies of (η⁵-cyclohexadienyl)Mn(CO)₃ complexes, see for example: (a) Lee, T. Y.; Bae, H. K.; Chung, Y. K.; Hallowse, W. A.; Sweigart, D. A. *Inorg. Chim. Acta* **1994**, *224*, 147. (b) Dullaghan, C. A.; Carpenter, G. B.; Sweigart, D. A. *Chem. Eur. J.* **1997**, *3*, 75. (c) Balssa, F.; Gagliardini, V.; Lecorre-Susanne, C.; Rose-Munch, F.; Rose, E.; Vaissermann, J. *Bull. Chem. Soc. Fr.* **1997**, *134*, 537. (d) Renard, C.; Valentic, R.; Rose-Munch, F.; Rose, E. *Organometallics* **1998**, *17*, 1587. (e) Rose, E.; Lecorre-Susanne, C.; Rose-Munch, F.; Renard, C.; Gagliardini, V.; Tedji, F.; Vaissermann, J. *Eur. J. Inorg. Chem.* **1999**, *3*, 421. (f) Son, S. U.; Paik, S. J.; Park, K. H.; Lee, Y. A.; Lee I. S.; Chung, Y. K., Sweigart, D. A. *Organometallics* **2002**, *21*, 239.