

PII: S0040-4020(97)00693-5

Synthesis and Cytotoxic Activity of Steroid-Anthraquinone Hybrids

Francesco De Riccardis*, Irene Izzo, Marcello Di Filippo and Guido Sodano*

Dipartimento di Chimica, Università di Salerno, 84081 Baronissi, Salerno - Italy

Fulvio D'Acquisto and Rosa Carnuccio

Dipartimento di Farmacologia Sperimentale, Università di Napoli "Federico II", Via Montesano 49, Napoli - Italy

Abstract: Synthesis of cytotoxic steroidal derivatives containing a quinone moiety is described. The synthetic strategy is based on an unsual A + CD -> ABCD Diels-Alder approach which generates 9β -H cholestane analogs. The adducts formed are efficiently aromatized in basic media to give steroid-anthaquinones hybrids showing interesting cytotoxic activity on four tumor cell lines. © 1997 Elsevier Science Ltd.

Novel potentially bioactive compounds can be designed combining in a single molecule different classes of natural products. This approach has been succesfully applied *e.g.* in the design of hybrid molecules containing a neotropsin-type minor groove binder linked to neocarzionostatin chromophore analogs,¹ or by combination of classic β -lactam antibiotics and gyrase inhibitors, which give rise to extremely potent antibiotics.² Other examples of potentially biologically active hybrid compounds synthetized are: taxamicins,³ lactendiynes,⁴ calichearubicins,⁵ retinoid-anthraquinones hybrids,⁶ β -lactam-nucleosides.⁷ Attention has also been paid to the synthesis of drug-steroid hybrids, as for example estramycins,⁸ steroidal polyamines,⁹ a cholesterol-baccatin III hybrid¹⁰ and opioid-steroid hybrids.¹¹

In consideration of the powerful antineoplastic activity of the anthracycline antibiotics great efforts have been made in the design and synthesis of analogs of these compounds presenting low molecular complexity and similar mechanism of action. Starting from doxorubicin (Fig. 1, 1), the most commonly used intercalating agent in the treatment of cancer,¹² new synthetic anthracenediones have been developed. Mitoxantrone (2), in particular, has shown clinical effectiveness against human malignancies.¹³



Figure 1

10871

Because of the importance of steroids as vectors for biological activities (see, for example, the potent antiviral colasane)¹⁴ as a part of a development of new approaches to compounds related to steroids, we designed, synthesized and evaluated the cytotoxic activity of a novel class of steroid-anthraquinone hybrids of general formula 3 (Fig. 2). The new compounds were synthesized in two series differing for the side chains. The number and position of hydroxyl groups were also varied in order to gain information on the requirements for cytotoxic activity.



Figure 2

For the synthesis of the steroid-anthraquinone hybrids we took advantage of a recent application of the rare¹⁵ A + CD -> ABCD Diels-Alder approach for the construction of cholestane analogs. Using this strategy we synthesized remarkable cholestane analogs, such as a C_2 symmetric *bis*-cholestane derivative and a fullerene adduct.¹⁶ The CD ring fragments used in the Diels-Alder reaction were dienes **4a** and **4b**, the latter recently prepared for the synthesis of (17R)-17-methylincisterol,¹⁷ which were obtained respectively from vitamin D₃ and vitamin D₂.





We report here full experimental details on the preparation of 4a and 4b, their reaction with a series of naphthoquinones and the aromatization of the resulting adducts to give antiproliferative steroid-anthraquinone hybrids.

Dienes 4a and 4b were obtained respectively from vitamin D_2 (65% overall yield) and vitamin D_3 (55% overall yield), through a general protocol reported in Scheme 1.



Sc	heme	1	

The ready kinetic enolization of Grundmann's ketones¹⁸ **5a** and **5b** was achieved at -78° C using the bulky base sodium *bis*(trimethylsilyl)amide. The resulting enolates were quenched with Nphenyltrifluoromethanesulfonimide¹⁹ to afford the triflates **6a** and **6b**. Stille reaction²⁰ between **6a** and **6b** and vinyltributyltin, catalyzed by tetrakis(triphenylphosphine)palladium(0) gave the dienes **4a** and **4b**.

The reaction of both 4a and 4b with different anthraquinones in refluxing toluene afforded stereochemically pure 9β -H unnatural cholestane analogues 7-10 (Table 1) as the sole detectable Diels-Alder adducts. The stereochemistry of the reaction was elucidated by comparison of the NMR data of 7b-10 with those of the adduct 7a, whose structure was secured by X-ray diffraction analysis.¹⁶



Table 1

Aromatization of adducts 7-10 was easily accomplished by the action of air in presence of DBU (1,8-diazabicyclo[5.4.0]undec-7-ene)²¹ to afford compounds 11-14, hydrogen peroxyde being concevaibly the other product.

The reaction of 4a,b with 5-hydroxynaphthoquinone afforded the two pairs 8a-9a and 8b-9b. The single components of each pair were separated by silica gel column chromatography and the structures were assigned by comparison of the ¹³C chemical shift values of the carbonyl carbons in the compounds 12a,b and 13a,b with those of 14a,b. In fact, in the ¹³C NMR spectrum of 14a the two carbonyl resonances at δ 190.1 and 187.1 were assigned to C-4 and C-1, respectively, since the latter suffers the γ -effect due to the C-11. In compound 12a the same carbonyl carbons resonate at δ 191.8 and 182.9, while in 13a the chemical shift values were 188.9 and 185.0, making clear that the variation of the chemical shift values at both carbons in the pair 12a-13a, as compared to 14a, was due to the presence or the absence of the *peri*-OH group.

Compounds 11-14 were tested *in vitro* on four tumor cell lines for the evaluation of their antiproliferative activity^{22,23} in comparison with doxorubicin. The results are reported in Table 2 as IC_{50} (concentration of compound expressed in µg/ml required to inhibit the cellular growth by 50% after 24, 48 and 72 hours of drug exposure).

Table 2. In vitro cytotoxic activity (IC₅₀, μ g/ml) of compounds 11-14 and doxorubicin on J774 (murine monocyte/macrophage), GM7373 (bovine aortic endothelial), IGR-1 (human melanoma) and P388 (murine leukemia) cell lines.^a

Cell-line h	11a	11b	12a	12b	13a	13b	14a	14b	doxorub.
J774									
24	n. d.	100.1±8.2	n. d.	30.7±5.1	117.6±9.6	n. d.	143.7±2.3	19.7±4.1	86.3±9.3
48	112.6±14.9	25.3±5.4	95.9±2.1	10.1±1.1	62.5±4.3	43.4±3.4	133.3±21.7	10.9±1.3	54.2±7.1
72	83.3±7.2	16.6±2.1	80.9 ±6 .4	0.35±0.01	25.9±7.2	35.4±4.1	80.1±4.3	7.7±1.1	12.1±3.9
GM7373									
24	n. d.	n. d.	n. d.	n. d.	n. d.	n. d.	n. d.	n. d.	69.7±9.4
48	71.4±2.9	35.0±10.3	85.3±4.2	22.3±1.7	50.8±6.6	54.7±4.2	44.1±5.2	16.3±2.1	17.6±3.5
72	35.7±5.5	12.5±2.4	8.8±2.5	0.55±0.08	30.7±5.2	33.3±3.1	31.7±2.3	8.8±2.2	4.6±2.9
IGR-1									
24	n. d.	n. d.	n. d.	87.0±6.2	n. d.	150.9±19.3	n. d.	46.7±4.1	22.9 ±6 .4
48	n. d.	44.1±5.3	111.1±11.3	30.7±5.2	19.7±3.3	87.0±14.1	231.3±22.6	10.9±2.3	11.5±1.4
72	83.3±10.3	31.7±4.3	81.3±6.4	9.5±1.9	12.5±2.4	50.6±3.4	66.6±12.3	8.2±1.3	0.34±0.09
P388									
24	n. d.	n. d.	n. d.	43.4±3.4	n. d.	n. d.	n. d.	33.3±2.1	52.3±7.3
48	274.0±12.3	181.1±9.2	164.0±14.1	34.5±4.1	75.0±10.3	125.0±15.8	174.0±5.7	23.2±1.6	10.3±3.9
72	132.0±7.1	111.1±5.3	77.1±6.3	18.5±1.3	50.0±8.1	133.3±21.6	153.8±4.1	12.5±0.9	2.1±0.9

n.d.; activity not detectable. Bolded data refer to IC_{50} lower than that found for doxorubicin

^a The results are expressed as mean ± S.E.M. of three separate experiments in triplicate.

The results show that compounds with the ergosterol side chain (11b-14b) were more cytotoxic than those with the saturated side chain (11a-14a). In particular, the most active compounds were those possessing a C-3' hydroxyl group, *e.g.* 12b and 14b, which show cytotoxic activities comparable to those of doxorubicin.

In conclusion the *in vitro* data appears quite promising and warrant further investigation to fully assess the potentials of these hybrids.

Acknowledgements. This work has been supported by MURST (40%) and CNR (Rome). The mass spectra were obtained from the "Servizio di Spettrometria di Massa del CNR e dell'Università di Napoli"; the staff is gratefully acknowledged. We thank Dr. Gianluca Giorgi ("Centro Interdipartimentale di Analisi e Determinazioni Strutturali" of the University of Siena) for accurate high resolution mass spectra. We wish to thank Dr. Paolo Lombardi (Menarini Ricerche Sud, Pomezia, Roma), Prof. Manlio Palumbo (University of Padova) for helpful discussions and miss Rossella De Caro for valuable aid in the experimental work.

EXPERIMENTAL SECTION

General techniques

All reactions were carried out under a dry argon atmosphere using freshly distilled solvents unless otherwise noted. Tetrahydrofuran was distilled from sodium and benzophenone. Toluene and methylene chloride were distilled from calcium hydride. Glassware was flame dried (0.05 torr) before use. When necessary, compounds were dried by azeotropic removal of water with toluene under reduced pressure. Reactions were monitored by thin layer chromatography (TLC) on Merck silica gel plates (0.25 mm) and visualized using UV light, spraying with H_2SO_4 -Ce(SO₄)₂ solution and drying. Reaction temperatures were measured externally. Flash chromatography was performed on Merck silica gel (60, particle size: 0.040-0.063 mm). Yields refer to chromatographically and spectroscopically (¹H-NMR) pure materials.

NMR spectra were recorded in CDCl₃ solutions on a Bruker AM-250 spectrometer at ambient temperature. Chemical shifts are reported relative to the residual solvent peak (CHCl₃: δ =7.26, ¹³CDCl₃: δ =77.0). Optical rotations were recorded in CHCl₃ solutions on a JASCO DIP-370 or JASCO DIP-1000 polarimeters. Mass spectra (E.I., 70 eV) were recorded on a VG TRIO 2000 mass spectrometer. High resolution mass spectra were recorded on a VG 70-250SE (70eV, source temp.: 200°C, resolution: 5000 m/ Δ m, 10% valley). IR were recorded on a Nicolet FT 5DXB spectrometer. UV-Vis spectra were recorded in dioxane on a Varian DMS 90 spectrometer. For the cytotoxic assays the optical density (OD) of the wells were measured with a microplate spectrophotometer (Titertek Multiscan MCC/340) equipped with a 620 nm filter. All the reagents for cell culture were from Cellbio; MTT and doxorubicin were purchased from Sigma.

Preparation and selected data of compounds.

Ketones 5a and 5b.18

5b. To a -20°C cooled solution of vitamin D_2 (5.00 g, 13.1 mmol) in absolute ethanol (500 ml) was slowly added a solution of KMnO₄ (4.20 g, 26.5 mmol) in distilled water (125 ml). The resulting mixture was stirred for additional 15 min, and then warmed at 40°C for 30 min. Filtration through a shorth path af celite and washing of the solid with absolute ethanol afforded a yellow solution. Removal of the solvent in vacuo afforded a residue which was flash chromatographed (silica gel, 2-3% methanol in choroform) to give the corresponding triol (3.25 g, 65%) as a white solid. $[\alpha]_D = +63.4^{\circ}$ (c=0.44, CHCl₃); ¹H NMR δ : 0.80 (3H, s, Me-18), 0.81 (3H, d, J= 6.6 Hz, Me-26), 0.82 (3H, d, J= 6.6 Hz, Me-27), 0.90 (3H, d, J= 6.6 Hz, Me-28), 0.97 (3H, d, J= 6.6 Hz, Me-21), 2.19 (1H, ddd, J= 13.5, 12.0, 1.6 Hz, H-4), 2.44 (1H, ddd, J= 13.0, 4.0, 3.9 Hz, H-1), 2.58 (1H, bdd, J=12.0, 3.5 Hz, H-4'), 3.78 (1H, m, H-3), 4.91 (1H, d, J= 9.7 Hz, H-7), 4.92 (1H, bs, H-19), 5.01 (1H, bs, H-19'), 5.16 (2H, m, H-22 and H-23), 5.56 (1H, dd, J=9.7, 1.6 Hz, H-6); ¹³C NMR δ : 13.0, 17.1, 19.6, 20.0, 20.4, 20.6, 21.8, 27.7, 33.1, 33.2, 36.0, 37.6, 39.9, 40.0, 42.8, 44.0, 46.5, 57.3, 59.9, 70.5, 70.7, 74.9, 111.4, 125.2, 132.1, 135.4, 140.7, 145.3; I.R. (film) v_{max} 3282, 1045 cm⁻¹, MS, *m/z*: 412 (M⁺+H₂0), 154, 136.

To a -15°C cooled solution of triol (0.20 g, 0.46 mmol) in dry methylene chloride (4 ml), $Pb(OAc)_4$ (0.24 g, 0.55 mmol) was added in portions. The resulting mixture was stirred for 20 min at the same temperature and then without the cooling bath for additional 20 min. Filtration through a short column of celite-silica gel (1:1) and

washing of the solid with methylene chloride afforded a solution which was concentrated in vacuo. The residue was flash chromatographed (silica gel; 5-30% ethyl acetate in petroleum ether) to give **5b** (0.107 g, 93% yield) as a colorless oil. $[\alpha]_D$ = +11.6° (c=1.01, CHCl₃); ¹H NMR & 0.61 (3H, s, Me-18), 0.78 (3H, d, J= 6.6 Hz, Me-26), 0.79 (3H, d, J= 6.6 Hz, Me-27), 0.87 (3H, d, J= 6.6 Hz, Me-28), 1.00 (3H, d, J= 6.6 Hz, Me-21), 2.41 (1H, dd, J= 10.9, 7.6 Hz, H-14), 5.18 (2H, m, H-22 and H-23); ¹³C NMR & 12.6, 17.5, 19.0, 19.6, 19.9, 20.9, 24.0, 27.7, 33.0, 38.8, 39.8, 40.9, 42.8, 49.7, 56.5, 62.0, 132.4, 134.8, 211.9; I.R. (film) v_{max} 1717 cm⁻¹; MS, *m/z*: 276 (M⁺), 261, 233, 205, 179, 151.

5a. To a -20°C cooled solution of vitamin D₃ (15.0 g, 39.0 mmol) in absolute ethanol (1350 ml) was slowly added a solution of KMnO₄ (12.4 g, 78.5 mmol) in distilled water (450 ml). The resulting mixture was stirred for additional 75 min, and then warmed at 40°C for 30 min. Filtration through a shorth path af celite and washing of the solid with absolute ethanol afforded a yellow solution. Removal of the solvent in vacuo afforded a residue which was dried by azeotropic removal of water with toluene under reduced pressure and dissolved in dry methylene chloride (400 ml). This solution was cooled to -15°C and Pb(OAc)₄ (24.4 g, 55.0 mmol) was added in four portions. The resulting mixture was stirred for 15 min at the same temperature and without the cooling bath for additional 60 min. Filtration through a short column of celite-silica gel (1:1) and washing of the solid with methylene chloride afforded a solution which was concentrated in vacuo. The residue was flash chromatographed (silica gel, 5-30% ethyl acetate in petroleum ether) to give **5a** (8.51 g, 82.5% yield) as a colorless oil. $[\alpha]_D = +8.9^\circ$ (c=1.59, CHCl₃); Lit.:^{18b} $[\alpha]_D = +7.9^\circ$ (c=2.80, CHCl₃); ¹H NMR δ : 0.62 (3H, s, Me-18), 0.86 (6H, d, J= 6.6 Hz, Me-26 and -27), 0.94 (3H, d, J= 6.6 Hz, Me-21), 2.44 (1H, dd, J= 10.9, 7.5 Hz, H-14); MS, *m/z*: 264 (M⁺), 221, 179, 151.

Vinyl triflates 6a and 6b.19b

6b. To a solution of sodium *bis*(trimethylsily)amide (0.75 ml, 1.0 M in THF, 0.75 mmol) at -78°C **5b** (85.0 mg, 0.30 mmol) dissolved in dry THF (1.0 ml) was added. After 1 h, N-phenyltrifluoromethanesulfonimide (260 mg, 0.72 mmol) was added to the resulting enolate. After 20 min the reaction mixture was slowly warmed at r.t. and stirred for 2 h. The reaction was quenched with water, concentrated in vacuo to remove the excess of THF, and extracted with petroleum ether. The organic phase was dried (Na₂SO₄), filtered and concentrated in vacuo. The residue was flash chromatographed (silica gel, petroleum ether) to give **6b** (115 mg, 93% yield) as a colorless oil. $[\alpha]_D$ = +85.8° (c=2.31, CHCl₃); ¹H NMR δ : 0.80 (3H, s, Me-18), 0.84 (3H, d, J= 6.6 Hz, Me-26), 0.86 (3H, d, J= 6.6 Hz, Me-27), 0.94 (3H, d, J= 6.6 Hz, Me-28), 1.05 (3H, d, J= 6.6 Hz, Me-21), 2.31 (2H, m, H₂-11), 2.50 (1H, m, H-14), 5.18 (1H, dd, J= 15.2, 8.6 Hz, H-22 or H-23), 5.30 (1H, dd, J= 15.2, 7.0 Hz, H-23 or H-22), 5.58 (1H, d, J= 7.4, 3.7 Hz, H-9); ¹³C NMR δ : 11.4, 17.5, 19.9, 20.8, 21.5, 21.7, 23.8, 33.1, 34.8, 40.3, 42.9, 45.1, 50.3, 54.2, 118.6 (q, J= 312 Hz), 132.9, 135.0, 149.9; MS, *m/z*: 410 (M⁺+2), 408 (M⁺), 365, 327, 283, 259.

6a^{19b} was synthesized from **5a** as described above for **6b** (85% yield, colorless oil). $[\alpha]_D = +18.2^{\circ}$ (c=1.02, CHCl₃); ¹H NMR δ : 0.78 (3H, s, Me-18), 0.88 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 0.94 (3H, d, J= 6.6 Hz, Me-21), 2.40 (2H, m H-11), 2.48 (1H, m, H-14), 5.58 (1H, dd, J= 7.4, 3.7 Hz, H-9); MS, *m/z*: 398 (M⁺+2), 396 (M⁺).

Dienes 4a and 4b

4b. To a suspention of $Pd(PPh_3)_4$ (85 mg, 0.07 mmol), LiCl (475 mg, 11.3 mmol) and vinyltributyltin (0.85 ml, 2.94 mmol), a solution of **6b** (1.0 g, 2.45 mmol) in dry THF (10 ml) was slowly added. The reaction

mixture was stirred at reflux for 2 h and then quenched with water (10 ml), concentrated in vacuo to remove the excess of THF and extracted with petroleum ether. The organic phase was washed with water, a 10% aqueous solution of NH₄OH and brine. The organic phase was dried (Na₂SO₄), filtered and concentrated in vacuo. The residue was flash chromatographed (silica gel, petroleum ether) to give **4b** (696 mg, 98% yield) as a colorless oil. $[\alpha]_D$ = -11.5° (c=2.35, CHCl₃); ¹H NMR & 0.70 (3H, s, Me-18), 0.84 (3H, d, J= 6.6 Hz, Me-26), 0.85 (3H, d, J= 6.6 Hz, Me-27), 0.93 (3H, d, J= 6.6 Hz, Me-28), 1.06 (3H, d, J= 6.6 Hz, Me-21), 4.84 (1H, d, J= 11.2, H-6), 5.20 (1H, d, J= 17.2 Hz, H-6'), 5.21 (2H, m, H-23 and H-24), 5.67 (1H, m, H-9), 6.20 (1H, dd, J= 17.2, 11.2 Hz, H-7); ¹³C NMR & 11.3, 17.4, 19.5, 19.8, 20.9, 23.7, 24.6, 28.7, 32.9, 35.6, 40.2, 42.5, 42.7, 49.7, 53.7, 111.3, 125.7, 131.8, 135.5, 138.0, 138.7; MS, *m/z*: 286 (M⁺), 259, 161.

4a was synthesized from 6a as described above for 4b (93% yield, colorless oil). $[\alpha]_D = -21.0^\circ$ (c=0.86, CHCl₃); ¹H NMR δ: 0.78 (3H, s, Me-18), 0.89 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 0.96 (3H, d, J= 6.6 Hz, Me-21), 4.83 (1H, d, J= 11.2, H-6), 5.20 (1H, d, J= 17.2 Hz, H-6'), 5.68 (1H, m, H-9), 6.20 (1H, dd, J= 17.2, 11.2 Hz, H-7); ¹³C NMR δ: 11.2, 18.9, 22.5, 22.8, 23.9, 24.0, 24.8, 28.0, 28.7, 35.9, 36.1, 36.2, 39.5, 42.7, 49.8, 54.0, 111.4, 125.8, 138.2, 138.8; MS, *m/z*: 274 (M⁺).

Diels-Alder adducts 7-10

General procedure: diene (1.0 mmol) and dienophile (1.0 mmol) were dissolved in anhydrous toluene (3.0 ml) and heated at 100°C until the reaction was complete (5-12 h, TLC analysis). Concentration under reduced pressure gave a residue which was flash chromatographed (silica gel) to give adducts **7-10**.

7a was eluted with petroleum ether - toluene 1:1 (98% yield) as a colorless solid and recrystallized from ethyl acetate; m. p.: 159 - 160°C. $[\alpha]_D$ = +190.9° (c=0.49, CHCl₃); ¹H NMR & 0.71 (3H, s, Me-18), 0.86 (6H, d, J= 6.5, Me-26 and Me-27), 0.91 (3H, d, J= 5.8 Hz, Me-21), 3.40 (2H, m, H-5 and H-10), 5.06 (1H, m, H-7), 7.70 (2H, dd, J= 5.8, 3.3 Hz, H-3' and H-6'), 7.90 (1H, dd, J= 5.8, 3.3 Hz, H-4' or H-5'), 8.03 (1H, dd, J= 5.8, 3.3 Hz, H-5' or H-4'); ¹³C NMR & 18.4 (C-18), 18.8 (C-21), 22.5, 22.8 (C-26, C-27), 22.8 (C-11), 23.9 (C-23), 24.5 (C-15), 27.1 (C-6), 28.0 (C-25), 28.7 (C-16), 35.9 (C-22), 36.1 (C-20), 36.7 (C-9), 37.7 (C-12), 39.4 (C-24), 41.6 (C-13), 49.2 (C-14), 50.7 (C-5), 50.8 (C-10), 57.1 (C-17), 114.3 (C-7), 126.2, 127.0 (C-3', C-6'), 132.5, 135.9 (C-2, C-3), 133.8, 134.2 (C-4', C-5'), 142.0 (C-8), 198.0 (C-1), 199.2 (C-4); UV: λ = 210-240 (ϵ = 9654-9221), λ_{max} = 257 (ϵ = 6311), λ = 295-307 (ϵ = 1412-1175); MS, *m/z*: 432 (M⁺), 417, 319.

7b was eluted with petroleum ether - toluene 1:1 (71% yield) as a colorless solid. $[\alpha]_D = +109.8^{\circ}$ (c=0.21, CHCl₃); ¹H NMR δ : 0.72 (3H, s, Me-18), 0.82 (3H, d, J= 6.6 Me-26), 0.83 (3H, d, J= 6.6, Me-27), 0.92 (3H, d, J= 6.6 Hz, Me-28), 1.02 (3H, d, J= 6.6 Hz, Me-21), 3.40 (2H, m, H-5 and H-10), 5.08 (1H, m, H-7), 5.20 (2H, m, H-22 and H-23), 7.70 (2H, dd, J= 5.8, 3.3 Hz, H-3' and H-6'), 7.90 (1H, dd, J= 5.8, 3.3 Hz, H-4' or H-5'), 8.02 (1H, dd, J= 5.8, 3.3 Hz, H-5' or H-4'); MS, m/z: 444 (M⁺).

8a, 9a were eluted with petroleum ether - ethyl acetate - acetic acid 99:1:0.1 as yellow solids in 73% (8a) and 13% (9a) yield, respectively.

8a: $[\alpha]_{D}$ = +220.0° (c=0.45, CHCl₃); ¹H NMR δ : 0.71 (3H, s, Me-18), 0.87 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 0.92 (3H, d, J= 5.8 Hz, Me-21), 3.38 (2H, m, H-5 and H-10), 5.08 (1H, m, H-7), 7.22 (1H, dd, J= 7.5, 1.3 Hz, H-4'), 7.53 (1H, dd, J= 7.5, 1.3 Hz, H-6') 7.60 (1H, t, J= 7.5 Hz, H-5'), 11.90 (1H, s, OH); ¹³C NMR δ : 18.4 (C-18), 18.8 (C-21), 22.5, 22.7 (C-26, C-27), 22.8 (C-11), 23.9 (C-23), 24.8 (C-15), 26.9 (C-6), 28.0 (C-25), 28.7 (C-16), 35.9 (C-22), 36.1 (C-20), 37.0 (C-9), 37.7 (C-12), 39.4 (C-24), 41.7 (C-13), 49.2 (C-14), 49.9 (C-5), 50.3 (C-10), 57.1 (C-17), 114.5 (C-7), 117.7 (C-3), 118.4, 124.0, 136.4

(C-4', C-5', C-6'), 132.7 (C-2), 142.3 (C-8), 160.8 (C-3'), 198.6 (C-1), 205.3 (C-4); UV: λ = 200-233 (ε= 8912-8710), λ_{max} = 260 (ε= 5754), λ = 323-348 (ε= 6166-6761); MS, *m/z*: 448 (M⁺), 335.

9a: $[\alpha]_{D}$ = +216.7° (c=0.24, CHCl₃); ¹H NMR δ : 0.70 (3H, s, Me-18), 0.87 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 0.92 (3H, d, J= 5.8 Hz, Me-21), 3.36 (2H, m, H-5 and H-10), 5.08 (1H, m, H-7), 7.22 (1H, d, J= 8.4 Hz, H-5'), 7.39 (1H, d, J= 7.3 Hz, H-3'), 7.60 (1H, dd, J= 8.4, 7.3 Hz, H-4') 12.02 (1H, s, OH); ¹³C NMR δ : 18.4 (C-18), 18.8 (C-21), 22.5, 22.8 (C-26, C-27), 22.8 (C-11), 23.9 (C-23), 24.4 (C-15), 28.0 (C-6), 28.0 (C-25), 28.7 (C-16), 35.9 (C-22), 36.1 (C-20), 36.4 (C-9), 37.6 (C-12), 39.5 (C-24), 41.7 (C-13), 49.2 (C-14), 50.1 (C-5), 50.7 (C-10), 57.2 (C-17), 114.1 (C-7), 115.1 (C-2), 117.5, 123.4, 136.9 (C-3', C-4', C-5'), 136.1 (C-3), 142.0 (C-8), 162.1 (C-6'), 197.0 (C-4), 206.4 (C-1); UV: λ = 200-230 (ϵ = 9332-8511), λ_{max} = 260 (ϵ = 5880), λ = 339-346 (ϵ = 5623-6309); MS, *m/z*: 448 (M⁺), 335.

8b, 9b were isolated as above as yellow solids in 64% and 10% yield, respectively.

8b: $[\alpha]_{D}$ = +85.2° (c=0.64, CHCl₃); ¹H NMR δ : 0.72 (3H, s, Me-18), 0.82 (3H, d, J= 6.6, Me-26), 0.83 (3H, d, J= 6.6, Me-27), 0.91 (3H, d, J= 6.6 Hz, Me-28), 1.02 (3H, d, J= 6.6 Hz, Me-21), 3.37 (2H, m, H-5 and H-10), 5.08 (1H, m, H-7), 5.20 (2H, m, H-22 and H-23), 7.22 (1H, dd, J= 7.5, 1.3 Hz, H-4'), 7.53 (1H, dd, J= 7.5, 1.3 Hz, H-6') 7.60 (1H, t, J= 7.5 Hz, H-5'), 11,88 (1H, s, OH); ¹³C NMR δ : 17.6, 18.9, 19.6, 19.9, 20.6, 22.7, 24.7, 26.7, 29.1, 33.0, 37.0, 37.4, 40.6, 41.5, 42.8, 49.3, 49.9, 50.3, 56.9, 114.6, 117.7, 118.5, 124.1, 132.2, 132.7, 135.3, 136.4, 142.2, 160.8, 198.6, 205.3; MS, *m/z*: 460 (M⁺), 335, 317, 203.

9b: $[\alpha]_{D}$ = +119.7° (c=0.15, CHCl₃); ¹H NMR δ : 0.73 (3H, s, Me-18), 0.82 (3H, d, J = 6.6, Me-26), 0.83 (3H, d, J = 6.6, Me-27), 0.92 (3H, d, J = 6.6 Hz, Me-28), 1.02 (3H, d, J = 6.6 Hz, Me-21), 3.37 (2H, m, H-5 and H-10), 5.08 (1H, m, H-7), 5.20 (2H, m, H-22 and H-23), 7.21 (1H, d, J= 8.4 Hz, H-5'), 7.38 (1H, d, J= 7.3 Hz, H-3'), 7.60 (1H, dd, J= 8.4, 7.3 Hz, H-4') 12.01 (1H, s, OH); ¹³C NMR δ : 17.6, 18.9, 19.6, 19.9, 20.6, 22.8, 24.3, 27.9, 29.1, 33.0, 36.3, 37.4, 40.6, 41.5, 42.8, 49.3, 50.0, 50.6, 56.9, 114.6, 115.1, 117.5, 123.4, 132.1, 135.3, 136.1, 136.9, 141.9, 162.1, 197.0, 206.4; MS, *m/z*: 460 (M⁺), 335, 317, 203.

10a was eluted as a yellow solid with petroleum ether - ethyl acetate - acetic acid 99:1:0.1 in 81% yield. $[\alpha]_{D}$ = +280.0° (c=0.38, CHCl₃); ¹H NMR δ : 0.72 (3H, s, Me-18), 0.88 (6H, d, J=6.6 Hz, Me-26 and Me-27) 0.92 (3H, d, J= 5.8 Hz, Me-21), 3.33 (2H, m, H-5 and H-10), 5.12 (1H, m, H-7), 7.22 (2H, s, H-4' and H-5'), 11.69 (1H, s, OH), 11.88 (1H, s, OH); ¹³C NMR δ : 18.4 (C-18), 18.8 (C-21), 22.7, 22.8 (C-26, C-27), 22.8 (C-11), 23.9 (C-23), 24.7 (C-15), 27.7 (C-6), 28.0 (C-25), 28.7 (C-16), 35.9 (C-22), 36.1 (C-20), 36.7 (C-9), 37.6 (C-12), 39.4 (C-24), 41.6 (C-13), 49.2 (C-14), 49.3 (C-5), 49.4 (C-10), 57.1 (C-17), 114.1 (C-7), 112.2, 114.7 (C-2, C-3), 127.8, 128.5 (C-4', C-5') 142.0 (C-8), 154.2, 155.7 (C-3', C-6'), 203.8 (C-1), 205.2 (C-4); UV: λ = 212-245 (ε = 15849-14125), λ_{max} = 260 (ε = 5888), λ = 394 (ε = 8318) λ (s)= 413 (ε = 6760); MS, *m/z*: 464 (M⁺), 446, 351.

10b. Yellow solid isolated as above in 66% yield. ¹H NMR δ : 0.72 (3H, s, Me-18), 0.82 (3H, d, J= 6.6 Me-26), 0.83 (3H, d, J= 6.6, Me-27), 0.91 (3H, d, J= 6.6 Hz, Me-28), 1.01 (3H, d, J= 6.6 Hz, Me-21), 3.32 (2H, m, H-5 and H-10), 5.10 (1H, m, H-7), 5.20 (2H, m, H-22 and H-23), 7.20 (2H, s, H-4' and H-5'), 11.66 (1H, s, OH), 11.85 (1H, s, OH); MS, *m*/*z*: 476 (M⁺).

Steroid-anthraquinone hybrids 11-14

General procedure: to a solution of Diels-Alder adducts 7-10 (1 mmol) in methylene chloride (15 ml), DBU (1,8-diazabicyclo[5.4.0]undec-7-ene) was added (1 mmol for 7, 2.5 mmol for 8-10). The resulting dark

solution was stirred for 3 h in an open flask. The reaction was quenched with HCl (0.5 N, 7 ml) and water (10 ml). The organic layer was washed with water and dried (Na_2SO_4) , filtered and concentrated in vacuo. The

residue was flash cromatographed (silica gel) to give hybrids 11-14.

11a. Yellow solid. Eluted with petroleum ether - toluene 1:1; 90% yield. $[\alpha]_D = +27.6^{\circ}$ (c=0.42, CHCl₃); ¹H NMR δ : 0.57 (3H, s, Me-18), 0.88 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 1.01 (3H, d, J= 5.8 Hz, Me-21), 1.74 (1H, dt, J= 14, 8.5 Hz, H-12), 2.13 (2H, m, H₂-15), 2.30 (1H, dt, J= 14.0, 5.7 Hz, H-12), 2.76 (1H, dd, J= 14.2, 6.8 Hz, H-14), 3.53 (2H, dd, J= 8.5, 5.7 Hz, H₂-11), 7,44 (1H, d, J= 7.8 Hz, H-7), 7.74 (2H, m, H-4' and H-5'), 8.19 (1H, d, J= 7.8 Hz, H-6), 8.23 (2H, m, H-3' and H-6'); ¹³C NMR δ : 11.5 (C-18), 18.7 (C-21), 22.5, 22.8 (C-26, C-27), 23.8 (C-23), 24.3 (C-15), 27.4 (C-11), 28.0 (C-25), 28.5 (C-16), 36.0 (C-22), 36.1 (C-20), 37.0 (C-12), 39.4 (C-24), 41.4 (C-13), 52.5 (C-14), 55.6 (C-17), 125.6, 126.4, 127.2, 131.5, 133.2, 133.9 (C-3', C-4', C-5', C-6', C-6, C-7), 130.1, 132.7, 132.9, 135.2 (C-2, C-3, C-5, C-10), 141.2 (C-9), 149.1 (C-8), 183.7 (C-1), 185.7 (C-4); UV: λ = 200-218 (ϵ = 6607), λ_{max} = 266 (ϵ = 14454), λ = 320-328 (ϵ = 3311-3467); MS, *m*/*z*: 428 (M⁺), 315; HRMS, *m*/*z*: 428.2704 (428.2715 calcd for C₃₀H₃₆O₂).

11b. Yellow solid. Eluted with petroleum ether - toluene 1:1; 90% yield. $[\alpha]_D = +26.8^{\circ}$ (c=1.73, CHCl₃); ¹H NMR δ : 0.59 (3H, s, Me-18), 0.85 (3H, d, J= 6.6 Hz, Me-26), 0.86 (3H, d, J= 6.6 Hz, Me-27), 0.95 (3H, d, J= 6.6 Hz, Me-28), 1.11 (3H, d, J= 6.6 Hz, Me-21), 2.74 (1H, dd, J= 14.2, 6.8 Hz, H-14), 3.53 (2H, dd, J= 8.5, 5.7 Hz, H₂-11), 5.25 (2H, m, H-22 and H-23), 7,42 (1H, d, J= 7.8 Hz, H-7), 7.74 (2H, m, H-4'and H-5'), 8.17 (1H, d, J= 7.8 Hz, H-6), 8.23 (2H, m, H-3' and H-6'); MS, *m/z*: 440 (M⁺), 315; HRMS, *m/z*: 440.2710 (440.2715 calcd for C₃₁H₃₆O₂).

12a. Yellow solid. Eluted with petroleum ether - AcOEt - AcOH 95:5:0.1; 75% yield. $[\alpha]_D = +48.5^{\circ}$ (c=0.68, CHCl₃); ¹H NMR δ : 0.56 (3H, s, Me-18), 0.88 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 1.01 (3H, d, J= 5.8 Hz, Me-21), 2.12 (2H, m, H₂-15), 2.30 (1H, dt, J= 14.2, 5.7 Hz, H-12), 2.70 (1H, dd, J= 14.2, 6.8 Hz, H-14), 3.50 (2H, dd, J= 8.5, 5.7 Hz, H₂-11), 7.24 (1H, d, J= 8.4 Hz, H-4'), 7.41 (1H, d, J= 7.9 Hz, H-7), 7.58 (1H, dd, J= 8.4, 7.4 Hz, H-5') 7.72 (1H, d, J= 7.4 Hz, H-6'), 8.13 (1H, d, J= 7.9 Hz, H-6), 12.92 (1H, s,OH); ¹³C NMR δ : 11.4 (C-18), 18.7 (C-21), 22.5, 22.8 (C-26, C-27), 23.8 (C-23), 24.4 (C-15), 27.8 (C-11), 28.0 (C-25), 28.5 (C-16), 36.0 (C-22), 36.1 (C-20), 37.0 (C-12), 39.5 (C-24), 41.3 (C-13), 52.5 (C-14), 55.6 (C-17), 117.4 (C-3), 118.5 (C-4'), 124.3 (C-6'), 125.8 (C-6), 129.6 (C-5), 132.1 (C-7), 132.9 (C-2, C-10), 135.8 (C-5'), 141.8 (C-9), 149.1 (C-8), 162.3 (C-3'), 182.9 (C-1), 191.8 (C-4); UV: λ = 208-252 (ϵ = 13803-13182), $\lambda_{max} = 263$ (ϵ = 25119), λ (sh)= 274 (ϵ = 10000), λ = 360-404 (ϵ = 7244-7413) λ = 384 (ϵ = 7762); MS, *m/z*: 444 (M⁺), 331, 289; HRMS, *m/z*: 444.2662 (444.2664 calcd for C₃₀H₃₆O₃).

12b. Yellow solid. Eluted with petroleum ether - AcOEt - AcOH 95:5:0.1, 85% yield. $[\alpha]_D = +45.2^{\circ}$ (c=0.50, CHCl₃); ¹H NMR δ : 0.59 (3H, s, Me-18), 0.84 (3H, d, J= 6.6 Hz, Me-26), 0.87 (3H, d, J= 6.6 Hz, Me-27), 0.94 (3H, d, J= 6.6 Hz, Me-28), 1.11 (3H, d, J= 6.6 Hz, Me-21), 2.75 (1H, dd, J= 14.2, 6.8 Hz, H-14), 3.54 (2H, m, H₂-11), 5.25 (2H, m, H-22 and H-23), 7.27 (1H, d, J= 8.4 Hz, H-4'), 7.44 (1H, d, J= 7.9 Hz, H-7), 7.61 (1H, dd, J= 8.4, 7.4 Hz, H-5') 7.75 (1H, d, J= 7.4 Hz, H-6'), 8.17 (1H, d, J= 7.9 Hz, H-6), 12.96 (1H, s,OH); MS, *m/z*: 456 (M⁺), 332, 331, 329, 289; HRMS, *m/z*: 456.2669 (456.2664 calcd for C₃₁H₃₆O₃).

13a. Yellow solid. Eluted with petroleum ether - AcOEt - AcOH 95:5:0.1, 61% yield. $[\alpha]_D$ = +47.4° (c=0.19, CHCl₃); ¹H NMR δ : 0.59 (3H, s, Me-18), 0.89 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 1.00 (3H, d, J= 5.8 Hz, Me-21), 2.16 (2H, m, H₂-15), 2.30 (1H, m, H-12), 2.76 (1H, m, H-14), 3.52 (2H, m, H₂-11), 7.24 (1H, d, J= 8.4 Hz, H= 5'), 7.48 (1H, d, J= 7.9 Hz, H-7), 7.65 (1H, dd, J= 8.4, 7.4 Hz, H-4'), 7.76

10881

(1H, d, J= 7.4 Hz, H-3'), 8.22 (1H, d, J= 7.9 Hz, H-6), 12.58 (1H, s, OH); ¹³C NMR δ : 11.5 (C-18), 18.7 (C-21), 22.5, 22.8 (C-26, C-27), 23.8 (C-23), 24.4 (C-15), 27.4 (C-11), 28.0 (C-25), 28.5 (C-16), 36.0 (C-22), 36.1 (C-20), 36.9 (C-12), 39.5 (C-24), 41.5 (C-13), 52.6 (C-14), 55.6 (C-17), 115.7 (C-2), 119.3 (C-5'), 123.0 (C-3'), 125.3 (C-6), 130.2 (C-5), 131.5 (C-7), 132.5 (C-10), 135.3 (C-3), 136.5 (C-4') 141.7 (C-9) 150.0 (C-8), 161.8 (C-6'), 185.0 (C-4), 188.9 (C-1); UV: λ = 200-228 (ϵ = 13833-13148), λ_{max} = 258 (ϵ = 20892), λ (s)= 273 (ϵ = 10471), λ = 398-413 (ϵ = 7079-6918); MS, *m*/*z*: 444 (M⁺), 331, 289; HRMS, *m*/*z*: 444.2661 (444.2664 calcd for C₃₀H₃₆O₃).

13b. Yellow solid. Eluted with petroleum ether - AcOEt - AcOH 95:5:0.1, 85% yield. $[\alpha]_D = +28.6^{\circ}$ (c=0.88, CHCl₃); ¹H NMR δ : 0.59 (3H, s, Me-18), 0.85 (3H, d, J= 6.6 Hz, Me-26), 0.88 (3H, d, J= 6.6 Hz, Me-27), 0.94 (3H, d, J= 6.6 Hz, Me-28), 1.11 (3H, d, J= 6.6 Hz, Me-21), 2.75 (1H, dd, J= 14.2, 6.8 Hz, H-14), 3.53 (2H, m, H₂-11), 5.25 (2H, m, H-22 and H-23), 7.26 (1H, d, J= 8.4 Hz, H= 5'), 7.47 (1H, d, J= 7.9 Hz, H-7), 7.65 (1H, dd, J= 8.4, 7.4 Hz, H-4'), 7.76 (1H, d, J= 7.4 Hz, H-3'), 8.22 (1H, d, J= 7.9 Hz, H-6), 12.61 (1H, s, OH); MS, *m/z*: 456 (M⁺), 331, 332, 290, 289, 275; HRMS, *m/z*: 456.2667 (456.2664 calcd for C₃₁H₃₆O₃).

14a. Red solid. Eluted with petroleum ether - AcOEt - AcOH 90:10:0.1; 90% yield. $[\alpha]_D = + 82.3^{\circ}$ (c=0.53, CHCl₃); ¹H NMR δ : 0.54 (3H, s, Me-18), 0.87 (6H, d, J= 6.6 Hz, Me-26 and Me-27), 1.00 (3H, d, J= 5.8 Hz, Me-21), 2.12 (2H, m, H₂-15), 2.28 (1H, m, H-12), 2.80 (1H, m, H-14), 3.48 (2H, m, H₂-11), 7.20 (2H, s, H= 4' and H= 5'), 7.41 (1H, d, J= 7.9 Hz, H= 7), 8.13 (1H, d, J= 7.9 Hz, H-6), 12.87 (1H, s, OH), 13.12 (1H, s, OH); ¹³C NMR δ : 11.4 (C-18), 18.7 (C-21), 22.5, 22.8 (C-26, C-27), 23.8 (C-23), 24.4 (C-15), 27.8 (C-11), 28.0 (C-25), 28.5 (C-16), 36.1 (C-20, C-22), 37.0 (C-12), 39.4 (C-24), 41.2 (C-13), 52.6 (C-14), 55.6 (C-17), 112.5, 113.9 (C-2, C-3), 125.3 (C-6), 127.9, 129.1 (C-4', C-5'), 129.7 (C-5), 132.0 (C-7), 132.6 (C-10), 142.2 (C-9) 149.8 (C-8), 156.8, 157.2 (C-3', C-6'), 187.1 (C-1), 190.1 (C-4); UV: λ = 200-234 (ϵ = 14125-131804), λ_{max} = 272 (ϵ = 13490), λ (s)= 283 (ϵ = 5370), λ_{max} = 356 (ϵ = 1995), λ = 467-494 (ϵ = 5754-5495); MS, *m*/*z*: 460 (M⁺), 347; HRMS, *m*/*z*: 460.2615 (460.2614 calcd for C₃₀H₃₆O₄).

14b. Red solid. Eluted with petroleum ether - AcOEt - AcOH 90:10:0.1; 90% yield. $[\alpha]_D = + 91.5^{\circ}$ (c=0.60, CHCl₃); ¹H NMR δ : 0.59 (3H, s, Me-18), 0.84 (3H, d, J= 6.6 Hz, Me-26), 0.85 (3H, d, J= 6.6 Hz, Me-27), 0.94 (3H, d, J= 6.6 Hz, Me-28), 1.11 (3H, d, J= 6.6 Hz, Me-21), 2.79 (1H, dd, J= 14.2, 6.8 Hz, H-14), 3.57 (2H, m, H₂-11), 5.25 (2H, m, H-22 and H-23), 7.26 (2H, s, H= 4' and H= 5'), 7.48 (1H, d, J= 7.9 Hz, H= 7), 8.24 (1H, d, J= 7.9 Hz, H-6), 12.95 (1H, s, OH), 13.21 (1H, s, OH); MS, *m*/z: 472 (M⁺), 347, 305, 291; HRMS, *m*/z: 472.2628 (472.2614 calcd for C₃₁H₃₆O₄).

Determination of the cytotoxic activity

Cells. J774 cells (murine monocyte/macrophage cells) were grown in suspension culture, in Techne stirrer bottles, spun at 25 rpm and incubated at 37°C in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FBS (heat-inactivated foetal bovine serum), 25 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid), glutamine (2mM), penicillin (100 U/ml) and streptomycin (100 μ g/ml). GM7373 cells (bovine aortic endothelial cell line) were maintained in adhesion flask with MEM (Eagle's Minimum Essential Medium) with Earle's salts, supplemented with 10% FBS, 25 mM HEPES, glutamine (2mM), penicillin (100 U/ml), streptomycin (100 μ g/ml) and enriched with 2% MEM non essential aminoacids and 2% MEM vitamin solution. IGR-1 cells (human melanoma cell lines) were grown in adhesion on

Petri dishes with MEM supplemented with 10% FBS, 25 mM HEPES, penicillin (100 U/ml) and streptomycin (100 µg/ml). P388 cells (murine leukemia cell lines) were grown in adhesion on Petri dishes with L-15 (Leibovitz) medium supplemented with 10% FBS, 25 mM HEPES, penicillin (100 U/ml) and streptomycin (100 $\mu g/ml$).

Cytotoxicity assay. J774, GM7373, IGR-1, P388, (1x10⁴ cells) were plated on 96-well microtiter plates and allowed to adhere at 37°C in 5% CO₂/95% air for 2h. Thereafter the medium was replaced with 50µl of fresh medium and 75 µl aliquot of 1:2 v/v serial dilution of each test compound (dissolved in culture medium plus 10% dimethoxyethane) was added and then the cells incubated for 24, 48, 72 h. The cells viability was assessed through an [3-(4,5-dimethyltiazol-2yl)-2,5-phenyl-2H-tetrazolium bromide] (MTT) conversion assay.²² Briefly, 25 µg of MTT (5 mg/ml) was added and the cells were incubated for additional 3 h. Following this time the cells were lysed and the dark blue crystals solubilized with 100 μ l of a solution containing 50% (v:v) N,Ndimethylformamide, 20% (w:v) SDS (sodium dodecyl sulfate) with an adjusted pH of 4.5.²³ The optical density (OD) of each well was measured with a microplate spectrophotometer equipped with a 620 nm filter. The viability of each cell line in response to treatment with compounds 11-14 and doxorubicin was calculated as:

% dead cell = $100 - (OD \text{ treated/OD control}) \times 100; (OD: optical density).$

REFERENCES

- 1. Tokuda, M.; Fujiwara, K.; Gomibuchi, T.; Hirama, M.; Uesugi, M.; Sugiura, H. Tetrahedron Lett. 1993. 34. 669.
- 2. O'Challagan, C. H.; Sykes, R. B.; Stanforth, S. E. Antimicrob. Agents Chemoter . 1976, 10, 254. Greenwood, D.; O'Grady, F. ibid 1976, 10, 249. Corraz, A. J.; Dax, S. L.; Dunlap, N. K.; Georgiopapadoukou, N. H.; Keith, D. D.; Pruess, D. L.; Rossman, P. L.; Then, R.; Unowsky, J.; Wei, C.-C. J. Med. Chem. 1992, 35, 1828.
- 3.
- 4.
- Lu, Y.-F.; Harwig, C. W.; Fallis, A. G. J. Org. Chem. **1993**, 58, 4202. Banfi, L.; Guanti G. Angew. Chem., Int. Ed. Engl. **1995**, 34, 2393. Depew, K. M.; Zeman, S. M.; Boyer, S. H.; Denhart, D. J.; Ikemoto, N.; Danishefski, S. J.; Crothers, 5. D. M. Angew Chem., Int. Ed. Engl. 1996, 35, 2797.
- 6. Wada, A.; Tode, C.; Hiraishi, S.; Tanaka, Y, Ohfusa, T.; Ito, M. Synthesis 1995, 1107.
- 7. Domling, A.; Starnecker, M.; Ugi, I. Angew Chem., Int. Ed. Engl. 1995, 34, 2238.
- Wang, J.; De Clercq, P. J. Angew Chem., Int. Ed. Engl. 1995, 34, 1749. 8.
- Sadownik, A.; Deng, G.; Janout, V.; Regen, S. L.; Bernard, E. M.; Kikuchi, K.; Armstrong, D; J. Am. 9. Chem. Soc. 1995, 117, 6138.
- 10. Masters, J. J.; Jung, D. K.; Danishefsky, S. J.; Snyder, L. B.; Park, T. K.; Isaacs, R. C. A.; Alaimo, C. A.; Young, W. B. Angew Chem., Int. Ed. Engl. 1995, 34, 452. Kolb, V. M.; Hua, D. H.; Duax, W. L. J. Org. Chem. 1987, 52, 3003.
- 11.
- 12. Priebe, W. Ed. Anthacycline Antibiotics, ACS symposium series 574, Am. Chem. Soc., Washington, DC, 1995 and references cited therein.
- Faulds, D.; Balfour, J. A.; Chrisp, P.; Langtry, H. D. Drugs 1991, 41, 400.
 Golebiewski, W. M.; Keyes, R. F.; Cushman, M. Bioorg. Med. Chem. 1996, 4, 1637.
- 15. Akhrem, A. A.; Titov, A. Y Total Synthesis of Steroid; Israel Program for Scientific Translations, Jerusalem, 1969.
- De Riccardis, F.; Izzo, I.; Tedesco. C.; Sodano, G. Tetrahedron Lett. 1997, 38, 2155.
 De Riccardis, F.; Spinella, A.; Izzo, I.; Giordano A.; Sodano, G. Tetrahedron Lett. 1995, 36, 4303.
- (a) Toh, H. T.; Okamura, W. H. J. Org. Chem. 1983, 48, 1414. (b) Yong, W.; Vandewalle, M. Syn. 18. Lett. 1996, 911.
- 19. (a) McMurry, J. E.; Scott, W. J. Tetrahedron Lett. 1983, 24, 979. (b) Mascareñas, J. L.; Sarandeses, L. A.; Castedo, L.; Mouriño, A. Tetrahedron Lett. 1991, 47, 3485.
- 20. Scott, W. J., Stille, J. K. J. Am. Chem. Soc. 1986, 108, 3033.
- Larsen, D. S.; O'Shea, M. D.; Brooker, S. Chem. Commun. 1996, 203. Mosman, T. J. Immunol. Methods 1983, 65, 55. 21
- 22.
- 23. Opipari, A. W.; Hu, H. M.; Yabkowitz, R.; Dixit, V. M. J. Biol. Chem. 1992, 267, 12424.

(Received in UK 23 April 1997; accepted 12 June 1997)