

# Selective Formation of Heterodimeric Bis-Bidentate Amino-thiol-Oxometal Complexes of Rhenium(V)

Dae Yoon Chi and John A. Katzenellenbogen\*

Department of Chemistry, University of Illinois  
1209 West California Street, Urbana, Illinois 61801

Received March 31, 1993

Complexes of  $\gamma$ -emitting isotopes of technetium and rhenium have been studied extensively in the development of agents for diagnostic imaging and radiotherapy.<sup>1</sup> In many cases, these complexes involve a chelate molecule that donates four heteroatoms (most often two nitrogen and two sulfur atoms) to an oxometal(V) core. Useful complexes may be formed from bis-bidentate (NS) and tetradentate (N<sub>2</sub>S<sub>2</sub>) ligands. While several bis-bidentate complexes have been prepared, for example, with S<sub>2</sub> (ethane-1,2-dithiol,<sup>2</sup> thiomercaptoacetic acid<sup>3</sup>), OS (2-mercaptoethanol<sup>4</sup>), and NS (2-mercaptoethylamine<sup>5</sup>) ligands, a number of factors have limited the further development of these systems: (1) introduction of substituents onto the carbon atom backbone of the bis-bidentate ligand creates stereogenic centers, with the number of possible stereoisomeric complexes being higher in the bis-bidentate than in the tetradentate case; (2) because the bis-bidentate complex is symmetrical, one cannot easily modify just one side of the complex in a controlled fashion; and (3) the bis-bidentate complex may be more labile to ligand exchange<sup>6</sup> and may therefore be less stable than the tetradentate complex.<sup>7</sup> Because of these difficulties, bis-bidentate technetium and rhenium oxo complexes have not been studied extensively.

We have recently described the preparation of two tetradentate rhenium and technetium complexes conjugated to a synthetic progestin.<sup>8</sup> While they retained nanomolar binding affinity for the progesterone receptor, the overall size of these conjugates is about twice that of normal ligands for steroid receptors, a fact that appears to impair their receptor-selective distribution in vivo; this compromises the potential of these radiolabeled complexes for diagnostic (Tc-99m) and therapeutic (Re-186 and -188) applications. Therefore, we have begun to investigate the preparation of heterodimeric bis-bidentate amino-thiol complexes of technetium and rhenium that might in themselves more closely mimic the size and shape of certain steroidal and nonsteroidal ligands for steroid receptors, in particular for the estrogen receptor.

\* Address correspondence to: John A. Katzenellenbogen, Department of Chemistry, University of Illinois, 461 Roger Adams Laboratory, Box 37, 600 S. Mathews Ave., Urbana, IL 61801. Telephone: (217) 333-6310. FAX: (217) 333-7325. BITNET: KATZENELL@UIUCSCS. INTERNET: KATZENELL@B.SCS.UIUC.EDU.

(1) (a) Steigman, J.; Eckelman, W. *The Chemistry of Technetium in Medicine*; National Academy Press: Washington, DC, 1992. (b) Clarke, M. J.; Podbielski, L. *Coord. Chem. Rev.* **1987**, *78*, 253–331.

(2) (a) Davison, A.; Orvig, C.; Trop, H. S.; Sohn, M.; DePamphilis, B. V.; Jones, A. G. *Inorg. Chem.* **1980**, *19*, 1988. (b) Bandoli, G.; Nicolini, M.; Mazzi, U.; Spies, H.; Munze, R. *Transition Met. Chem.* **1984**, *9*, 127.

(3) DePamphilis, B. V.; Jones, A. G.; Davis, M. A. *J. Am. Chem. Soc.* **1978**, *100*, 5570.

(4) Jones, A. G.; DePamphilis, B. V.; Davison, A. *Inorg. Chem.* **1981**, *20*, 1617.


(5) (a) Burns, H. D.; Dannals, R. F.; Marzilli, L. G.; Wagner, H. N., Jr. *J. Nucl. Med.* **1979**, *20*, 641. (b) Bandoli, G.; Gerber, T. I. A. *Inorg. Chim. Acta* **1987**, *126*, 205.

(6) Dannals, R. F.; Burns, H. D.; Marzilli, L. G.; Dannals, T. E.; Kramer, A. V.; Wagner, H. N., Jr. In *Application of Nuclear and Radiochemistry*; Lambrecht, R. M.; Morcos, N., Eds.; Pergamon Press: New York, 1982; pp 127–138.

(7) While most N<sub>2</sub>S<sub>2</sub> technetium(V) oxo complexes used for in vivo imaging are tetradentate, complexes of technetium in other valence states may be bis-tridentate (e.g., Tc(III)–HIDA (Loberg, M. D.; Fields, A. T. *Int. J. Appl. Radiat. Isotopes* **1978**, *29*, 167)) or even hexakis-monodentate (e.g., Tc(I)–hexaisonitrile (Abrams, M. J.; Davison, A. J.; Jones, A. G.; Costello, C. E.; Pang, H. *Inorg. Chem.* **1983**, *22*, 2798).

(8) (a) DiZio, J. P.; Fiaschi, R.; Davison, A.; Jones, A. G.; Katzenellenbogen, J. A. *Bioconjugate Chem.* **1991**, *2*, 353. (b) DiZio, J. P.; Anderson, C. J.; Davison, A.; Ehrhardt, G. J.; Carlson, K. E.; Welch, M. J.; Katzenellenbogen, J. A. *J. Nucl. Med.* **1992**, *33*, 558.

## Scheme I




Here we report the synthesis of some novel homodimeric oxorhenium(V) complexes of aminothiols, as well as the first selective synthesis and characterization of two heterodimeric complexes. The preparation of these complexes, particularly the heterodimeric ones, raises interesting issues of competitive complexation kinetics of different aminothiols.

The general method for the preparation of rhenium complexes in solution is by substitution reaction of oxotrichlorobis(triphenylphosphine)rhenium(V) (1, Cl<sub>3</sub>ReO(PPh<sub>3</sub>)<sub>2</sub>) in the presence of coordinating ligand. 1-(4'-Methoxyphenyl)amino-2-methylpropane-2-thiol (2) was prepared according to procedures similar to those used by D'Amico and Corbin.<sup>9</sup> The ligands 2 (42.2 mg, 0.20 mmol) and pyridine-2-methanethiol<sup>10</sup> (3, 25.0 mg, 0.20 mmol) are added to 1 (166.6 mg, 0.20 mmol) in 2 mL of 1 M NaOAc in methanol (Scheme I). The mixture is heated at 75 °C for 20 min and cooled to ambient temperature, resulting in the formation of a brownish purple precipitate. Recrystallization of collected precipitate from MeOH/CHCl<sub>3</sub> provides crystalline (N-(2-mercapto-2-methylpropyl)-N-(4'-methoxyphenyl)aminato)-(2-mercaptomethylpyridinato)oxorhenium(V) (4)<sup>11</sup> in essentially quantitative yield.

The results shown in Table I demonstrate the selective formation of heterodimer 4 at various concentrations and ratios of the two aminothiols 2 and 3. As the concentration of ligand 3 is increased, the yield of heterodimer 4 is reduced to some degree (entries 3–1), but even at a 1:4 ratio of 2:3, the heterodimer 4 forms in 50% yield (entry 1). Despite this preference for heterodimer formation, the reaction of 1 (166.6 mg, 0.20 mmol) with only ligand 2 (84.4 mg, 0.40 mmol) under the same conditions (entry 7) does provide a quantitative yield of the homodimer complex bis(N-(2-mercapto-2-methylpropyl)-N-(4'-methoxyphenyl)aminato)oxorhenium(V) (5).<sup>11</sup> In the absence of ligand 2, 2 equiv of ligand 3 will react with 1, providing a powdery tan product which is very polar but has not yet been characterized;<sup>12</sup> this material is not detected in our HPLC analysis.

The experiments presented in Table I demonstrate that heterodimeric complex 4 is formed in strong preference to the homodimeric complex 5, as the homodimeric complex 5 begins to form only when the ratio of 2:3 becomes very large (entries

(9) (a) D'Amico, J. J.; Dahl, W. E. *J. Org. Chem.* **1975**, *40*, 1224. (b) Corbin, J. L.; Work, D. E. *J. Org. Chem.* **1976**, *41*, 489. (c) The synthesis of amino-thiol ligand 2 is as follows:



(10) Pyridine-2-methanethiol (10% in EtOH) was purchased from Penta Manufacturing Company (Fairfield, NJ).

(11) <sup>1</sup>H NMR, <sup>13</sup>C NMR, low- and high-resolution mass spectra, and some 2D NMR spectroscopic data are available as supplementary material.


**Table I.** Product Distributions with Various Concentration of Ligands for the Reaction Shown in Scheme 1<sup>a</sup>

entry	2 ( $\mu$ mol)	3 ( $\mu$ mol)	ratio 2:3 (relative to 1)	yield of 4 (%) <sup>b</sup>	yield of 5 (%) <sup>b</sup>
1	20	80	1:4	50	0
2	20	40	1:2	85	0
3	20	20	1:1	99	0
4	40	20	2:1	97	2
5	80	20	4:1	86	4
6	80	80	4:4	99	0
7	40		2:0	0	99

<sup>a</sup> Reaction condition: the ligands 2 and 3 are added to 1 (16.6 mg, 20  $\mu$ mol) and NaOAc (200  $\mu$ mol) in 1 mL of methanol. The mixture is heated at 75 °C for 20 min. <sup>b</sup> Product total yields were determined by isolation, and product ratios by normal-phase HPLC (cyanopropyl silica gel column) on a completely soluble sample of crude product. Under these conditions, the proposed homodimeric complex from 3<sup>12</sup> is too polar to elute. Yields are based on the quantity of 1, and product ratios are based on response facts at 400 nm, calibrated against standard samples of 4 and 5.

4 and 5). The preferential formation of the heterodimeric complex 4 under these relatively mild conditions reflects a strong kinetic preference.


We have synthesized another heterodimer (*S*)-(2-mercaptomethylpyridinato)-(2-mercaptomethylpyrrolidinato)oxorhenium-(V) (7)<sup>11</sup> from ligand 3 and (*S*)-2-mercaptomethylpyrrolidine hydroiodide (6).<sup>13</sup> Because ligand 6 was prepared as the hydroiodide salt, 1 equiv of Et<sub>3</sub>N is added as an acid scavenger. Complex 7 is formed in 63% yield in the presence of Et<sub>3</sub>N and in 41% yield without any base.




In principle, NS-bis-bidentate complexes can exist as *cis* or *trans* isomers. In the case of other homodimeric NS-bis-bidentate complexes reported in the literature,<sup>5,14</sup> only the *trans* isomer has been reported; in the case of one OS-bis-bidentate complex, the *cis* isomer could be isolated, although the *trans* isomer was also detected by NMR in the unfractionated reaction product.<sup>4</sup>

With our heterodimeric bis-bidentate complexes 4 and 7, isolation of the major product, which appears as one spot on silica gel and basic alumina TLC and one peak by HPLC on cyanopropyl silica gel, gives a material that shows only one set of NMR signals. On the other hand, <sup>1</sup>H NMR of homodimeric bis-bidentate complex 5 shows two sets of signals in the ratio of 85:15, which may indicate the presence of a small amount of the *cis* isomer. The results of an X-ray structure determination of a single crystal


(12) We expected to isolate the homodimeric complex from 1 and 3 alone as the salt shown below. A molecular ion peak was not detected by FI or FAB mass spectroscopy, although the <sup>1</sup>H NMR spectrum of this unidentified product shows two doublets due to CH<sub>2</sub>S at  $\delta$  4.14 ( $J$  = 16.8 Hz) and 4.81 ( $J$  = 16.8 Hz), consistent with the structure shown.



(13) (*S*)-2-Mercaptomethylpyrrolidine was prepared from (*S*)-prolinol according to the method of Searles, Jr., et al.: Searles, S., Jr.; Roelofs, G. E.; Tamres, M.; McDonald, R. N. *J. Org. Chem.* 1965, 30, 3443. We have modified the reaction scheme shown below to provide higher yield.



(14) Baldas, J.; Bonnyman, J.; Williams, G. A. *Inorg. Chem.* 1986, 25, 150.

**Figure 1.** Molecular structure of 4.

of complex 4 (Figure 1),<sup>15</sup> obtained from MeOH/CHCl<sub>3</sub>, shows the *trans* configuration; the other complexes 5 and 7 are assumed to be *trans* by analogy. This is consistent with the general preference for the dative bond (i.e., the Re-NHR<sub>2</sub> bond in 5 and the Re-N(pyridine) bond in 4) to be *trans* to the most electron-withdrawing donor (N as opposed to S), the "trans effect".<sup>16</sup>

Bis-bidentate complexes generally have lower kinetic and thermodynamic stability than tetradentate complexes.<sup>6,7</sup> The heterodimeric rhenium complexes 4 and 7 are, however, remarkably stable under ambient conditions, and, in a preliminary study, a technetium-99m complex analogous to 4 appears to have reasonable stability in vivo as well (Chi, D. Y.; O'Neil, J. P.; Anderson, C. J.; Welch, M. J.; Katzenellenbogen, J. A., unpublished results).

It is hoped that this work, which demonstrates that NS-bis-bidentate heterodimeric complexes of oxorhenium can be prepared in a selective and controlled fashion, will stimulate a more thorough investigation of structurally novel square pyramidal technetium and rhenium oxo complexes, with the ultimate aim of designing complexes whose size, shape, and disposition of functional groups provide good mimics of ligands for binding sites in important biomolecules. We are currently investigating the stability and distribution properties of these homo- and heterodimeric technetium and rhenium complexes in vivo.

**Acknowledgment.** This work was supported by a grant from the National Institutes of Health (PHS 5R01 CA25836). We are grateful to Dr. Scott R. Wilson and Ms. Teresa Prussak-Wieckowska for X-ray diffraction studies and Dr. James P. O'Neil for COSY and HETCOR NMR experiments to assign peaks of compound 7. Dr. Alan Davison provided helpful advice.

**Supplementary Material Available:** Description of the spectroscopic characteristics of compounds 4, 5, and 7 (selected <sup>1</sup>H and <sup>13</sup>C NMR, 1D and 2D spectra) and crystal data and selected structural parameters for 4 (12 pages). Ordering information is given on any current masthead page.

(15) Crystal data and selected structural parameters for 4 are available as supplementary material.

(16) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 5 ed.; John Wiley & Sons: New York, 1988; pp 1299-1300.