


# Access to Stable Quaternary Phosphiranium Salts by P-Alkylation and P-Arylation of Phosphiranes

Julien Gasnot<sup>a</sup>  
 Clément Botella<sup>a</sup>  
 Sébastien Comesse<sup>a</sup>  
 Sami Lakhdar<sup>b</sup>  
 Carole Alayrac<sup>b</sup>  
 Annie-Claude Gaumont<sup>b</sup>  
 Vincent Dalla<sup>\*a</sup>  
 Catherine Taillier<sup>\*a</sup>

<sup>a</sup> Normandie Univ., URCOM, UNIHAVRE, FR 3032, EA 3221,  
 25 Rue Philippe Lebon, BP 540, 76058 Le Havre, France  
 vincent.dalla@univ-lehavre.fr

<sup>b</sup> Normandie Univ., LCMT, ENSICAEN, UNICAEN, CNRS,  
 6 Boulevard Marechal Juin, 14000 Caen, France



Received: 22.12.2019

Accepted after revision: 02.03.2020

Published online: 12.03.2020

DOI: 10.1055/s-0040-1708000; Art ID: st-2020-b0683-1


**Abstract** We report the preparation of phosphiranium salts by quaternization of phosphiranes, a class of sensitive, highly strained, and poorly nucleophilic cyclic phosphines. High-yielding introduction of a varied set of alkyl groups including methylene ester arms was accomplished under mild conditions. A Cu-catalyzed electrophilic arylation of phosphiranes using diaryl iodonium reagents was also achieved to yield unprecedented *P,P*-diaryl phosphiranium salts with good efficiency.

**Key words** alkylation, arylation, phosphirane, phosphiranium salt, alkylating agent, diaryliodonium salt

Nowadays, quaternary phosphonium salts (QPS) represent an important class of organophosphorus compounds which have found a broad spectrum of applications ranging from catalysis, organic synthesis to biological and medical applications.<sup>1</sup> It is therefore not surprising that the access to a large panel of QPS has been widely documented. Until recently, most of the QPS syntheses exploited the nucleophilic properties of phosphorus nonbonded lone pair of electrons to form new bonds with a variety of electrophilic species.<sup>2</sup> In particular, the direct introduction of alkyl groups on tertiary phosphines using alkyl halides (Cl, Br, I) or related sulfonates (Ms, Ts, Tf) as electrophilic partners has been extensively exploited over the years.

Compared to regular phosphines, the reactivity of phosphiranes, which are rather uncommon subtypes of cyclic phosphines, was only scarcely studied (Scheme 1). These three-membered phosphorus heterocyclic systems have been viewed for long as elusive species being largely neglected by the synthetic community. One main reason is

that phosphiranes were generally appreciated as highly strained species, moderately stable thermodynamically and, in many instances, subject to ring fragmentations or rearrangements.<sup>3</sup>



Moreover, theoretical calculations also revealed that strain of the small P-heterocycles would cause partial delocalization of the phosphorus lone pair over the cycle, resulting in a decrease of nucleophilicity in comparison to tertiary phosphines.<sup>4</sup> In light of these structural characteristics and the resultant reactivity uncertainties, the chemistry of phosphiranes, and in particular quaternization has been barely approached. To our knowledge, only a handful of such quaternizations has been reported, all of which being performed using the highly electrophilic alkylating agent methyl trifluoromethanesulfonate.<sup>5</sup> Hence, the assessment that phosphiranes can undergo quaternization with reliability and wide scope is still awaited. We have recently endeavored to tackle this problem and now report our preliminary results showcasing that phosphiranes are a


promising platform for easy synthesis of quaternary phosphiranium salts (QPrS) by alkylative and arylative functionalization. In particular, an assortment of new derivatives, including *P*-alkyl, *P*-methylene ester, and *P*-aryl units, with appealing and untapped potential reactivity in organic synthesis has been prepared.

Since we observed at the early stage of this study that introduction of large substituents on phosphorus significantly mitigated instability issues, we first selected as a model substrate phosphirane **1a** incorporating a mesityl (Mes = 2,4,6-trimethylphenyl) substituent on P, for imparting the ideal balance of thermal stability/resistance to air oxidation/reactivity. Phosphirane **1a** was prepared from its parent primary phosphine MesPH<sub>2</sub> using a straightforward one-step reaction procedure.<sup>6</sup>

A preliminary study first focused on the screening of classical methylating agents and reaction conditions for *P*-methylation of **1a**.<sup>7</sup> Of the set of reagents tested, only the highly electrophilic methyl trifluoromethanesulfonate (MeOTf) appeared sufficiently reactive to enable alkylation and form the desired phosphiranium salt **2a** in almost quantitative isolated yield (conditions A, Scheme 2).<sup>8</sup> This stands in full agreement both with literature precedents and the expected weak nucleophilicity of **1a** resulting from the sterics and the ring strain/electronics relationship discussed above.

Like any strained three-membered ring systems, phosphiranium triflate **2a** is characterized by extreme high field <sup>1</sup>H, <sup>13</sup>C, and <sup>31</sup>P chemical shifts as well as large <sup>1</sup>J<sub>C-P</sub> coupling constants.<sup>9,10</sup> It should also be noticed that, in the absence of air and water, the phosphiranium **2a** appeared completely stable over several months when stored at -20 °C. Modulation of aryl substituents on phosphorus was next evaluated. Preparation of the corresponding *P*-methyl-*P*-supermesityl phosphiranium **2b** and its *P*-phenyl analogue **2c** was again performed using MeOTf (1 equiv) at room temperature in dichloromethane. Both reactions were rapid and clean, requiring only 2 h of stirring at room temperature to reach completion and give the desired compounds in high yields (conditions A, 95% and 80%, respectively, from phosphiranes **1b** and **1c**). The high-yielding synthesis of phosphiranium **1c** adorned with a simple phenyl deserves particular mention, given the relatively modest shielding embedding this substituent.

The quaternarization of C-substituted phosphiranes was next considered. Phosphiranes **1d** and **1e**, respectively, incorporating one or two methyl groups on the ring were then reacted with methyl trifluoromethanesulfonate (conditions A). Fast reactions were again observed leading to full conversion of both phosphiranes **1d** and **1e** within 1 h to form the corresponding salts **2d** and **2e**.<sup>11</sup> On one hand, the monosubstituted phosphiranium **2d** could be isolated in 98% yield as a mixture of diastereoisomers (dr



**Scheme 2** Preparation of *P*-alkyl phosphiranium salts **2** and **3**. Conditions A: MeOTf (1–2 equiv), CH<sub>2</sub>Cl<sub>2</sub>. Conditions B: Me<sub>2</sub>O·BF<sub>4</sub> (1 equiv), CH<sub>2</sub>Cl<sub>2</sub>. Conditions C: ROTf<sup>a</sup> (1–1.2 equiv), CH<sub>2</sub>Cl<sub>2</sub>. <sup>a</sup> ROTf formed in situ by reaction of Tf<sub>2</sub>O (1.2 equiv) and the corresponding alcohols (allylOH or BnOH, 1.2 equiv) in the presence of DIPEA (1.2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> at -30 °C (see Supporting Information for full details). <sup>b</sup> Mes\* = 2,4,6-(*t*-Bu)<sub>3</sub>Ph. <sup>c</sup> Reaction performed from the diastereoisomeric mixture of *syn*- and *anti*-1-mesityl-2-methylphosphirane. For clarity, only the major *anti* diastereoisomer of **2d** was represented. <sup>d</sup> Reaction performed from the racemic *trans*-1-mesityl-2,3-dimethylphosphirane (**1e**). <sup>e</sup> Crude yield estimated (2 steps, from ROH). <sup>f</sup> Reaction run at 50 °C in DCE; n.d. = not determined.

*anti/syn* = 77:23).<sup>12</sup> As observed in the parent phosphirane series, the C-substituted phosphiranium salt **2d** appeared to be much less stable than its unsubstituted analogue **2a**, and partial degradation occurred after only a few hours at room temperature. These instability issues increased even further with additional substitution as shown by the significant proportion of degradation products formed when attempting to isolate the disubstituted phosphiranium salt **2e** from the reaction mixture.<sup>13</sup>


To verify whether the nature of the counteranion could also be modified, *P*-methylation of **1a** was next tested using the Meerwein reagent (Me)<sub>3</sub>O<sup>+</sup>BF<sub>4</sub><sup>-</sup>. The latter was found equally effective as methyl triflate to form the corresponding phosphiranium tetrafluoroborate salt **2f** in high yield within 2 h (96%, conditions B).<sup>8</sup>

Rewardingly, it was also shown that the quaternarization step could expand beyond methylation. Indeed, we were pleased to succeed in the *P*-allylation and *P*-benzylation of the phosphirane **1a** to afford the phosphiranium triflates **3a** and **3b**, which were isolated in good yields when a sequential one-pot triflation/alkylation protocol was applied from allyl and benzyl alcohols, respectively (conditions C).<sup>14,15</sup>

Formation of the *P*-ethyl phosphiranium salt **3c** was also attempted from phosphirane **1a** using the commercially available ethyl triflate.<sup>16</sup> In this case, a prolonged reaction time of 16 h and heating at 50 °C (conditions C) were found critical to reach completion and, although the desired QPrS **3c** could be formed as the major product, extensive decomposition was also observed preventing its isolation in pure form.<sup>17</sup>

To broaden diversity and open new perspectives for further functionalization, introduction of methylene ester (–CH<sub>2</sub>CO<sub>2</sub>R) groups on phosphirane **1a** was next considered. Since we expected an increased electrophilicity of the –CH<sub>2</sub> group adjacent to the proximal ester with respect to the methyl, allyl, and benzyl halides, we first tested the use of the commercially available bromo- and iodoacetate derivatives. Apart from avoiding the tedious glycolic esters derived triflate route while providing a simplified protocol, these experiments were also intended to bring valuable information refining our knowledge of phosphiranes general reactivity.

In line with earlier observations in methyl, allyl, and benzyl series, a lack of reactivity was, unfortunately, observed with only partial conversion of **1a** and degradation observed after prolonged reaction times. So, we finally had to resort to synthesizing the corresponding triflate reagents from the parent hydroxy esters. The triflation step was, somehow surprisingly, a clean process which enabled easy and high yielding (70–82%) isolation of the desired triflates **4a–d**. Their reaction with **1a** all proceeded smoothly to give the expected methylene ester phosphiranium triflates **5a–d** in high yields (80–95%) irrespective of the ester group (Scheme 3).<sup>18,19</sup>



**Scheme 3** Preparation of *P*-methylene ester phosphiranium triflates **5a–e**. Crude yields are given.

In all cases, slow reaction rates were observed, and the reactions required an extended reaction time of 24 h (vs 2 h for methylation) at room temperature to reach completion. Heating and solvent change were also tested without any improvement. As a whole, triflyloxy acetates **4a–d** turned out to be much more reactive than the corresponding halides, but clearly less than methyl triflate, thereby confirm-

ing the weak nucleophilicity of phosphiranes while demonstrating the dominant influence of sterics on reactivity. Disappointingly and for some unclear reasons, attempts to switch the ester function by a trifluoromethyl group completely suppressed the alkylation, and no trace of the corresponding quaternary phosphiranium salt **5e** could be detected when mixing **1a** with the trifluoroethyl triflate **4e**.

To further expand our phosphiranium salts portfolio, we next focused on the introduction of aryl groups on phosphirane **1a**. We postulated that hypervalent iodine derivatives should be suitable reagents for diarylphosphiranium salt synthesis through electrophilic arylation of phosphirane substrates,<sup>20,21</sup> thus opening new perspectives in phosphirane functionalization. We were delighted to be able to demonstrate that this was indeed the case.

Table 1 shows selected results of our preliminary exploration of a Cu/CuCl-catalyzed arylation quaternarization reaction of **1a** with diaryliodonium triflates **6a–c**.<sup>20d,22</sup> Reaction of **1a** with **6a** was initially conducted in 1,1,2,2-tetrachloroethane at 140 °C. Under these relatively drastic reaction conditions the diaryl phosphiranium salt **7a** was isolated in only 12% yield from the reaction mixture, in spite of complete conversion (entry 1). To limit decomposition, we next performed a series of reactions at lower temperatures and observed progressive yields enhancement following gradual decrease of the temperature (48% at 80 °C vs 81% at 50 °C, entries 2 and 3). A complete absence of reactivity was, however, observed at room temperature (entry 4). It is worth mentioning that *P,P*-diarylphosphiranium salt **7a** proved less stable than its *P*-Me,*P*-aryl analogue **2a**, with faster decomposition occurring in solution even at room temperature. The *P,P*-diarylphosphiranium salt **7a** was, however, found stable enough to be isolated by flash column chromatography on silica gel and could be stored neat for weeks at –20 °C, in the absence of air and water. Preparation of the hexafluorophosphate salt analogue **7b** was run under the same conditions using diphenyliodonium salt **6b**. In stark contrast with the triflate case, temperature influenced the reaction outcome only marginally (54% isolated yield at 80 °C vs 58% at 50 °C, entries 2 and 3 respectively), thereby suggesting a greater instability of *P,P*-diaryl QPrs paired with a hexafluorophosphate anion.

Last, introduction of the electron-rich and bulky *p*-*tert*-butylphenyl group on **1a** was attempted. Although the starting phosphirane **1a** was rapidly consumed either at 80 °C or 50 °C, the phosphiranium hexafluorophosphate salt **7c** was isolated in only 14% and 27% yields, respectively (entries 2 and 3). At this stage, it is not clear whether these low yields reflect an exacerbated instability of QPrS **7c** under the reaction conditions or are the results of an unfavorable aryl transfer.

In conclusion, we have described the preparation of unprecedented quaternary phosphiranium salts, an intriguing and barely explored class of quaternary phosphonium salts. To this end, the singular reactivity of their phosphirane pre-

**Table 1** Arylation of Phosphirane **1a** with Diaryliodonium Reagents: Optimization of Reaction Conditions

Entry	Temp (°C)	Yield <sup>a</sup> of <b>7a</b> (%)	Yield <sup>a</sup> of <b>7b</b> (%)	Yield <sup>a</sup> of <b>7c</b> (%)
1	140	12	–	–
2	80	48	54	14
3	50	81	58	27
4	20	NR	NR	NR

<sup>a</sup> Isolated yields are given; NR = no reaction.

cursors, which are highly strained and poorly nucleophilic cyclic phosphines, was studied with regard to quaternarization. Introduction of a varied set of alkyl substituents, ranging from methyl to methylene ester groups, could be accomplished in high yields when using the triflate reagents as the alkylating agents. For triflates which are not commercially available and highly sensitive, a straightforward and efficient one-pot triflation/alkylation procedure was developed. We have also demonstrated that use of hypervalent iodine reagents in a Cu-catalyzed electrophilic arylation of phosphirane substrates could offer a facile access to bisarylated compounds, thus broadening our library of phosphiranium salts. This work allowed access to known and new structures with fascinating yet untapped synthetic potential for new phosphorus chemistry. Studies on those aspects are currently ongoing in our laboratory and will be reported in due course.

## Funding Information

We gratefully acknowledge the region Haute-Normandie, the réseau CRUNCH for grants to J.G. and C.B.; J.G. also thanks the University Le Havre Normandie for financial support.

## Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0040-1708000>.

## References and Notes

- Berchel, M.; Jaffrès, P.-A. In *Organophosphorus Chemistry: From Molecules to Applications, 1st ed*; Iaroshenko, V., Ed.; Wiley-VCH: Weinheim, **2019**, 59–111; and references cited therein.
- (a) Virieux, D.; Volle, J.-N.; Pirat, J.-L. In *Science of Synthesis, Vol. 42*; Mathey, F.; Trost, B. M., Ed.; Thieme: Stuttgart, **2009**, 503–594. (b) Tebby, J. C.; Allen, D. W. In *Science of Synthesis, Vol. 31*;

Ramsden, C. A., Ed.; Thieme: Stuttgart, **2007**, 2083–2104. (c) For the first example of QPS synthesis utilizing phosphine oxides as electrophilic partners and Grignard reagents as nucleophiles, see: Vetter, A. C.; Nikitin, K.; Gilheany, D. G. *Chem. Commun.* **2018**, 54, 5843.

- (a) Mathey, F. *Chem. Rev.* **1990**, 90, 997. (b) Quin, L. D. In *A Guide to Organophosphorus Chemistry*; Quin, L. D., Ed.; John Wiley & Sons: New York, **2000**, 234. (c) Mathey, F.; Regitz, M. In *Phosphorus-Carbon Heterocyclic Chemistry: The Rise of a New Domain*; Mathey, F., Ed.; Elsevier Science: Amsterdam, **2001**, 17.
- (4) Henderson, W. A.; Buckler, S. A. *J. Am. Chem. Soc.* **1960**, 82, 5794.
- (5) (a) Hockless, D. C. R.; McDonald, M. A.; Pabel, M.; Wild, S. B. *J. Chem. Soc., Chem. Commun.* **1995**, 257. (b) Hockless, D. C. R.; McDonald, M. A.; Pabel, M.; Wild, S. B. *J. Organomet. Chem.* **1997**, 529, 189. For another recent example of isolated QPrS, see: (c) Ficks, A.; Martinez-Botella, I.; Stewart, B.; Harrington, R. W.; Clegg, W.; Higham, L. J. *Chem. Commun.* **2011**, 47, 8274.
- (6) For the synthesis of phosphirane substrates, see: (a) Mézailles, N.; Fanwick, P. E.; Kubiak, C. P. *Organometallics* **1997**, 16, 1526. (b) Yang, F.; Fanwick, P. E.; Kubiak, C. P. *Organometallics* **1999**, 18, 4222.
- (7) Partial conversions of phosphirane **1a** to the corresponding P-methyl phosphiranium salts were observed after prolonged reaction times using other methylating agents by <sup>31</sup>P NMR monitoring. Attempts to complete the methylation reactions all failed to give the products in satisfactory yields. Prolonged reaction times and heating were rather found to favor degradation of both substrate **1a** and quaternarization products. Results not shown.
- (8) **General Procedure for the Preparation of P-Me Phosphiranium Salts **2** (Conditions A or B)**  
To a solution of phosphirane **1** (1.0 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 M) was added methyl trifluoromethanesulfonate or trimethyloxonium tetrafluoroborate (1.0–2.0 equiv) dropwise at room temperature. The mixture was then stirred at room temperature and conversion of the starting material was monitored by <sup>31</sup>P NMR (approx. 2 h). The reaction mixture was then concentrated under reduced pressure to give the title compounds **2**.  
**1-Mesityl-1-methylphosphiran-1-ium Trifluoromethanesulfonate (**2a**)**  
Prepared from 1-mesitylphosphirane (**1a**, 0.84 mmol) using the general procedure with methyl trifluoromethanesulfonate (2.0 equiv). Reaction time: 2 h. The title compound **2a** was isolated as a white solid (290 mg, 99% yield). R<sub>f</sub> = 0.40 (CH<sub>2</sub>Cl<sub>2</sub>/EtOAc = 60:40); mp 58–60 °C. IR (neat): ν = 2941, 1604, 1250, 1031, 940 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, 20 °C): δ = 6.99 (d, <sup>4</sup>J<sub>H-P</sub> = 6.0 Hz, 2 H), 2.74–2.60 (m, 2 H), 2.57 (d, <sup>4</sup>J<sub>H-P</sub> = 1.6 Hz, 6 H), 2.26 (s, 3 H), 2.20 (d, <sup>2</sup>J<sub>H-P</sub> = 17.8 Hz, 3 H), 2.15–2.03 (m, 2 H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>, 20 °C): δ = 146.9 (d, <sup>4</sup>J<sub>C-P</sub> = 3.0 Hz, Cq), 144.5 (d, <sup>3</sup>J<sub>C-P</sub> = 11.0 Hz, 2 CH), 130.0 (d, <sup>2</sup>J<sub>C-P</sub> = 13.2 Hz, 2 Cq), 120.3 (q, <sup>1</sup>J<sub>C-P</sub> = 320.2 Hz, Cq), 110.7 (d, <sup>1</sup>J<sub>C-P</sub> = 93.8 Hz, Cq), 22.0 (d, <sup>3</sup>J<sub>C-P</sub> = 9.1 Hz, 2 CH<sub>3</sub>), 21.3 (d, <sup>5</sup>J<sub>C-P</sub> = 1.7 Hz, CH<sub>3</sub>), 9.3 (d, <sup>1</sup>J<sub>C-P</sub> = 4.4 Hz, 2 CH<sub>2</sub>P), 5.4 (d, <sup>1</sup>J<sub>C-P</sub> = 50.1 Hz, CH<sub>3</sub>P). <sup>31</sup>P NMR (121 MHz, CDCl<sub>3</sub>, 20 °C): δ = -116.1. <sup>19</sup>F NMR (282 MHz, CDCl<sub>3</sub>, 20 °C): δ = -78.4. HRMS (ESI): m/z calcd for C<sub>12</sub>H<sub>18</sub>P [M]<sup>+</sup>: 193.1146; found: 193.1144.
- (9) The <sup>31</sup>P NMR spectrum shows a single peak at -116.4 ppm for **2a**, while the methyl group on P appears as a doublet at 2.20 ppm (<sup>2</sup>J<sub>P-H</sub> = 17.8 Hz) in the <sup>1</sup>H NMR spectrum. In comparison, the acyclic phosphonium analogue of **2a** gives a signal at +21.2 ppm in the <sup>31</sup>P NMR spectrum, see: Sattler, A.; Parkin, G. *Chem. Commun.* **2011**, 47, 12828.

- (10) Similarly, phosphiranes give the most upfield  $^{31}\text{P}$  NMR signals known for tertiary phosphines. As a representative example, our model phosphirane **1a** gives a signal at  $-240.4$  ppm in the  $^{31}\text{P}$  NMR spectrum.
- (11) (a) The stereochemistry *anti* and *syn* refers to the relative position of the substituent on the ring carbon to the mesityl substituent at phosphorus. The reaction was performed from a diastereoisomeric mixture of *syn*- and *anti*-1-mesityl-2-methylphosphirane (**1d**). (b) The reaction was performed from the racemic *trans*-1-mesityl-2,3-dimethylphosphirane (**1e**).
- (12) (a) Stereochemistry of the products **2d** and **2e** are proposed by analogy and in accordance with Gaspar and co-workers' work who described the stereoselective synthesis of the parent phosphiranes **1d** and **1e**, see: Li, X.; Robinson, K. D.; Gaspar, P. P. *J. Org. Chem.* **1996**, *61*, 7702. (b) It is worth noting that due to instability issues, which differ from one diastereomer to another in both the phosphirane and the phosphiranium series, significant variations in the *dr* values of the isolated products may be observed.
- (13) It should be noted that due to their increased instability: (a) HRMS and IR data are missing from the characterization of phosphiranium **2d**. (b) neither determination of a reliable chemical yield nor full characterization of the quaternarization product **2e** could be done.
- (14) Isolation of both allyl and benzyl triflate could not be realized. Therefore, in situ formation of the triflate reagents and direct addition of the reaction mixture onto a dichloromethane solution of phosphirane **1a** at  $-30$  °C was realized. For a similar procedure involving formation and direct use of allyl triflate, see: Corey, E. J.; Helal, C. J. *Tetrahedron Lett.* **1996**, *37*, 5675.
- (15) **General Procedure for the Preparation of P-Alkyl Phosphiranium Triflate 3 (Conditions C)**  
To a solution of diisopropylethylamine (1.2 equiv) and the corresponding alkyl alcohol (1.2 equiv) in dry  $\text{CH}_2\text{Cl}_2$  (0.6 M) at  $-30$  °C was added dropwise trifluoromethanesulfonic anhydride (1.2 equiv) under argon atmosphere. The reaction mixture was successively stirred 10 min at  $-30$  °C, warmed to  $0$  °C, and stirred for 10 min. The reaction mixture was again cooled to  $-30$  °C. Then, dry  $\text{Et}_2\text{O}$  was added to precipitate the ammonium salts. After decantation, the supernatant solution was added dropwise to a solution of phosphirane **1** (1 equiv) in dry  $\text{CH}_2\text{Cl}_2$  (1 M) at  $-50$  °C. The reaction mixture was then allowed to warm to room temperature over a period of 4 h, and stirring was pursued at room temperature. Conversion of the starting material was monitored by  $^{31}\text{P}$  NMR spectroscopy. Evaporation of the solvent under reduced pressure then afforded the title product **3**.  
**1-Allyl-1-mesitylphosphiranium Trifluoromethanesulfonate (3a)**  
Prepared from 1-mesitylphosphirane (**1a**, 0.84 mmol) using the general procedure with allyl alcohol (1.2 equiv). Reaction time: 4 h. The title compound **3a** was isolated as a thick colorless oil (350 mg, 95% crude yield),  $R_f = 0.15$  (EtOAc). IR (neat):  $\nu = 2944, 1605, 1250, 1153, 1025, 940$   $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = 7.03$  (d,  $^4J_{\text{H-P}} = 5.7$  Hz, 2 H), 5.70 (m, 1 H), 5.55–5.39 (m, 2 H), 3.41 (dd,  $^2J_{\text{H-P}} = 18.7$  Hz and  $^3J_{\text{H-H}} = 7.6$  Hz, 2 H), 2.78–2.68 (m, 2 H), 2.58 (br s, 6 H), 2.31 (s, 3 H), 2.20–2.10 (m, 2 H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = 147.3$  (d,  $^4J_{\text{C-P}} = 3.2$  Hz, Cq), 145.2 (d,  $^3J_{\text{C-P}} = 10.8$  Hz, 2 CH), 130.2 (d,  $^2J_{\text{C-P}} = 13.3$  Hz, 2 Cq), 125.3 (d,  $^3J_{\text{C-P}} = 16.2$  Hz,  $\text{CH}_2$ ), 123.5 (d,  $^2J_{\text{C-P}} = 14.0$  Hz, CH), 120.3 (q,  $^1J_{\text{C-F}} = 319.6$  Hz,  $\text{CF}_3$ ), 109.3 (d,  $^1J_{\text{C-P}} = 86.1$  Hz, Cq), 25.8 (d,  $^1J_{\text{C-P}} = 42.6$  Hz,  $\text{CH}_2\text{P}$ ), 22.5 (d,  $^3J_{\text{C-P}} = 7.9$  Hz, 2  $\text{CH}_3$ ), 21.5 (d,  $^5J_{\text{C-P}} = 1.2$  Hz,  $\text{CH}_3$ ), 9.0 (m, 2  $\text{CH}_2\text{P}$ ).  $^{31}\text{P}$  NMR (121 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = -114.1$ .  $^{19}\text{F}$  NMR (282 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = -78.5$ . HRMS (ESI):  $m/z$  calcd for  $\text{C}_{14}\text{H}_{20}\text{P}$  [ $\text{M}$ ] $^+$ : 219.1303; found: 219.1298.
- (16) Crude yield estimated from  $^1\text{H}$  NMR analysis of the crude mixture.
- (17) Formation of *P*-ethyl phosphiranium triflate **3c** as a major product was observed by monitoring progress of the quaternarization reaction starting from phosphirane **1a** by  $^1\text{H}$  NMR and  $^{31}\text{P}$  NMR spectroscopy. The  $^{31}\text{P}$  NMR spectrum shows a single peak at  $\delta = -109.6$  ppm for **3c**, while the ethyl group on P appears as a dt at  $\delta = 1.42$  ppm (3 H,  $^3J_{\text{P-H}} = 27.0$  Hz and  $^3J_{\text{H-H}} = 7.6$  Hz) and a multiplet at  $\delta = 2.50$ – $2.25$  ppm (2 H) in the  $^1\text{H}$  NMR spectrum.
- (18) **General procedure for the preparation of P-methylene ester phosphiranium triflates 5:**  
a) **Triflate Reagents Formation:** To a solution of the corresponding glycolic ester (1 equiv) and pyridine (1.3 equiv) in  $\text{CH}_2\text{Cl}_2$  (0.25 M) at  $-10$  °C, was added triflic anhydride (1.15 equiv) dropwise, under argon atmosphere. Reaction was stirred 1 h at  $-10$  °C and 3.5 h at room temperature. Degassed water (0.2 M) was next added to the reaction mixture. After decantation, the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 5$  mL/mmol). The combined organic layers were then combined, dried over  $\text{MgSO}_4$ , filtered, and concentrated under vacuum to give the corresponding triflate reagents **4**.  
b) **Alkylation of Phosphirane 1:** To a solution of phosphirane (1 equiv) in MeCN (0.2 M) was added the corresponding triflate reagent **4** (1 equiv). The reaction mixture was then stirred at room temperature under argon atmosphere until complete conversion of phosphirane **1** ( $^{31}\text{P}$  NMR monitoring, approx. 24 h). The solvent was then removed under vacuum to give the crude methylene ester phosphiranium salts **5**.  
**1-Mesityl-1-(2-ethoxy-2-oxoethyl)phosphiranium Trifluoromethanesulfonate (5b)**  
Prepared from 1-mesitylphosphirane (**1a**, 0.40 mmol) using the general procedure with the triflate reagent derived from ethyl glycolate (1.0 equiv). Reaction time: 24 h. The title compound **5b** was isolated as a thick colorless oil (394 mg, 95% crude yield).  $R_f = 0.23$  ( $\text{CH}_2\text{Cl}_2/\text{acetone} = 50:50$ ). IR (neat):  $\nu = 2985, 1724, 1606, 1243, 1223, 1156, 1027$   $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = 7.09$  (d,  $^4J_{\text{P-H}} = 6.2$  Hz, 2 H), 4.19 (q,  $^3J_{\text{H-H}} = 7.2$  Hz, 2 H), 4.03 (d,  $^2J_{\text{P-H}} = 16.8$  Hz, 2 H), 3.17–3.07 (m, 2 H), 2.71 (s, 6 H), 2.37 (s, 3 H), 2.28–2.18 (m, 2 H), 1.24 (t,  $^3J_{\text{H-H}} = 7.2$  Hz, 3 H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = 164.6$  (d,  $^2J_{\text{P-C}} = 5.1$  Hz, C=O), 147.7 (d,  $^4J_{\text{P-C}} = 3.2$  Hz, Cq), 145.7 (d,  $^3J_{\text{P-C}} = 11.4$  Hz, 2 CH), 130.5 (d,  $^2J_{\text{P-C}} = 14.0$  Hz, 2 Cq), 120.4 (q,  $^1J_{\text{F-C}} = 319.4$  Hz,  $\text{CF}_3$ ), 109.1 (d,  $^1J_{\text{P-C}} = 93.0$  Hz, Cq), 63.5 ( $\text{CH}_2$ ), 30.0 (d,  $^1J_{\text{P-C}} = 56.2$  Hz,  $\text{CH}_2\text{P}$ ), 22.4 (d,  $^3J_{\text{P-C}} = 8.5$  Hz, 2  $\text{CH}_3$ ), 21.7 (d,  $^5J_{\text{P-C}} = 1.6$  Hz,  $\text{CH}_3$ ), 13.8 ( $\text{CH}_3$ ), 10.3 (d,  $^1J_{\text{P-C}} = 2.7$  Hz, 2  $\text{CH}_2\text{P}$ ).  $^{31}\text{P}$  NMR (121 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = -118.6$ .  $^{19}\text{F}$  NMR (282 MHz,  $\text{CDCl}_3$ ,  $20$  °C):  $\delta = -78.5$ . HRMS (ESI):  $m/z$  calcd for  $\text{C}_{15}\text{H}_{22}\text{O}_2\text{P}$  [ $\text{M}$ ] $^+$ : 265.1357; found: 265.1360.
- (19) Crude yields are given since purification attempts only led to the decomposition of the products.
- (20) For examples of metal-catalyzed arylation of tertiary phosphines, see: (a) Ziegler, C. B. Jr.; Heck, R. F. *J. Org. Chem.* **1978**, *43*, 2941. (b) Marcoux, D.; Charette, A. B. *J. Org. Chem.* **2008**, *73*, 590. (c) Marcoux, D.; Charette, A. B. *Adv. Synth. Catal.* **2008**, *350*, 2967. (d) Hanamoto T., Kiguchi Y., Shindo K., Matsuoka M., Kondo M.; *Chem. Commun.*; **1999**, 151. For a photoredox reaction mediated by a Ru-based photosensitizer, see: (e) Fearnley, A. F.; An, J.; Lindovska, P.; Denton, R. M. *Chem. Commun.* **2016**, *52*, 4987.

- (21) For examples of metal-free arylation methods, see: (a) Wittig, G.; Matzura, H. *Angew. Chem., Int. Ed. Engl.* **1964**, *3*, 231. (b) Rémond, E.; Tessier, A.; Leroux, F. R.; Bayardon, J.; Jugé, S. *Org. Lett.* **2010**, *12*, 1568. (c) Huang, W.; Zhong, C.-H. *ACS Omega* **2019**, *4*, 6690. (d) Bugaenko, D. I.; Volkov, A. A.; Livantsov, M. V.; Yurovskaya, M. A.; Karchava, A. V. *Chem. Eur. J.* **2019**, *25*, 12502.
- (22) **General Procedure for the Preparation of Diaryl Phosphiranium Salts 6**

To a solution of phosphirane **1** (1 equiv) in 1,1,2,2-tetrachloroethane (0.1 M) were added diphenyliodonium salt (1.2 equiv), copper chloride (10 mol%), and copper metal wire (cat.) at room temperature. The reaction mixture was then heated at 50 °C. After cooling down to room temperature, the resulting crude mixture was directly purified by flash column chromatography on silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub> then CH<sub>2</sub>Cl<sub>2</sub>/acetone, 60:40) to yield the title product **7**.

**1-Mesityl-1-phenylphosphiranium Trifluoromethanesulfonate (7a)**

Prepared from 1-mesitylphosphirane (**1a**, 0.14 mmol) using the general procedure with diphenyliodonium triflate (1.2 equiv). Reaction time: 1 h. The title compound **7a** was isolated as a colorless oil (42 mg, 81% crude yield). *R<sub>f</sub>* = 0.40 (CH<sub>2</sub>Cl<sub>2</sub>/acetone = 60:40). IR (neat): 1603, 1257, 1169, 1034, 636 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, acetone-*d*<sub>6</sub>, 20 °C): δ = 7.93–7.82 (m, 2 H), 7.81–7.68 (m, 3 H), 7.35 (d, <sup>4</sup>*J*<sub>H-P</sub> = 5.9 Hz, 2 H), 3.36–3.23 (m, 2 H), 3.08–2.96 (m, 2 H), 2.70 (d, d, <sup>4</sup>*J*<sub>H-P</sub> = 1.7 Hz, 6 H), 2.44 (s, 3 H). <sup>13</sup>C NMR (75 MHz, acetone-*d*<sub>6</sub>, 20 °C): δ = 148.5 (d, <sup>4</sup>*J*<sub>C-P</sub> = 3.1 Hz, Cq), 147.1 (d, <sup>2</sup>*J*<sub>C-P</sub> = 10.7 Hz, 2 Cq), 135.8 (d, <sup>4</sup>*J*<sub>C-P</sub> = 4.5 Hz, CH), 133.4 (d, <sup>3</sup>*J*<sub>C-P</sub> = 13.3 Hz, 2 CH), 131.6 (d, <sup>3</sup>*J*<sub>C-P</sub> = 16.4 Hz, 2 CH), 131.1 (d, <sup>2</sup>*J*<sub>C-P</sub> = 13.4 Hz, 2 CH), 122.3 (q, <sup>1</sup>*J*<sub>C-F</sub> = 321.8 Hz, CF<sub>3</sub>), 115.2 (d, <sup>1</sup>*J*<sub>C-P</sub> = 90.5 Hz, Cq), 109.2 (d, <sup>1</sup>*J*<sub>C-P</sub> = 97.4 Hz, Cq), 22.6 (d, <sup>3</sup>*J*<sub>C-P</sub> = 8.9 Hz, 2 CH<sub>3</sub>), 21.6 (d, <sup>5</sup>*J*<sub>C-P</sub> = 1.4 Hz, CH<sub>3</sub>), 12.9 (d, <sup>1</sup>*J*<sub>C-P</sub> = 3.2 Hz, 2 CH<sub>2</sub>P). <sup>31</sup>P NMR (121 MHz, acetone-*d*<sub>6</sub>, 20 °C): δ = -113.8. <sup>19</sup>F NMR (282 MHz, acetone-*d*<sub>6</sub>, 20 °C): δ = -78.9. HRMS (ESI): *m/z* calcd for C<sub>17</sub>H<sub>20</sub>P [M]<sup>+</sup>: 255.1302; found: 255.1303.