

Figure 2. $[H_5O_2]^+$ groups assembled into a ring which is a link of a chain extending along b. The repeating unit is $[H_9O_4]^+$. The H₂O molecules which bind the $[H_5O_2]^+$ groups together are rather loosely held and display large thermal motion. Large circles are oxygen and small circles hydrogen atoms.



Figure 3. Configuration of the $[H_5O_2]^+$ ion. The two O(1) and the two $\frac{1}{2}H(1)$ sites are coplanar. The four H(2) sites are symmetry related. The group configuration is strictly *trans*. The O-O separation is 2.57 \pm 0.01 Å and the O-H···O angle is 172°.

which is a link in the protonated water molecule chain. Hydrogen-bonding interactions between adjacent chains as well as those between adjacent $(AuCl_4)^-$ and $[(H_5O_2)^+ \cdot 2H_2O]$ layers appear to be weak or absent.

The most interesting structural feature is the $(H_5O_2)^+$ group which is illustrated in Figure 3. The required symmetry of the group is 2/m and the configuration is *trans* as indicated. The bridging proton, originally expected to be centered, definitely refined to disordered, off-bond-line positions with a site separation of 0.62 Å.

This rather remarkable configuration raises interpretative problems. There is an intriguing similarity between the now well-known pyramidal oxonium ion and the two symmetry related halves of the $(H_5O_2)^+$ group described here. This suggests that $(H_5O_2)^+$ is simply $(H_3O^+ \cdot H_2O)$. The static disorder which this implies and the nonequivalence of the three $O-H \cdot \cdot \cdot O$ bonds involved tend to weaken this interpretation. We prefer to consider the grouping a variant of the diaquated proton⁸ with an unusual symmetrical double-well potential. The dynamic disorder implied in this interpretation should be verifiable by spectroscopic measurements. Such measurements are now in progress.

(8) J. M. Williams, "Proceedings of the Second Materials Research Symposium, 1967," Special Publication No. 301, National Bureau of Standards, Washington, D. C., in press.

Jack M. Williams, S. W. Peterson

Chemistry Division, Argonne National Laboratory Argonne, Illinois 60439 Received November 12, 1968 The Geometry of the Transition State of the Retrograde Homo Diels-Alder Reaction. A Stereospecific Conversion of 1,3- to 1,4-Dienes¹

Sir:

The efficiency with which a small ring transmits conjugative and orbital symmetry influences can be evaluated from a knowledge of the stereochemistry of the homo Diels-Alder reaction $(1 \rightleftharpoons 2)$. The present work describes a solution of this problem for the retrograde $(2 \rightarrow 1)$ monohomo case (n = 1).² The results demonstrate that a cyclopropane ring exerts strict control of the stereochemistry of the olefinic product, even in the face of strongly countervailing steric effects. They also provide a stereospecific synthetic method for homologation of 1,3-dienes to 1,4-dienes.



Model studies of ordinary Diels-Alder retrogression show a high degree of stereospecificity. Thus, oxidation of the cis hydrazo compound 3 by air, or preferably of an ether solution by yellow mercuric oxide or manganese dioxide in the presence of anhydrous sodium sulfate at 25°, gives a nearly quantitative yield of nitrogen, presumably via the very unstable azo compound 4. Vapor chromatography (vpc) of the solution shows trans, trans-2, 4-hexadiene (5) as the only volatile organic product under analytical conditions that would reveal 0.1% cis, trans- or cis, cis-2, 4-hexadiene. Similarly, oxidation of trans-hydrazo compound 6 gives only cis, trans-2, 4-hexadiene (8). 3-8 The ultraviolet chromophore of the presumed azo compounds 4 and 7 is not observed, even at -50° . The exceptionally high rate9 and stereospecificity suggest double bond participation in both the rate- and product-determining steps of the decomposition of 4 and 7. Although one cannot strictly exclude a two-step mechanism with an intermediate (e.g., 11) in which loss of nitrogen is much faster than bond rotation, a simple and conceptually economical alternative would picture the reaction as a concerted, orbital-symmetry-allowed¹⁰ retro-Diels-Alder process.

(1) This work was supported in part by grants from the Petroleum Research Fund and the AFOSR (SRC)-OAR, U. S. Air Force Grant No. 68-1358.

(2) During our study, W. L. Mock, 155th National Meeting of the American Chemical Society, San Francisco, Calif., April 1968, Abstract P-20, reported some stereochemical results of the analogous decomposition of 3-bicyclo[3.1.0]thiahexane 3,3-dioxides.

(3) Hydrazo compounds 3 and 6 result from hydrolytic decarboxylation of the corresponding esters, 9⁶ and 10.⁵ Compound 9 is formed in virtually quantitative yield from the reaction of 5 and dimethyl azodicarboxylate;⁶ 8 and dimethyl azodicarboxylate react more sluggishly to give a 4:1 mixture of 10 and 9 (boiling benzene, 18 hr).

(4) Diazenes⁷ and sulfolenes⁸ behave analogously.

(5) Elemental analysis confirms the composition.

(6) R. Daniels and K. A. Roseman, *Tetrahedron Letters*, 1335 (1966).
(7) D. M. Lemal and S. D. McGregor, J. Am. Chem. Soc., 88, 1335 (1966).

(8) (a) S. D. McGregor and D. M. Lemal, *ibid.*, 88, 2858 (1966);
(b) W. L. Mock, *ibid.*, 88, 2857 (1966).

(9) Saturated analogs do not decompose at appreciable rates below 140° (see, *inter alia*, P. D. Bartlett and N. A. Porter, *ibid.*, 90, 5317 (1968)].

(10) R. Hoffmann and R. B. Woodward, Accounts Chem. Res., 1, 15 (1968).



The CuCl-catalyzed action of diazomethane on the *trans* diester 10 in ether-pentane gives the *trans*-cyclopropane 12;⁵ on the *cis* diester 9, it gives a mixture of



one part of *syn-cis*-cyclopropane 13⁵ and eight parts of *anti-cis* 14.⁵ Separation of 13 and 14 is achieved by vpc on a Carbowax column.



The nmr spectra of the cyclopropanation products in the *cis* series 13 and 14 do not lead to an unambiguous assignment of stereochemistry, but the minor product is shown to be 13 by independent synthesis from the cycloheptatriene–N-phenyltriazolinedione adduct 15.¹¹ The assignment¹¹ to 15 of the indicated orientation of the cyclopropane ring is confirmed by the observation that the nmr chemical shifts of the cyclopropane methylene hydrogens in 15 lie at much higher fields (0.5 ppm for *endo* and 0.2 ppm for *exo*) than those in the dihydro derivative obtained by diimide reduction of the double bond. Successive treatment of 15 with ozone



(11) R. C. Cookson, S. S. H. Gilani, and I. D. R. Stevens, Tetrahedron Letters, 615 (1962).

Journal of the American Chemical Society | 91:3 | January 29, 1969

and NaBH₄ gives the crystalline diol⁵ 16, from which 13 is derived by the steps shown.

Procedures analogous to those used in the olefinic series convert the three bicyclic diesters 12, 13. and 14 to the corresponding hydrazo compounds 17, 18, and 19, which are oxidized as before. The azo compounds presumed to result are unstable at 25° and decompose smoothly to nitrogen and 2,5-heptadiene. Olefin formation is again highly stereospecific (Table I).

Table I. Products of Oxidation of Hydrazo Compounds

			//
	cis, trans ^a	trans, trans ^a	cis.cis ^a
17 (trans)	100	0%	0*
18 (syn-cis)	ە0	1	99
19 (anti-cis)	05	98	2

^a Dienes independently synthesized by reductions of 2,5-heptadiyne with stereospecific alkylaluminum hydride reagents [cf. G. Zweifel and R. B. Steele, J. Am. Chem. Soc., **89**, 5085 (1967)]. The nmr spectrum of the *trans,trans* diene is accurately simulated by a spectrum computed with the LAOCOON II program [A. A. Bothner-By and S. Castellano, J. Chem. Phys., **41**, 3863 (1964)], using reasonable coupling constants. ^b As little as 0.5% would be detectable.

The small amounts of cross products observed from 18 and 19 probably arise from small amounts of mutual cross-contamination of the two precursors 13 and 14, which are difficultly separable by vpc.

The results are consistent with a concerted retrograde homo Diels-Alder mechanism. A strong preference is manifest for a transition-state geometry in which the nitrogen molecule departs *anti* to the methylene group of the cyclopropane ring. Especially noteworthy is the exclusive formation of *cis,cis*-2,5-heptadiene from *syn-cis*-hydrazo compound **18**, a reaction which passes over transition state **20** rather than its conformational isomer **21**. That **20** must involve a severe methyl-



methyl repulsion is evident from the complete absence of *cis,cis*-2,4-hexadiene in the analogous oxidation of olefinic *cis*-hydrazo compound **3**. Clearly, some overriding influence is present in the bicyclic transition state. Although compensatory steric effects may make small contributions, the major factor is probably electronic. Apparently, in the bicyclic case, the requirement for orbital overlap of the three breaking bonds¹² is so stringent as to overcome the strain completely.

(12) The large enhancement in the rate of thernial decarbonylation of *endo*-tricyclo[3.2.1.0² 4]octen-8-one over its *exo* isomer is interpreted similarly [B. Halton, M. A. Battiste, R. Rehberg, C. L. Deyrup, and M. E. Brennan, *J. Am. Chem. Soc.*, **89**, 5964 (1967)].

(13) This investigation was supported in part by National Institutes of Health Postdoctoral Fellowship No. 2 FO2CA25233-02 from the National Cancer Institute.

> Jerome A. Berson, Stephen S. Olin¹³ Department of Chemistry, University of Wisconsin Madison, Wisconsin 53706 Received November 16, 1968