

ONE-POT SYNTHESIS OF DIHYDROXYCYCLOPENTANES FROM ALLYLSULFONES AND CHIRAL DIEPOXIDES



Hiroaki MIYAOKA and Masahiro KAJIWARA*

Department of Medicinal Chemistry, Meiji College of Pharmacy, 1-22-1 Yato-cho, Tanashi-shi, Tokyo 188, Japan












Allylphenylsulfonyl lithium, generated from allyl phenyl sulfone and n BuLi, reacts with chiral diepoxides to give dihydroxycyclopentane derivatives in good yields. The synthesis of the aldehyde **2**, which is a key intermediate for brefeldin A, was achieved using this reaction.

KEYWORDS one-pot synthesis; dihydroxycyclopentane derivative; allyl phenyl sulfone; chiral diepoxide; brefeldin A

Sulfone compounds are useful in carbon-carbon bond-forming reactions, particularly in cyclization.¹⁻⁴ For example, Eisch and co-workers reported the conversion of 4-bromo-1,2-epoxybutane and [(phenylsulfonyl)methylene] dilithium to 3-(phenylsulfonyl)-cyclopentanol in 84% yield.³ But simultaneous formation of two new carbon-carbon bonds has not yet been reported. We wish to report a new reaction of phenylsulfonyl compounds with chiral diepoxides for the formation of optically active cyclopentane derivatives. This reaction was applied to a synthesis of the aldehyde **2**,^{5c} which is a key synthetic intermediate of (+)-brefeldin A (**1**).⁵



The lithio derivatives of sulfone compounds were prepared from the sulfone **6** (2.5 eq) and n -butyllithium (2.4 eq) in tetrahydrofuran (THF) at -78°C for 1 h. The diepoxide (1.0 eq) in THF was then added dropwise at -78°C . The reaction mixture was stirred at -78°C for 1 h, and then kept at room temperature with stirring for 1 h. The reaction of two allyl phenyl sulfones gave the dihydroxycyclopentane derivatives as diastereomeric mixtures in the ratio of 1:1 via selective *exo* cyclization in 78% and 95% yields based on the diepoxide **3**^{7,8} (entries 1, 2). The reaction of phenyl propargyl sulfone gave the cyclopentane derivative as a diastereomeric mixture in the ratio of 6:1 in 84% yield (entry 3). The diepoxide **4**,⁹ a diastereomer of **3**, reacted with allyl phenyl sulfone to form the cyclopentane as a sole product in 71% yield (entry 4). The diepoxide **5**,⁹ which was protected with benzylether, gave the cyclopentane derivative in the ratio of 1:1 in 95% yield (entry 5). The diepoxide

Entry	Sulfone	Diepoxide	Products	Yield, ^{a)} %
1		 3		78 ^{b)}
2		3		95 ^{b)}
3		3		84 ^{c)}
4		 4		71 ^{d)}
5		 5		95 ^{b)}
6		 6	 7	95 ^{e)}

a) Isolated yield based on the diepoxide.

b) A 1:1 mixture of diastereomer was generated.

c) A 6:1 mixture of diastereomer was generated.


d) A single compound, whose stereochemistry was not determined.

e) A single compound.

6⁹⁾ reacted with allyl phenyl sulfone to form the cyclopentane 7 as a single product, whose stereochemistry was determined by NOE experiment, in 95% yield (entry 6).

Based on this approach to constructing dihydroxycyclopentane derivatives, a synthetic study for brefeldin A was carried out. The hydroxy group of 7 was protected with 2-methoxyethoxymethyl chloride to give 8. The terminal olefin of 8 was converted to an aldehyde 9. The phenylsulfonyl group of 9 was removed by treatment with SmI₂ in the presence of HMPA. The isomerization of the formyl group was carried out by treatment with K₂CO₃ in methanol to give the aldehyde 2.¹⁰⁾ The aldehyde 2 has already been synthesized by Taber's group in their synthesis of (+)-brefeldin A.

The present results clearly demonstrate the potential of this method for syntheses of not only brefeldin A but also other natural products.




- a) MEM-Cl, $i\text{-Pr}_2\text{NEt}$, $\text{CH}_2\text{ClCH}_2\text{Cl}$, 50°C , b) O_3 , $\text{CH}_2\text{Cl}_2\text{-MeOH}$, -78°C , then Me_2S , r.t.,
c) i) SmI_2 , THF-HMPA , r.t., ii) K_2CO_3 , MeOH , r.t.,

ACKNOWLEDGMENT We are grateful to Prof. D. F. Taber, University of Delaware, for $^1\text{H-NMR}$ spectrum of the aldehyde **2**. We thank Takasago Research Institute for the generous supply of (R)-methyl 4-chloro-3-hydroxybutyrate.

REFERENCES AND NOTES

- 1) a) B. Corbel, and T. Durst, *J. Org. Chem.*, **41**, 3648 (1976); b) B. Corbel, J. M. Decesare, and T. Durst, *Can. J. Chem.*, **56**, 1415 (1978); c) J. M. Decesare, B. Corbel, T. Durst, and J. F. Blount, *Can. J. Chem.*, **59**, 1415 (1981).
- 2) a) Y. Gaoni, *Tetrahedron Lett.*, 503 (1976); b) Y. Gaoni, *Tetrahedron Lett.*, 4339 (1981); c) Y. Gaoni, *J. Org. Chem.*, **47**, 2564 (1982).
- 3) J. J. Eisch, S. K. Dua, and M. Behrooz, *J. Org. Chem.*, **50**, 3674 (1985).
- 4) J. H. Babler, *J. Org. Chem.*, **52**, 4614 (1987).
- 5) Recent synthetic studies on (+)-brefeldin A: a) E. J. Corey, and P. Carpino, *Tetrahedron Lett.*, **31**, 7555 (1990); b) J. Nokami, M. Ohkura, Y. Dan-oh, and Y. Sakamoto, *Tetrahedron Lett.*, **32**, 2409 (1991); c) D. F. Taber, L. J. Silverberg, and E. D. Robinson, *J. Am. Chem. Soc.*, **113**, 6639 (1991) and references cited therein.
- 6) Phenylsulfonyl compounds were prepared by treating the corresponding halides with sodium benzenesulfinate in DMF (80-95% yield).
- 7) Diepoxides **3** was synthesized as follows from (R)-methyl 4-chloro-3-hydroxybutyrate (96.6% ee):



- a) i) TBS-Cl, ii) DIBAL-H, toluene, -78°C , iii) $\text{Ph}_3\text{P=CHCO}_2\text{Et}$, iv) DIBAL-H, CH_2Cl_2 , -78°C , b) i) TBHP, $\text{Ti}(\text{O}^i\text{Pr})_4$, L-(+)-DET, ii) MOM-Cl, iii) $n\text{-Bu}_4\text{NF}$, iv) K_2CO_3 , MeOH
- 8) Data for **3**: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm: 1.79 (1H, dt, $J=15.0$, 5.9 Hz), 2.07 (1H, dt, $J=15.0$, 4.2 Hz), 2.60 (1H, dd, $J=4.9$, 2.8 Hz), 2.99 (1H, m), 3.04 (1H, m), 3.07 (1H, m), 3.38 (3H, s), 3.57 (1H, dd, $J=11.7$, 5.5 Hz), 3.77 (1H, dd, $J=11.7$, 3.3 Hz), 4.66 (2H, s); $[\alpha]_{\text{D}}^{-13.8^\circ}$ ($c=4.37$, CHCl_3).
 - 9) Diepoxides **4**, **5** and **6** were synthesized by a similar method to that described for **3**.
 - 10) Data for **2**: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm: 1.71 (1H, ddd, $J=13.6$, 10.3, 4.9 Hz), 1.96 (1H, m), 2.0 - 2.1 (2H, m), 2.68 (1H, m), 2.78 (1H, m), 3.37 (3H, s), 3.38 (3H, s), 3.5-3.8 (10H, m), 3.80 (1H, m), 4.25 (1H, m), 4.46 (1H, d, $J=11.8$ Hz), 4.50 (1H, d, $J=11.8$ Hz), 4.67 (1H, d, $J=7.0$ Hz), 4.67 (1H, d, $J=7.0$ Hz), 4.75 (1H, d, $J=6.9$ Hz), 4.88 (1H, d, $J=6.9$ Hz), 7.25 - 7.4 (5H, m), 9.60 (1H, d, $J=2.3$ Hz); IR (neat): 1720 cm^{-1} ; $[\alpha]_{\text{D}}^{-27.3^\circ}$ ($c=0.66$, CHCl_3).

(Received April 16, 1992)