


Pd(OAc)₂-Catalyzed Macrocyclization of 1,2-Diazonaphthoquinones with Cyclic Ethers

Mitsuru Kitamura,* Masato Kisanuki, Koichi Kanemura, and Tatsuo Okauchi

Department of Applied Chemistry, Kyushu Institute of Technology, 1-1 Sensuicho, Tobata, Kitakyushu, 804-8550 Japan


S Supporting Information

ABSTRACT: Pd(OAc)₂ was found to be an efficient catalyst for the macrocyclization of 1,2-diazonaphthoquinones and cyclic ethers. This transformation serves as an efficient method for the synthesis of protected 1,2-naphthalenediols.



α -Diazocarbonyl compounds are easily decomposed to the corresponding α -carbonylcarbenes or metal carbenes via photoirradiation, electrophilic activation, or the treatment of metal complexes.¹ Thus formed, these carbenes are electrophilic and readily react with various nucleophilic Lewis bases to generate the corresponding ylides, which are widely used in organic synthesis.¹ For example, oxonium ylides are formed via the reaction of these carbenes and cyclic ethers, and then Stevens-type rearrangement,² polymerization,³ or macrocyclization^{4–6} proceeds subsequently. In the presence of a Rh-catalyst, carbenes formed from relatively stable diazo compounds, such as 2-diazo-1,3-dicarbonyl compounds **1**, react with cyclic ethers followed by preferential macrocyclization via oxonium ylide **2** to give the large cyclic ethers **3** and **4** with more than two molecules incorporated (Scheme 1, Paths a and b).^{5,6} Although there are few reports on the formation of medium-sized ring products **5** via the intramolecular cyclization of ylide **2** (Path c),^{5a} recently, Rh-catalyzed cyclization of 2-diazo-1,3-dicarbonyl

Scheme 1. Rh-Catalyzed Formation of Macro Cyclic Ethers via the Reaction of 2-Diazo-1,3-dicarbonyl Compounds **1** and Small Cyclic Ethers




compounds **1** and five- or six-membered ring cyclic acetals was achieved, which was designed by [N]-*endo-trig* cyclization ([N] = 8 or 9).^{5b}

Previously, we developed an efficient synthetic method for the preparation of 1,2-diazonaphthoquinones via diazo-transfer with 2-azido-1,3-dimethylimidazolinium salts⁷ and have been investigating the metal-catalyzed synthesis of substituted-naphthol derivatives using these products.⁸ Pd(OAc)₂ is found to be an efficient catalyst for several coupling reactions of 1,2-diazonaphthoquinones.^{8a,b} During the evaluation of the stability of 2-diazonaphthoquinone **6a**⁹ in the presence of a catalytic amount of Pd(OAc)₂ in various solvents (CH₂Cl₂, toluene, benzene, CH₃CN, and THF) at reflux, the unexpected formation of 8-membered cyclic ether **7a** and 13-membered cyclic ether **8a** was observed in THF. We were intrigued by this uncommon formation of medium-sized rings and anticipated that this transformation could be a new method for the synthesis of protected 1,2-naphthalenediols. Similar to catechol derivatives, 1,2-naphthalenediols are attractive candidates as aromatic functional materials (or their building blocks), such as for use in solar cells, metal ligands, and antioxidants.¹⁰ However, to date, only a few synthetically useful processes have been reported for the synthesis of 1,2-naphthalenediols.¹¹ Therefore, the generality and efficiency of this Pd(OAc)₂-catalyzed cyclization of 1,2-diazonaphthoquinones with THF were explored. In addition, the Pd(II) catalyst was efficient for the macrocyclization of 1,2-diazonaphthoquinones and oxetane.

Initially, the reaction of 2-diazonaphthoquinone **6a** and THF was examined under several reaction conditions (Table 1). First, several palladium catalysts were examined (Table 1, entries 1–8). When 10 mol % Pd(OAc)₂ was used, **6a** was consumed within 10 min, and cyclic ethers **7a** and **8a** were

Received: December 6, 2013

Published: March 3, 2014

Table 1. Metal-Catalyzed Macrocyclization of 2-Diazonaphthoquinone 6a with THF^a


entry	metal cat.	time	7a (%)	8a (%)
1	Pd(OAc) ₂ (10 mol %)	10 min	29	26
2	Pd(OAc) ₂ (5 mol %)	40 min	27	28
3	Pd(OAc) ₂ (2.5 mol %)	1.5 h	33	36
4	Pd(OAc) ₂ (1 mol %)	6.5 h	13	9
5 ^b	Pd(OAc) ₂ (2.5 mol %)	3.5 h	21	31
6 ^c	Pd(OAc) ₂ (2.5 mol %)	2.5 h	0	0
7	PdCl ₂ (2.5 mol %)	2.5 h	20	21
8	Pd(OCOCF ₃) ₂ (2.5 mol %)	1 h	21	25
9	Rh ₂ (OAc) ₄ (1 mol %)	3 h	trace ^d	trace
10	Rh ₂ (OCOCF ₃) ₄ (1 mol %)	2.5 h	4 ^e	13
11	Rh ₂ (Oct) ₄ (1 mol %) ^f	2 h	11 ^g	5
12	Cu(OAc) ₂ (2.5 mol %)	3 h	0 ^h	0
13	CuCl ₂ (2.5 mol %)	2.5 h	0 ⁱ	0
14	Cu(OTf) ₂ (2.5 mol %)	3 h	0 ^j	0
15	Cu(OTf)·C ₆ H ₆ (2.5 mol %)	2.5 h	0 ^k	0

^aReaction conditions: **6a** (0.5 mmol) in THF (2 mL) at reflux. ^bThe reaction was performed at 45 °C. ^cLiCl (1.0 equiv) was added. ^d**6a** was recovered in 28% yield. ^e**6a** was recovered in 19% yield. ^fRh₂(OCOC₇H₁₅)₄. ^g**6a** was recovered in 39% yield. ^h**6a** was recovered in 92% yield. ⁱ**6a** was recovered in 80% yield. ^j**9** was obtained in 9% yield. **6a** was recovered in 32% yield. ^k**9** was obtained in 3% yield. **6a** was recovered in 25% yield.

obtained in 29% and 26% yields, respectively. As shown in Table 1, entries 1–4, the amount of Pd(OAc)₂ could be reduced to 2.5 mol % (Table 1, entry 3). At 45 °C, the yields of


the cyclization products were decreased slightly (Table 1, entry 5). The addition of a halide anion has been reported to be effective in some Pd-catalyzed reactions.¹² Although **6a** was consumed with 2.5 mol % Pd(OAc)₂ in the presence of an equimolar amount of LiCl, cyclic ethers **7a** and **8a** were not obtained (Table 1, entry 6). Use of PdCl₂ and Pd(OCOCF₃)₂ resulted in low yields of both products (Table 1, entries 7 and 8).

Interestingly, while only the Rh catalysts were reported to be effective for the metal-catalyzed macrocyclization reaction of cyclic ethers and α -carbonylcarbenes,^{5,6} they were not efficient for the cyclization of **6a** and THF (Table 1, entries 9–11).

The series of Cu reagents examined for the reaction, including Cu(OAc)₂, CuCl₂, Cu(OTf)₂, and Cu(OTf)·C₆H₆, also did not provide the cyclized products **7a** and **8a** (Table 1, entries 12–15), although a minor amount of C–H insertion product **9** was formed in the reactions with Cu(OTf)₂ and Cu(OTf)·C₆H₆ (Table 1, entries 14 and 15, respectively).

On the basis of these results, the optimum cyclization conditions were determined to be Pd(OAc)₂ (2.5 mol %) in THF at reflux.

Next, the scope and limitations of the Pd(OAc)₂-catalyzed cyclization of 1,2-diazonaphthoquinones and THF were examined (Table 2). The substrate 2-diazonaphthoquinone **6b** bearing an electron-withdrawing group at the C-4 position gave cyclization products **7b** and **8b** in 16% and 21% yields, respectively (Table 2, entry 1). On the other hand, 4-methoxy-2-diazonaphthoquinone **6c** was completely consumed within 20 min, but the expected cyclization products were not generated, and dimerization product **11** was obtained in 10% yield (Table 2, entry 2). Introduction of a C-3 substituent was, however, efficient for the selective formation of 8-membered ring cyclic ethers **7**, as shown in Table 2, entries 3 and 4. The reactions of 1-diazonaphthoquinones **10**⁹ also proceeded smoothly and selectively to give 8-membered cyclic ethers **7** in good to high yields (Table 2, entries 5–9). Simple 1-diazonaphthoquinone **10a** gave **7a** in 45% yield, accompanied by 2% of the 13-membered cyclic ether **8a** (Table 2, entry 5). The reaction of 6-bromo-1-diazonaphthoquinone **10b** gave 8-membered ether **7f**

Table 2. Pd(OAc)₂-Catalyzed Macrocyclization of 1,2-Diazonaphthoquinones 6 and 10 with THF^a



entry	R ¹	R ²	R ³	6 or 10	time	7 (%)	8 (%)
1	H	Cl	H	6b	1 h	7b 16	8b 21
2	H	OMe	H	6c	20 min	7c 0 ^b	8c 0
3	CH ₂ OTBS	H	H	6d	2 h	7d 60	8d 0
4	CO ₂ Me	H	H	6e	1 h	7e 55	8e 5
5	H	H	H	10a	1.5 h	7a 45	8a 2
6	H	H	Br	10b	1.5 h	7f 53	8f 0
7	CH ₂ OTBS	H	H	10c	2 h	7d 78	8d 0
8	CO ₂ Me	H	H	10d	20 min	7e 84	8e 0
9	CO ₂ Ph	H	H	10e	1.5 h	7g 35	8g 0

^aReaction conditions: **6** or **10** (0.5 mmol), Pd(OAc)₂ (2.5 mol %) in THF (2 mL) at reflux. ^b**11** was obtained in 10% yield.

as the sole product in 53% yield (Table 2, entry 6). With these substrates, introduction of an appropriate substituent at the C-3 position also clearly improved the yield of cyclic ether **7** (Table 2, entries 7 and 8). In the reaction of 3-siloxymethyl-1-diazonaphthoquinone **10c**, cyclic ether **7d** was formed in 78% yield (Table 2, entry 7), and the reaction of 3-methoxycarbonyl-1-diazonaphthoquinone **10d** gave 8-membered cyclic ether **7e** in 84% yield (Table 2, entry 8). However, the corresponding phenyl ester **10e** afforded cyclic ether **7g** in a lower yield (35%) (Table 2, entry 9).

We also examined the Pd(OAc)₂-catalyzed reaction of 1,2-diazonaphthoquinones and oxetane (Table 3). In the reaction

Table 3. Pd(OAc)₂-Catalyzed Macrocyclization of 1,2-Diazonaphthoquinones **6a** and **10a** with Oxetane^a



entry	6a or 10a	time	12 (%)	13 (%)
1	6a	3 h	66	0
2 ^b	6a	3.5 h	0 ^c	0
3	10a	20 h	26 ^d	5


^aReaction conditions: **6a** or **10a** (0.5 mmol), Pd(OAc)₂ (2.5 mol %) in oxetane (2 mL) at reflux. ^bRh₂(OAc)₄ (1 mol %) was used instead of Pd(OAc)₂. ^c**6a** was recovered in 84% yield. ^d**10a** was recovered in 34% yield.

of 2-diazonaphthoquinone **6a**, 15-membered cyclic ether **12** was formed in 66% yield after stirring for 3 h at reflux.¹³ A similar Rh₂(OAc)₄-catalyzed reaction has been reported for 2-diazo-1,3-dicarbonyl compounds.^{6b} However, Rh₂(OAc)₄ was ineffective for the cyclization of diazonaphthoquinone **6a**, as shown in Table 3, entry 2. In addition, the reaction of 1-diazonaphthoquinone **10a** was slower than that of 2-diazonaphthoquinone **6a**, and 15-membered product **12** was formed in just 26% yield along with acyclic ether **13** (5%) after stirring for 20 h (Table 3, entry 3).


In addition, the 1,2-diazonaphthoquinones also reacted with tetrahydropyran (THP) in the presence of a catalytic amount of Pd(OAc)₂ to afford a mixture of 9-membered cyclic ether **14** and 15-membered ether **15** (Scheme 2).

In Scheme 3, a possible reaction mechanism is depicted for the Pd(OAc)₂-catalyzed formation of medium-sized cyclic ethers (**7**, **14**) and macrocyclic ethers (**8**, **15**) from 2-diazonaphthoquinone **6** in cyclic ether (THF, THP). First, Pd(OAc)₂ reacts with diazonaphthoquinone **6** to form Pd(II) carbene complex **I**.¹⁴ In the case of the reaction in THF, nucleophilic attack of THF on carbene complex **I** proceeds to form oxonium ylide **II**,¹⁵ which may be aromatic palladium naphtholate **II'**. Successively, **II** reacts with THF giving **III**. Since 5-*endo*-tet cyclization is highly disfavored commonly,¹⁶ 8-membered cyclic ether **7** is not formed directly from **II**, but is formed by 8-*exo*-tet cyclization of **III** (Path A). Macrocyclic ether **8** is formed from the same intermediate **III** by 10-*endo*-tet

Scheme 2. Pd(OAc)₂-Catalyzed Cyclization of 1,2-Diazonaphthoquinones **6a** and **10a** with THP



Scheme 3. Possible Reaction Mechanism



cyclization (Path B). The C-3 substituted substance R¹ can more easily attain the required conformation of **III** for 8-*exo*-cyclization as compared to the C-3 unsubstituted substance. For the reaction of 1-diazonaphthoquinone **10** and THF, the hydrogen at the C-8 position in **10** is assumed to function similarly to C-3 substituent R¹ in 2-diazonaphthoquinone **6**, resulting in the selective formation of *exo*-cyclization product **7**.

Compared to the reaction with THF, the reaction with THP gave lower yields of cyclic ethers, and different selectivity in the formation of cyclic ethers (medium-sized cyclic ether/macro cyclic ether) was observed, which would be attributed to the cyclization mode, that is 8-*exo*-tet/10-*endo*-tet cyclization for THF vs 9-*exo*-tet/11-*endo*-tet cyclization for THP.

In conclusion, we developed a new method for the synthesis of medium-sized/macrocyclic ethers via the reaction of diazonaphthoquinones in the presence of a catalytic amount of Pd(OAc)₂. This transformation also serves as an efficient method for the preparation of protected 1,2-naphthalenediols.

Further studies on the Pd(OAc)₂-catalyzed cyclization reaction of diazonaphthoquinones with heterocyclic compounds are in progress.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and characterization data, including ¹H and ¹³C NMR spectra for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: kita@che.kyutech.ac.jp.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Support for this work by the Naito Foundation is gratefully acknowledged.

■ REFERENCES

- (1) For reviews, see: (a) Ye, T.; McKervey, M. A. *Chem. Rev.* **1994**, 94, 1091. (b) Doyle, M. P.; Ye, T.; McKervey, M. A. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*; John Wiley & Sons: New York, 1998. (c) Zhang, Z.; Wang, J. *Tetrahedron* **2008**, 64, 6577.
- (2) Nozaki, H.; Takaya, H.; Noyori, R. *Tetrahedron* **1966**, 22, 3393.
- (3) (a) Ihara, E.; Saiki, K.; Goto, Y.; Itoh, T.; Inoue, K. *Macromolecules* **2010**, 43, 4589. (b) Ihara, E.; Hara, Y.; Itoh, T.; Inoue, K. *Macromolecules* **2011**, 44, 5955.
- (4) Kirmse, W.; Lelgemann, R. *Chem. Ber.* **1991**, 124, 1865.
- (5) (a) Cenini, S.; Cravotto, G.; Giovenzana, G. B.; Palmisano, G.; Tollari, S. *Tetrahedron* **1999**, 55, 6577. (b) Ballesteros-Garrido, R.; Rix, D.; Besnard, C.; Lacour, J. *Chem.—Eur. J.* **2012**, 18, 6626.
- (6) (a) Zeghida, W.; Besnard, C.; Lacour, J. *Angew. Chem., Int. Ed.* **2010**, 49, 7253. (b) Rix, D.; Ballesteros-Garrido, R.; Zeghida, W.; Besnard, C.; Lacour, J. *Angew. Chem., Int. Ed.* **2011**, 50, 7308.
- (7) Kitamura, M.; Tashiro, N.; Sakata, R.; Okauchi, T. *Synlett* **2010**, 2503.
- (8) (a) Kitamura, M.; Sakata, R.; Okauchi, T. *Tetrahedron Lett.* **2011**, 52, 1931. (b) Kitamura, M.; Kisanuki, M.; Sakata, R.; Okauchi, T. *Chem. Lett.* **2011**, 40, 1129. (c) Kitamura, M.; Kisanuki, M.; Okauchi, T. *Eur. J. Org. Chem.* **2012**, 905. (d) Kitamura, M.; Araki, K.; Matsuzaki, H.; Okauchi, T. *Eur. J. Org. Chem.* **2013**, 5045.
- (9) In this manuscript, 2-diazonaphthoquinone means 2-diazo-1(2H)-naphthalenone derivative and 1-diazonaphthoquinone means 1-diazo-2(1H)-naphthalenone derivative.
- (10) (a) Stahl, P.; Kissau, L.; Mazitschek, R.; Huwe, A.; Furet, P.; Giannis, A.; Waldmann, H. *J. Am. Chem. Soc.* **2001**, 123, 11586. (b) Lu, T.; Shao, P.; Mathew, I.; Sand, A.; Sun, W. *J. Am. Chem. Soc.* **2008**, 130, 15782. (c) Madan, S.; Cheng, C. *J. Org. Chem.* **2006**, 71, 8312.
- (11) (a) Platt, K. L.; Oesch, F. *J. Org. Chem.* **1983**, 48, 265. (b) Zambrano, J. L.; Dorta, R. *Synlett* **2003**, 1545. (c) Crandall, J. K.; Zucco, M.; Kirsch, R. S.; Coppert, D. M. *Tetrahedron Lett.* **1991**, 32, 5441.
- (12) Fagnou, K.; Lautens, M. *Angew. Chem., Int. Ed.* **2002**, 41, 26.
- (13) For a photoinduced similar macrocyclization of the diazoquinone and oxetane, see ref 4. The maximum yield of the macrocyclic compound was 16%.
- (14) (a) Anciaux, A. J.; Hubert, A. J.; Noels, A. F.; Petiniot, N.; Teyssie, P. *J. Org. Chem.* **1980**, 45, 695. (b) Denmark, S. E.; Stavenger, R. A.; Faucher, A.-M.; Edwards, J. P. *J. Org. Chem.* **1997**, 62, 3375. (c) Greenman, K. L.; Carter, D. S.; Van Vranken, D. L. *Tetrahedron* **2001**, 57, 5219. (d) Bröring, M.; Brandt, C. D.; Stellwag, S. *Chem. Commun.* **2003**, 2344. (e) Albéniz, A. C.; Espinet, P.; Manrique, R.; Pérez-Mateo, A. *Chem.—Eur. J.* **2005**, 11, 1565. (f) Greenman, K. L.; Van Vranken, D. L. *Tetrahedron* **2005**, 61, 6438. (g) López-Alberca, M. P.; Mancheño, M. J.; Fernández, I.; Gómez-Gallego, M.; Sierra, M. A.; Torres, R. *Org. Lett.* **2007**, 9, 1757. (h) Devine, S. K. J.; Van Vranken, D. L. *Org. Lett.* **2007**, 9, 2047. (i) Barluenga, J.; Moriel, P.; Valdés, C.; Aznar, F. *Angew. Chem., Int. Ed.* **2007**, 46, 5587. (j) Chen, S.; Wang, J. *Chem. Commun.* **2008**, 4198. (k) Peng, C.; Wang, Y.; Wang, J. *J. Am. Chem. Soc.* **2008**, 130, 1566. (l) Xiao, Q.; Ma, J.; Yang, Y.; Zhang, Y.; Wang, J. *Org. Lett.* **2009**, 11, 4732. (m) Zhang, Z.; Liu, Y.; Gong, M.; Zhao, X.; Zhang, Y.; Wang, J. *Angew. Chem., Int. Ed.* **2010**, 49, 1139. (n) Zhang, Z.; Liu, Y.; Ling, L.; Li, Y.; Dong, Y.; Gong, M.; Zhao, X.; Zhang, Y.; Wang, J. *J. Am. Chem. Soc.* **2011**, 133, 4330. (o) Chen, Z.-S.; Duan, X.-H.; Zhou, P.-X.; Ali, S.; Luo, J.-Y.; Liang, Y.-M. *Angew. Chem., Int. Ed.* **2012**, 51, 1370. (p) Xiao, Q.; Wang, B.; Tian, L.; Yang, Y.; Ma, J.; Zhang, Y.; Chen, S.; Wang, J. *Angew. Chem., Int. Ed.* **2013**, 52, 9305. (q) Yang, K.; Zhang, J.; Cheng, B.; Zhao, L.; Zhai, H. *Org. Lett.* **2013**, 15, 808. (r) Khanna, A.; Premachandra, I. D. U. A.; Sung, P. D.; Van Vranken, D. L. *Org. Lett.* **2013**, 15, 3158. (s) Khanna, A.; Premachandra, I. D. U. A.; Sung, P. D.; Van Vranken, D. L. *Org. Lett.* **2013**, 15, 3694. (15) Padwa, A.; Hornbuckle, S. F. *Chem. Rev.* **1991**, 91, 263. (16) Baldwin, J. E. *J. Chem. Soc., Chem. Commun.* **1976**, 734.