

Notes

Tissue Distribution of Technetium-99m and Carbon-14 Labeled *N*-(2,6-Dimethylphenylcarbamoylmethyl)iminodiacetic Acid


Patrick S. Callery,* William C. Faith, Michael D. Loberg, Anna T. Fields, Elizabeth B. Harvey, and Malcolm D. Cooper

Department of Medicinal Chemistry, School of Pharmacy, and the Division of Nuclear Medicine, School of Medicine, University of Maryland, Baltimore, Maryland 21201. Received March 4, 1976

The synthesis, radiochemical labeling, and tissue distribution characteristics of *N*-(2,6-dimethylphenylcarbamoylmethyl)iminodiacetic acid are described. The radiopharmaceutical prepared by labeling with ^{99m}Tc was rapidly eliminated through the hepato-biliary system of mice. Parent ^{14}C compound was eliminated primarily through the kidney. The ^{99m}Tc ion appears to have a greater influence than the organic carrier molecule on the distribution of the radiopharmaceutical.

Numerous radiopharmaceutical agents composed of radioactive metal ions bound to chelating compounds have been developed as organ-imaging agents in nuclear medicine. Generally, these agents consist of a combination of γ -emitting radionuclidic cations with existing organic compounds which have established complexing properties. For example, the tetracyclines,¹ DTPA,² and other compounds³ have been chelated or complexed with reduced technetium-99m to form radiopharmaceuticals. Sundberg et al.⁴ have coupled an EDTA derivative to proteins for the purpose of binding radioactive metals to a macromolecule-bound chelating group.

Incorporation of a chelating moiety into a relatively low-molecular-weight drug analogue may create an agent capable of binding with technetium-99m while retaining the biological actions and tissue distribution characteristics of the parent drug. Iminodiacetic acid (1) is a metal



complexing moiety which strongly binds transition metals⁵ and can be readily incorporated into organic molecules. We wish to report the synthesis and biological distribution of an iminodiacetic acid containing radiopharmaceutical agent which is structurally related to the antiarrhythmic drug lidocaine (2).⁶

Results and Discussion

Substitution of the diethylamino group of 2 with iminodiacetic acid produced the drug-chelate analogue 3 which is capable of complexing with ^{99m}Tc and other organ-imaging radionuclides. Carrier compound 3 was synthesized by displacement of the halogen atom of the chloroacetyl compound 4 with the disodium salt of iminodiacetic acid (1). A complex between reduced technetium, prepared by stannous reduction of TcO_4^- , and 3 was formed under acidic conditions. Paper chromatography with physiological saline and paper electrophoresis of the Tc-3 chelate showed the absence of radioactive bands associated with pertechnetate or technetium-tin

colloid. While the exact radiochemical structure of Tc-3 remains to be determined, the paper electrophoretic data, which shows a single sharp peak with a migration distance intermediate between that of colloid and pertechnetate, suggests that Tc-3 exists in one discrete radiochemical form.

The tissue distribution of Tc-3 in mice was determined at various time intervals following intravenous administration. Table I shows the percent injected dose of radioactivity in blood, liver, intestine, and kidney. The technetium was rapidly extracted from the blood by the liver and was ultimately eliminated through the hepato-biliary system. After 60 min, a large portion of the dose (76%) was accounted for in the liver, gall bladder, and intestine. Urinary excretion of label was found to be a minor pathway for the elimination of Tc-3 in mice and dogs.⁷

The dissimilarity in tissue distribution and blood concentrations of Tc-3 compared to literature values for [^{14}C]lidocaine⁸ led to studies on the influence of the metal ion on the disposition of Tc-3. A comparison of the distribution of 3 with and without the technetium label was undertaken with the aid of [^{14}C]-3. The ^{14}C -label introduced into 3 was located in the methylene groups of the iminodiacetic acid moiety and was produced by an alternate pathway to 3 (see Experimental Section, method B). Amino compound 5, prepared by the action of ammonia on 4, was treated with chloroacetic acid- ^{14}C to produce [^{14}C]-3 with a specific activity of 1.12 $\mu\text{Ci}/\text{mg}$. Tissue distribution determinations for [^{14}C]-3 in mice were carried out by a similar procedure using the same milligram quantity as in the Tc-3 experiments. Clearance of [^{14}C]-3 from the blood was as rapid as Tc-3 blood clearance.

In Table I is shown the percent ^{14}C dose in blood, liver, intestine, and kidney. A comparison of the major routes of elimination of Tc-3 and [^{14}C]-3 in mice is shown in Figure 1. The main elimination pathway of radioactivity after administration of [^{14}C]-3 was via the kidney, which is more in keeping with the urinary elimination of lidocaine⁹ and its metabolites¹⁰ than with the hepato-biliary fate of Tc-3. At 60 min after injection, most (74%) of the ^{14}C dose was accounted for in the urine with less than 2% found in the liver or intestines.

Evidence supporting that a Tc-3 complex was being monitored and that extensive dissociation in vivo of Tc-3 to colloidal technetium or pertechnetate had not occurred was gained by a comparison of the distribution of Tc-3 to

Table I. Percent Injected Radioactivity at Various Times in Tissues of Mice Following Intravenous Administration of Tc-3 and [¹⁴C]-3^a

	Time, min	Blood	Liver	Intestine	Kidney	Urine
Tc-3	5	3.3 ^b (2.6-3.8 ^c)	28.2 (21.3-44.2)	38.3 (27.8-46.3)	1.8 (1.1-3.0)	
	30	1.1 (0.6-1.8)	10.2 (3.4-14.7)	60.2 (47.6-70.1)	0.6 (0.2-1.0)	
	60	0.8 (0.3-1.4)	8.0 (0.9-19.2)	68.3 (54.1-73.8)	0.8 (0.6-1.3)	
[¹⁴ C]-3	5	7.9 ^b (6.8, 8.9 ^d)	5.7 (6.1, 5.2)	3.8 (4.0, 3.6)	7.5 (7.6, 7.4)	31.9 (34.6, 29.3)
	30	0.5 (0.3, 0.7)	1.3 (1.0, 1.6)	1.8 (1.5, 2.1)	1.0 (0.3, 1.8)	74.2 (61.0-82.7 ^c)
	60	0.2 (0.1, 0.2)	0.4 (0.4, 0.5)	1.4 (1.5, 1.3)	1.7 (3.3, 0.2)	74.0 (58.9-82.8 ^c)

^a Mice were injected in the tail vein with Tc-3 (ca. 0.5 μ Ci) or [¹⁴C]-3 (2.23 μ Ci). Radioactivity in lung, spleen, or heart was less than 1% of the injected dose. In the Tc experiments, less than 2% of the dose was accounted for in the stomach.
^b Represents percent dose average value. ^c Represents range of four mice. ^d Represents mouse 1 and 2, respectively.




Figure 1. Major pathways of elimination of radioactivity in mice following intravenous injection of Tc-3 and [¹⁴C]-3. Each value is in terms of the total injected radioactive dose and represents an average of at least two mice. The hepato-biliary clearance of Tc (•-•-•) consists of the combined counts found in liver, gall bladder, and intestines. The urinary excretion of ¹⁴C (■-■-■) includes the urinary bladder and contents.

the literature values for colloidal technetium and pertechnetate distribution. The absence of significant quantities of radioactivity in the stomach and spleen indicates that the distribution of Tc-3 is different from the distribution of either pertechnetate¹¹ or colloidal technetium.¹² Furthermore, the short transit time of Tc-3 through the liver is inconsistent with the biological distribution of technetium-tin colloid which is known to have a biological half-life in the liver in excess of 2 days.¹² This suggests that the technetium remains in chelate form during its course through the animal. Additionally, the chromatographic and tissue distribution characteristics of the technetium eliminated in the bile 1 h after Tc-3 administration were evaluated as a measure of the in vivo stability of Tc-3. Paper chromatography of the radioactive biliary contents in saline produced a chromatogram identical with that of Tc-3 before injection. Tissue distribution studies obtained from reinjection of the radioactive biliary contents into a second set of mice showed a pattern well correlated with Tc-3 distribution.⁷ The distribution pattern of Tc-3 along with the properties of the technetium excreted in the bile after Tc-3 administration strongly suggests that the Tc-3 chelate remains intact in the animal and is eliminated in an unchanged form. The possibility of metabolism of Tc-3 to Tc-labeled metabolites of **3** which undergo biliary excretion has not been studied further.

The difference in fate of Tc-3 compared to [¹⁴C]-3 may be related to the formation of the radioactive metal-3 chelate. Although the effect of the technetium atom on the electronic and spatial characteristics of **3** and the possibility of dimer or trimer formation have yet to be

studied, it seems reasonable that, in the chelate, donation of the nitrogen unshared electron pair to the electropositive technetium atom is likely. The resulting partially positive iminodiacetic acid nitrogen or the positive character of the reduced technetium may possess properties similar to a cationic quaternary nitrogen and may play a role in the rapid hepato-biliary excretion of Tc-3. A relatively nonspecific active transport system which is well described for the excretion of cations¹³ may be responsible for the biliary elimination of a number of quaternary ammonium antiarrhythmic compounds.¹⁴ Tc-3 may be eliminated in the bile by such a transport process while [¹⁴C]-3 follows the usual urinary pathway for the excretion of polar substances.

Knowledge of the structural features which influence biological disposition is essential in the design of organ-imaging radiopharmaceuticals.¹⁵ By radiolabeling the carrier molecule **3** and preparing the Tc-3 chelate for distribution studies, we have shown that the fate of **3** is dramatically different from the disposition of Tc-3. The strong influence of the radiochemical ion on the target organ specificity of Tc-3 should be taken into account in the design of radiopharmaceuticals containing an iminodiacetic acid moiety.

Experimental Section

Melting points were determined on a Thomas-Hoover melting point apparatus and are uncorrected. Microanalyses were performed by Galbraith Laboratories, Inc., Knoxville, Tenn. ¹H NMR spectra were obtained on a Jeolco C-60HL NMR spectrometer and the chemical shifts are expressed in parts per million relative to Me₄Si or sodium 2,2-dimethyl-2-silapentane-5-sulfonate. Electron impact mass spectra were obtained with a Du Pont 21-490 mass spectrometer operating at 70 eV. A Beckman paper electrophoresis cell (Durrum Type) Model R-Series D system was used for paper electrophoresis. Chromatograms were scanned with a Packard Model 385 radiochromatogram scanner. Liquid scintillation counting and γ counting of samples were performed on a Packard Tri-Carb Model 2002 spectrometer.

ω -Chloro-2,6-dimethylacetanilide (4). Following the procedure of Lofgren,¹⁶ 2,6-dimethylaniline (4.85 g, 40.0 mmol) in glacial HOAc (30 ml) was treated with chloroacetyl chloride (4.97 g, 44.0 mmol) to yield 6.1 g (77%) of pure **4**: mp 147-149° (lit.¹⁷ mp 145-146°); ¹H NMR (CDCl₃) δ 2.21 (s, 6, CH₃), 4.18 (s, 2, CH₂Cl), and 7.10 ppm (s, 3, aromatic protons); mass spectrum *m/e* (rel intensity) 197 (55, M⁺), 148 (100, M⁺ - CH₂Cl).

ω -Amino-2,6-dimethylacetanilide Hydrochloride (5-HCl). A solution of **4** (4.4 g, 22.0 mmol) in 135 ml of absolute EtOH saturated with NH₃ was stirred at room temperature for 3 days. The mixture was evaporated to dryness and the residue was dissolved in 0.1 N HCl (50 ml) and then extracted with two 50-ml portions of CHCl₃. The aqueous layer was adjusted to pH 10 with 1 N NaOH and extracted with CHCl₃ (5 \times 50 ml) to yield, after evaporation of the combined CHCl₃ layers, crude **5** (1.1 g, 28%, mp 78-80°). The HCl salt of **5** was prepared by bubbling HCl gas through an EtOH-Et₂O solution of **5**. Recrystallization from EtOH gave an analytical sample: mp 302° dec; ¹H NMR (Me₂SO-*d*₆) δ 2.24 (s, 6, CH₃), 3.92 (s, 2, CH₂), and 7.22 ppm (s,

3, aromatic protons); mass spectrum m/e (rel intensity) 178 (100, M^+). Anal. ($C_{10}H_{15}N_2OCl$) C, H, N.

***N*-(2,6-Dimethylphenylcarbamoylmethyl)iminodiacetic Acid (3).** **Method A.** A solution of 4 (2.0 g, 10.0 mmol) and disodium iminodiacetic acid (1.8 g, 10.0 mmol) in EtOH-H₂O (3:1) was held at reflux for 48 h. The mixture was cooled and then evaporated to dryness to leave a yellow solid. The residue in 25 ml of H₂O was extracted with Et₂O (3 × 25 ml). The pH of the aqueous layer was then adjusted to 3.2 with concentrated HCl and the precipitate which formed on cooling was recrystallized from H₂O to give 0.5 g (20%) of 3: mp 215–216°; ¹H NMR (Me_2SO-d_6) δ 2.20 (s, 6, CH₃), 3.57 (s, 2, NCOCH₂N), 3.63 (s, 4, CH₂COO), 7.19 ppm (s, 3, aromatic protons). Anal. ($C_{14}H_{18}N_2O_5$) C, H, N.

Method B. A solution of 5 (1.1 g, 5.0 mmol) in 95% EtOH (50 ml) was added dropwise to a stirred solution of chloroacetic acid (0.95 g, 10.0 mmol) in 95% EtOH (50 ml). During the addition, the pH of the mixture was maintained above 7 with 4 N NaOH. The resulting mixture was heated to reflux for 24 h. The solvents were evaporated and the residue was recrystallized from H₂O buffered to pH 3 to give the final product (0.32 g, 25%) which gave ir and NMR spectra and melting point identical with a sample of 3 produced by method A. Anal. ($C_{14}H_{18}N_2O_5$) C, H, N.

***N*-(2,6-Dimethylphenylcarbamoylmethyl)iminodiacetic Acid-methylene-¹⁴C ([¹⁴C]-3).** Following the procedure of method B, a solution of 5 (147 mg, 0.69 mmol) in 95% EtOH (15 ml) was allowed to react with chloroacetic acid- α -¹⁴C (250 μ Ci/mg, Amersham Searle) diluted with carrier chloroacetic acid (147 mg, 1.5 mmol). The crude product was recrystallized from H₂O to give 104 mg (51%) of [¹⁴C]-3: mp 185–190°; 1.12 μ Ci/mg; chemical and radiochemical purity was confirmed by TLC on silica gel with BuOH-HOAc-H₂O (4:1:1) and CHCl₃-EtOH (1:1).

***N*-(2,6-Dimethylphenylcarbamoylmethyl)iminodiacetic Acid ^{99m}Tc Chelate (Tc-3).** A solution of 3 (0.15 g, 0.51 mmol) in 3 ml of 0.1 N NaOH was adjusted to pH 3.5 with 1 N HCl. To this mixture was added 0.3 ml of a 1 N HCl solution containing SnCl₂ (0.6 mg, 3.3 μ mol) while maintaining a pH of 3.5 by the dropwise addition of 1 N NaOH. After 5 min at room temperature, 5–50 μ Ci of ^{99m}TcO₄⁻ was added.

Radiochemical Purity. The radiochemical purity of Tc-3 was examined by ascending paper chromatography in physiological saline and by paper electrophoresis for 1.5 h at a constant voltage of 300 V in a 0.05 M phosphate buffer, pH 6.8. The radiochromatograms of Tc-3 were compared with those found for both the pertechnetate anion and for technetium-tin colloid prepared by the method of Lin and Winchell¹² and by the method used to prepare Tc-3 except that 3 was omitted from the reaction mixture. The two colloids appeared indistinguishable under both analytical systems. Paper chromatography yielded R_f values of 0.0, 0.74, and 0.95–1.0 for technetium-tin colloid, pertechnetate, and Tc-3, respectively. When subjected to paper electrophoresis, technetium-tin colloid, pertechnetate, and Tc-3 were observed to migrate 0.0, 7.3, and 3.2 cm, respectively. Less than 0.1% of Tc-3 was present as either technetium-tin colloid or as pertechnetate. Greater than 95% of the radioactivity was recovered after filtration of the Tc-3 solution through a 0.22- μ membrane filter (Millipore Corp.). Radiochemical analysis as above of the filtrate was identical with unfiltered Tc-3.

Tissue Distribution Studies. Male Swiss-Webster mice (20–25 g) were administered [¹⁴C]-3 or Tc-3 intravenously in the tail vein. Each mouse received a dose of 2.0 mg (4.96×10^6 dpm) of [¹⁴C]-3 in 0.25 ml of distilled H₂O adjusted to pH 5.7 with 1 N NaOH or 0.1–1.0 μ Ci of Tc-3 in 0.1 ml of the pH 3.5 SnCl₂ solution described above. Liver, lung, kidneys, spleen, heart, intestines, tail, and urine were counted at 5, 30, and 60 min after injection. In the ¹⁴C experiments, mice were clamped to prevent urination, and urinary bladders plus contents were counted.

Counts in the tail in all cases were found to be less than 5% of the injected dose and were subtracted from the total dose to obtain the total injected dose. Organ and blood samples in the Tc experiments were counted in test tubes. For ¹⁴C samples lung, kidney, spleen, and heart (100–200 mg) were solubilized with 2 ml of NCS (Amersham Searle) and 2 drops of H₂O in liquid scintillation vials with plastic tops by heating to 50° in a water bath overnight. The resulting solutions were counted in 13 ml of a toluene cocktail [0.2 mM POPOP (Packard) and 27 mM PPO (Packard) in reagent grade toluene]. All ¹⁴C values were corrected for efficiency using a toluene-¹⁴C internal standard. Blood was collected at the time of sacrifice using a 200- μ l capillary tube. A syringe containing NCS (2.0 ml) was used to wash the blood from the tube into a scintillation vial. After heating to 50° in a water bath overnight, 13 ml of cocktail was added and the samples were bleached with 0.6 ml of a 0.8 M solution of benzoyl peroxide in toluene. Additional NCS was sometimes necessary to afford a clear solution. Lung and spleen samples were bleached with 0.3 ml of benzoyl peroxide solution. Liver and intestine samples were homogenized in a 4:1 water to organ mixture with a Potter-Elvehjem tissue homogenizer and 0.3-ml aliquots were solubilized for counting. The urinary bladders containing urine were solubilized with NCS and aliquots were counted.

References and Notes

- (1) M. K. Dewanjee, C. Fliegel, S. Treves, and M. A. Davis, *J. Nucl. Med.*, **15**, 176 (1974).
- (2) W. Hauser, H. L. Atkins, K. G. Nelson, and P. Richards, *Radiology*, **94**, 679 (1970).
- (3) M. S. Lin, D. A. Goodwin, and S. L. Kruse, *J. Nucl. Med.*, **15**, 338 (1974); T. Higashi, M. Kanno, and K. Tomura, *ibid.*, **15**, 1167 (1974).
- (4) M. V. Sundberg, C. F. Meares, D. A. Goodwin, and C. I. Diamanti, *J. Med. Chem.*, **17**, 1304 (1974).
- (5) A. E. Martell and M. Calvin, "Chemistry of the Metal Chelate Compounds", Prentice-Hall, New York, N.Y., 1952, p 534.
- (6) Presented in part before the 169th National Meeting of the American Chemical Society, Philadelphia, Pa., April 1975, and the 22nd Annual Meeting of the Society of Nuclear Medicine, Philadelphia, Pa., June 1975.
- (7) M. Loberg, M. Cooper, E. Harvey, P. Callery, and W. Faith, *J. Nucl. Med.*, in press.
- (8) B. Akerman, A. Astrom, S. Ross, and A. Telc, *Acta Pharmacol. Toxicol.*, **24**, 389 (1966).
- (9) C. Y. Sung and A. P. Truant, *J. Pharmacol. Exp. Ther.*, **112**, 432 (1954).
- (10) J. B. Keenaghan and R. N. Boyes, *J. Pharmacol. Exp. Ther.*, **180**, 454 (1972).
- (11) P. V. Harper, K. A. Lathrop, and A. Gottschalk in "Radioactive Pharmaceuticals", G. A. Andrews, R. M. Kinsey, and N. N. Wagner, Jr., Ed., Symposium Series 6, CONF-651111, National Bureau of Standards, Springfield, Va., 1966, p 335.
- (12) M. S. Lin and H. S. Winchell, *J. Nucl. Med.*, **13**, 58 (1972).
- (13) A. Ryrfeldt and E. Hansson, *Acta Pharmacol. Toxicol.*, **30**, 59 (1971).
- (14) L. S. Shanker and H. M. Solomon, *Am. J. Physiol.*, **204**, 829 (1963); J. E. Axelson, M. Gibaldi, and W. D. Conway, *Xenobiotica*, **4**, 585 (1974); R. E. Counsell, T. Yu, V. V. Ranade, A. A. Buswink, E. A. Carr, Jr., and M. Carroll, *J. Nucl. Med.*, **15**, 991 (1974).
- (15) R. E. Counsell and R. D. Ice in "Drug Design", Vol. 6, E. J. Ariens, Ed., Academic Press, New York, N.Y., 1975, p 172.
- (16) N. Lofgren and B. J. Lundquist, *Chem. Abstr.*, **42**, P6378 (1948); U.S. Patent 2 441 498 42 (1948).
- (17) N. Lofgren, "Studies on Local Anesthetics: Xylocaine, A New Synthetic Drug", Hoeggstroms, Stockholm, 1948, p 28.