may open *in vivo* so that inactivity cannot be assigned to the aziridine *per se*, although the present compounds were unchanged by a 5-hr incubation at  $37^{\circ}$  in aqueous solution buffered at pH 7.3. The p $K_a$  of Ia (7.2) is little different from that of aziridine itself (8.0) but is somewhat less than that of the more basic amphetamine (9.90<sup>12</sup>). It is possible that this leads to inefficient binding of the title compounds to a receptor because of the reduced availability of the nitrogen lone pair, a condition previously proposed for the reduced activity of the even less basic 2-amino-3-phenyl-1,1.1-trifluoropropanes<sup>4</sup> and 1-cyanophenethylamines.<sup>5</sup>

## **Experimental Section†**

Phenylalaninyl Hydrogen Sulfates. A. To a cold suspension of 4-chlorophenylalaninol (18.3 g, 0.1 mole) in H<sub>2</sub>O (30 ml) was added cold concd H<sub>2</sub>SO<sub>4</sub> (10 g, 0.1 mole). The light yellow solution was heated at 120° to remove H<sub>2</sub>O, the final traces being removed on the rotary evaporator. Recrystallization of the brown residue from 40% aqueous EtOH, with concentration of the mother liquors, gave 19 g (67%) of yellow needles, mp 277-279°. Anal. (C<sub>9</sub>H<sub>12</sub>CINO<sub>4</sub> S: H<sub>2</sub>O) C, H, N.

Phenylalaninyl hydrogen sulfate, obtained in 70% yield by the same procedure, had mp 253-255° (lit.<sup>6</sup> 265-270°). Anal.  $(C_9H_{13}NO_4S \cdot H_2O)$  C, H, N.

**B.** Dicyclohexylcarbodiimide (24.7 g, 0.12 mole) in DMF (50 ml) was added at 0° to a solution of 4-methoxyphenylalaninol (6.2 g, 0.033 mole) in DMF (60 ml). Concd  $H_2SO_4$  (6 g, 0.033 mole) in DMF (25 ml) was then added dropwise over 30 min at 0°. The mixture was stirred for 90 min at room temperature, the solid dicyclohexylurea was filtered off, and the filtrate was evaporated to dryness. The residue was washed well with cold water and recrystallized from aqueous EtOH, mp 255-256°, yield 4.4 g (55%). Anal. (C<sub>10</sub>H<sub>15</sub>NO<sub>5</sub>S) C, H, N.

2-Benzylaziridine Hydrogen Maleates. A. 2-Benzylaziridine (Ia), bp 68-70° (0.5 mm), was prepared by continuous distillation from a mixture of aqueous NaOH and phenylalaninyl hydrogen sulfate.<sup>6</sup> 2-Benzylaziridine hydrogen maleate had mp 93-94° (EtOH-Et<sub>2</sub>O), yield 56%. Anal. ( $C_9H_{11}N \cdot C_4H_4O_4$ ) C, H, N.

**B.** Typically, 6.5 g of 4-chlorophenylalaninyl hydrogen sulfate and 50 ml of 35% aqueous NaOH were refluxed together for 2 hr. The mixture was cooled, extracted with Et<sub>2</sub>O, dried (MgSO<sub>4</sub>), and distilled as a pale yellow oil, bp 97-98° (0.5 mm). Addition of a saturated ethereal solution of maleic acid gave the hydrogen maleate (**Ib**), which was recrystallized from EtOH-Et<sub>2</sub>O as colorless plates, mp 94-95°, yield 3.0 g (42%). Anal. (C<sub>9</sub>H<sub>10</sub>ClN·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·H<sub>2</sub>O) C, H, N. 2-(4-Methoxybenzyl)aziridine hydrogen maleate (Ic) had mp 96-97° (EtOH-Et<sub>2</sub>O), yield 36%. Anal. (C<sub>10</sub>H<sub>13</sub>N·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>) C, H, N.

Acknowledgments. We thank Messrs. D. A. Buxton, D. M. Green, and A. Muir for the pharmacological data and Mr. D. B. Coult for  $pK_a$  determinations.

### References

- (1) O. C. Dermer and G. E. Ham, "Ethylenimine and Other Aziridines," Academic Press, London, 1969.
- (2) N. B. Chapman and J. D. P. Graham, Drugs Affecting Peripheral Nerv. Syst., 1, 473 (1967).
- (3) R. M. Pinder and A. Burger, J. Pharm. Sci., 56, 970 (1967).
- (4) R. M. Pinder, R. W. Brimblecombe, and D. M. Green, J. Med.
- Chem., 12, 322 (1969). (5) R. M. Pinder, A. Burger, and E. J. Ariens, Arzneim.-Forsch., 20, 245 (1970).
- (6) D. V. Kashelikar and P. E. Fanta, J. Amer. Chem. Soc., 82, 4930 (1960).
- (7) R. O. Mumma, Lipids, 1, 221 (1966).
- (8) K. Okada, K. Ichimura, and R. Sudo, Bull. Chem. Soc., Jap., 43, 1185 (1970).
- (9) R. W. Brimblecombe, Int. J. Neuropharmacol., 6, 423 (1967).
- (10) R. W. Brimblecombe, Psychopharmacologia, 4, 139 (1963).

(11) A. Burger, Fortschr. Arzneimittelforsch., 15, 227 (1971).

(12) T. B. Vree and J. M. Van Rossum, J. Pharm. Pharmacol., 21,

774 (1969).

# Central Nervous System Depressants. 10. 1-Carbamoylbenzodiazepines<sup>†,1</sup>

Robert Bruce Moffett\* and Allan D. Rudzik

Research Laboratories of The Upjohn Company, Kalamazoo, Michigan 49001. Received March 10, 1972

A considerable number of diazepam analogs have been reported in which other groups were substituted for the 1-methyl. In general, any such substituent containing more than 3 carbon atoms has been less active in tests thought to correlate with antianxiety activity.<sup>2</sup> In this work a series of 1-carbamoyl derivatives (I) is reported. These compounds



1, X = Cl; X' = H; R =  $CH_2CH=CH_2$ 

were prepared by treatment of 1-unsubstituted benzodiazepin-2-one with the appropriate isocyanate. This procedure is exemplified in the Experimental Section by the preparation of 1, and the compounds are listed in Table I.

After the start of this work it came to our attention that Usui, *et al.*, <sup>3</sup> had prepared a similar series of benzodiazepine derivatives. However, our series overlapped theirs in only one compound, the methylcarbamoyl analog (10). Even though this was one of their more active compounds, our compound 1 proved to be more active in mice on all the parameters used (Table II).

Pharmacology. The pharmacologic results obtained with this series of 1-carbamoylbenzodiazepines are presented in Table II, and the results are compared to those obtained with diazepam in the same test systems. Compound 1, the allylcarbamoyl analog, was the most active compound in this series. It was equipotent or more active than diazepam on all end points except the antagonism of pentylenetetrazolinduced clonic convulsions and the potentiation of ethanol narcosis. The activity of the allyl derivative (1) in the traction and strychnine tests may indicate potent muscle relaxant activity.

Substitution of a cyano group for a chloro group in position 7 (2) markedly decreased the pharmacologic activity. Compound 2 was inactive in antagonizing strychnine lethality and pentylenetetrazol-induced clonic convulsions and on all other end points it was weakly active. Substitution of a cyano group (RO5-4528) for the chloro group in the diazepam series produced a compound more potent on almost all test systems except the simple reflex tests (chimney, dish, and pedestal) and antagonism of pentylenetetrazolinduced seizures. It appears therefore that the structureactivity relationships for the 1-carbamoylbenzodiazepines may be different from those established for the 1-methyl derivatives.

<sup>†</sup>Melting points are uncorrected and were measured in an Electrothermal capillary apparatus. Satisfactory analyses (within  $\pm 0.4\%$  of the theoretical values) were obtained for all compounds, which were identified by ir and nmr spectroscopy.  $pK_a$  values were determined potentiometrically with a Radiometer Titrograph SBR 2c.

<sup>†</sup>Presented in part at the Sixth Great Lakes Regional Meeting of the American Chemical Society, Houghton, Mich., June 22, 1972.



No.	х	Χ'	R	Cryst solvent	Yield, <sup>a</sup> %	Mp, <sup>b</sup> °C	and analyses <sup>c</sup>
1	Cl	Н	CH <sub>2</sub> CH=CH	<i>i</i> -PrOH	88.6	103-105	C <sub>10</sub> H <sub>14</sub> ClN <sub>2</sub> O <sub>2</sub>
2	C≡N	Н	$CH_{2}CH=CH_{2}^{2}$	<i>i</i> -PrOH	75	137.5-139	C, H, NO
3	C1	Н	$CH_{2}CH=CHCH_{3}^{d}$	$C_{A}H_{1}$	23	114-116	C,H,CINO,
4	C1	Н	CHCH <sub>2</sub> CH <sub>2</sub> <sup>e</sup>	CH,Ċl,-C,H,,	32	213.5-214.5 <sup>e</sup>	C, H, CIN,O
5	C1	Н	CH(CH <sub>2</sub> ) <sub>4</sub> CH <sub>2</sub>	i-PrOH	46	134.5-136	C,,H,,CIN,O,
6	Cl	Н	CH <sub>2</sub> CH <sub>2</sub> Cl	i <b>-P</b> rOH	80	110-113	$C_1 H_1 C_1 N_2 O_2$
7	Cl	Н	CH <sub>2</sub> COOEt	<i>i</i> -PrOH	79	125-127.5	C,H,CIN,O
8	Cl	OCOCH,	CH <sub>3</sub>	EtCOMe	81.5	170-171	C, H, CIN,O
9	Cl	OCONHCH,	CH	<i>i</i> -PrOH	70	174-175	C, H, CIN O
10	Cl	H	CH <sup>f</sup>	Et <sub>2</sub> O	90	147.5–148.5 <sup><i>f</i></sup>	$C_{17}H_{14}CIN_{3}O_{2}$

<sup>a</sup>Yields are calculated on material melting not less than 2° below the highest mp obtained. <sup>b</sup>See footnote  $\ddagger$ . <sup>c</sup>All these benzodiazepines were analyzed for C, H, N, and Cl (except 2). See footnote  $\ddagger$ . <sup>d</sup>The crude product was purified by column chromatograph on SiO<sub>2</sub> and eluted with 5% MeOH in PhH. The appropriate fractions were combined, evaporated, and repeatedly fractionally crystallized from cyclohexane. <sup>e</sup>The crude product was purified by column chromatography on SiO<sub>2</sub> and eluted with 1% MeOH in CHCl<sub>3</sub>. The appropriate fractions were combined, evaporated, and crystallized first from cyclohexane, then from *i*-PrOH, and finally from a mixture of CH<sub>2</sub>Cl<sub>1</sub> and cyclohexane. The product exhibited a double mp, first melting at 144-145°, resolidifying at 146-148° and then melting at 213.5-214.5°. <sup>f</sup>Usui, et al., <sup>3</sup> report mp 150°.

#### Table II. Pharmacology<sup>a</sup>

No.	Tr₅₀ <sup>b</sup>	$\operatorname{Ch}_{\mathfrak{so}}^{b}$	D₅0 <sup>b</sup>	$P_{50}^{b}$	Nicotine (TE) antag <sup>b</sup>	Thiosemicarbazide antag <sup>c</sup>	Stry chnine antag <sup>c</sup>	Electroshock antag <sup>d</sup>	Pentylenetetrazol antag <sup>d</sup>	EtOH potentiation <sup>e</sup>
1	2.8	0.8	0.8	0.8	0.2	1.1	9.0	23	1.6	2.0
2	50	16	6.3	9.0	4.5	4.0	>50		>25	12
3	10	3.1	2.0	4.0	1.6	2.0	13	>100	3.2	
4	20	7.0	2.8	7.0	2.8	3.1	40	200	18	20
5	>100	71	56	89	45	>25	>50		>25	>100
6	16	>6.3	3.5	4.0	0.3	2.5	11	>50	2.8	2.3
7	45	6.0	9.0	10	3.0	2.8	>50	>50	10	7.0
8	112	36	13	15	4.0	25	>50	>50	28	>100
9	>200	45	45	71	126					
10 <sup>f</sup>	11	4.5	3.1	>6.3	1.1	2.0	32	>50	2.2	2.8
Diazepam <sup>g</sup>	7	2.0	0.7	1.3	0.28	0.7	8.0	50	0.8	0.9
RO5-4528 <sup>g, h</sup>	3.6	2.0	0.8	1.1	0.06	0.15	5.6	28	1.6	0.4

<sup>a</sup>Carworth Farms male, albino mice (CF-1), weighing 18-22 g, were used for all studies reported here. The test compounds were dissolved or suspended in 0.25% aqueous methylcellulose soln and administered ip. Values are  $ED_{50}$ 's expressed in mg/kg. <sup>b</sup>Procedures for measuring the effects of the compound on overt behavior, traction ( $Tr_{50}$ ), chimney ( $Ch_{50}$ ), dish ( $D_{50}$ ), pedestal ( $P_{50}$ ), and antagonism of nicotine-induced tonic extensor convulsions (TE) have been described previously.<sup>6</sup> <sup>C</sup>Procedures for the antagonism of thiosemicarbazide (TSC), electroshock convulsions, and strychnine lethality have been described.<sup>7</sup> <sup>d</sup>Procedures for measuring antagonism of pentylenetetrazole-induced convulsions have been described.<sup>8</sup> <sup>e</sup>Procedures for measuring potentiation of ethanol narcosis have been described.<sup>9</sup> <sup>f</sup>See ref 3. <sup>g</sup>The authors thank Hoffman-LaRoche Inc. for a sample. <sup>h</sup>2,3-Dihydro-1-methyl-2-0xo-5-phenyl-1*H*-1,4-benzodiazepine-7-carbonitrile.

Extending the length of the side chain from allyl (1) to crotyl (3) decreased the activity of the compound in all test systems. Substitution of cyclopropyl (4) or cyclohexyl (5) in place of the allyl group also markedly decreased the activity.

Two other compunds in this series had potent central nervous system depressant activity. These were the compounds where R was either  $CH_2CH_2Cl$  (6) or  $CH_3$  (10). However, both of these compounds were less active than the allyl derivatives (1). Compound 6, although much weaker than the allyl derivative (1) on simple reflexes and strychnine end points, approached the allyl derivative in potency on anticonvulsant end points (nicotine, thiosemicarbazide, and pentylenetetrazol antagonism) and also in its ability to potentiate ethanol narcosis. Thus, compound 6 may be an active anticonvulsant with weak effects as a muscle relaxant or CNS depressant. Compound 10 was also much more active on anticonvulsant end points than on those measuring CNS depression.

It appears from this small series of compounds that 1-carbamoylbenzodiazepines possess potent CNS depressant and anticonvulsant activity. They may have a different structure-activity relationship pattern than do 1-methyl-substituted benzodiazepines. The substantiation of this hypothesis requires the testing of more 7-substituted derivatives and also the effect of substitution on the 5-phenyl ring of the molecule.

#### Experimental Section<sup>‡</sup>

N-Allyl-7-chloro-2,3-dihydro-2-oxo-5-phenyl-1*H*-1,4-benzodiazepine-1-carboxamide (1). A mixture of 16.26 g (0.06 mole) of

<sup>&</sup>lt;sup>‡</sup>Melting points were taken in capillary tubes with a partial immersion thermometer. Calibration of the apparatus against standard compounds showed no need for correction. Ir (Nujol mull), usually nmr (in DMSO- $d_6$  or CDCl<sub>3</sub>), and often mass spectra were obtained on pure compounds and were in accordance with the proposed structure. Where analyses were indicated only by symbols of the elements of functions, analytical results obtained for these elements or functions were within ±0.4% of theoretical.

7-chloro-1,3-dihydro-5-phenyl-2H-1,4-benzodiazepin-2-one, 125 ml of THF, and 45 ml of allyl isocyanate (added 25 ml at start and 10 ml after 1 and after 3 days) was stirred under  $N_2$  and under reflux for 5 days. The reaction was followed by tlc (on SiO<sub>2</sub>, with 5% MeOH in CHCl<sub>2</sub>). The mixture was filtered and evaporated *in vacuo*. PhCH<sub>3</sub> was added and evaporated to remove excess isocyanate, and the resulting gum was crystallized from 250 ml of *i*-PrOH. Thus 18.8 g (88.6%) of white fluffy needles was obtained, mp 103-105°.

Crotyl Isocyanate. To a suspension of 41.54 g (0.256 mole) of 1,1'-carbonyldiimidazole in 100 ml of CH<sub>4</sub>Cl<sub>4</sub> was slowly added, with stirring during 0.5 hr at 25°, 18.0 g (0.256 mole) of crotylamine. After standing overnight, the solution was slowly distilled through an 18-cm, glass-helices-packed, column. After removing the CH<sub>4</sub>Cl<sub>2</sub>, a fraction boiling at 45-46° (56 mm) was collected. This was redistilled at atmospheric pressure yielding 9.7 g (39.9%) of colorless liquid, bp 113°. Ir and nmr support the structure, and gc indicates it was greater than 94.7% pure. Christophersen and Holm<sup>4</sup> report bp 116°.

Cyclopropyl Isocyanate. A suspension of 35.8 g (0.55 mole) of NaN<sub>3</sub> in 200 ml of triethylene glycol dimethyl ether in a 1-l. flask fitted with a stirrer, thermometer, dropping funnel, and a Vigreux column to which was attached a Dry Ice cooled, two-necked flask was cooled to 5°, and 52.25 g (0.5 mole) of cyclopropylcarbonyl chloride was slowly added under N<sub>4</sub> during 10 min. After stirring at 0-25° for 1 hr, the mixture was slowly heated in a water bath. At about 55° N<sub>2</sub> started to come off and it was evolved rapidly at 70-103°. The flask was then heated in an oil bath up to 171°, and the product was distilled at 56 mm. Solvent refluxed in the bottom of the column. The distillate was redistilled through an 18-cm, glass-helices-packed column giving 25.67 g (62%) of colorless liquid, bp 87° (atm),  $n^{26}$ D 1.4210;  $d_4^{25}$  1.00. Jones and Scott<sup>5</sup> apparently prepared this but did not isolate or characterize it.

Acknowledgments. The authors wish to thank the following people who contributed to this work: our Physical and Analytical Chemistry Unit for analytical and spectral data; Mr. Bharat V. Kamdar and Mr. Walter Friis for technical assistance.

# References

- (1) R. B. Moffett and A. D. Rudzik, J. Med. Chem., 14, 588 (1971). (paper 9).
- (2) L. H. Sternback, L. O. Randall, R. Banziger, and H. Lehr, "Drugs Affecting the Central Nervous System," Vol. 2, A. Burger, Ed., Marcel Dekker, New York, N.Y., 1968, Chapter 6, p 252.
- (3) Y. Usui, Y. Hara, I. Mikami, and T. Asuda, Takeda Kenkyusho Nempo, 30, 145 (1971).
- (4) C. Christophersen and A. Holm, Acta Chem. Scand., 24, 1852 (1970).
- (5) L. W. Jones and A. W. Scott, J. Amer. Chem. Soc., 44, 407 (1922).
- (6) G. H. Youngdale, D. G. Anger, W. C. Anthony, J. P. Da Vanzo, M. E. Greig, R. V. Heinzelman, H. H. Keasling, and J. Szmuszkovicz, J. Med. Chem., 7, 415 (1964).
- (7) H. H. Keasling, E. L. Schumann, and W. Veldkamp, *ibid.*, 8, 548 (1965).
- (8) J. B. Hester, A. D. Rudzik, and B. V. Kamdar, *ibid.*, 14, 1078 (1971).
- (9) J. B. Hester, A. D. Rudzik, H. H. Keasling, and W. Veldkamp, *ibid.*, 13, 23 (1970).

# Derivatives of Indole-1-acetic Acid as Antiinflammatory Agents

Karl J. Doebel\* and Jan W. F. Wasley

Research Department, Pharmaceuticals Division, Ciba-Geigy Corporation, Ardsley, New York 10502. Received January 7, 1972

In view of the activity of 1-(p-chlorobenzoyl)-5-methoxy-2-methylindole-3-acetic acid (1, Indomethacin) as an antiinflammatory agent, a study of molecular models showed 2 to be spatially similar to Indomethacin (1). The related isomer 3 was sufficiently similar to merit investigation also.



The publication of  $2^1$  prompts us to report our findings in this area.

## **Results and Discussion**

Preparation of 2 and 3 proceeded by similar routes, but for 2 the indole 4a was required. This was prepared by catalytic hydrogenation of 5.



The general synthetic pathway involved the condensation of p-chlorobenzonitrile with 4a or 4b to yield the imines 6a and 6b, respectively. It was anticipated that the imines would readily hydrolyze to the corresponding ketones 7a and 7b, but this was not the case.



Alkylation of 6a or 6b with ethyl bromoacetate in acetone in the presence of anhydrous  $K_2CO_3$  proceeded smoothly, and alkaline hydrolysis (3 N NaOH) of the intermediate esters gave the desired products 2 and 3.

A comparison of the  $pK_{mcs}$  and distribution coefficients indicated that 2 and 3 are significantly more acidic than Indomethacin (1), as can be seen from Table I.

In order to obtain molecules with acidity similar to indomethacin (1), the corresponding 1-propionic and 1-isobutyric acid derivatives 8 and 9 were prepared by the con-



densation of **6b** with methyl acrylate and methyl methacrylate, respectively, followed by alkaline hydrolysis of the intermediate esters. As can be seen from Table I, the acidity of these compounds now approximates that of 1.

Table I					
Compound	pK <sub>mcs</sub>	P (CHCl <sub>3</sub> -aqueous buffer, pH 6.9)			
1	6.45	37.2			
2	4.97	0.13			
3	5.00	0.13			
8	6.08	11.9			
9	6.26	60.0			