

Total Synthesis of (+)-Polyoxin J

Alessandro Dondoni,^a Federico Junquera,^b Francisco L. Merchán,^b Pedro Merino^{a,b} and Tomas Tejero^b^a Dipartimento di Chimica, Laboratorio di Chimica Organica, Università, Ferrara, Italy^b Departamento de Química Organica, ICMA, Universidad de Zaragoza, Zaragoza, Spain


The antifungal antibiotic, polyoxin J **1**, has been obtained (46.4%) by the coupling of a protected derivative of 5-*O*-carbamoylpolyoxamic acid **2** with thymine polyoxin C **3** which were prepared by a stereoselective aminohomologation strategy of sugar aldehydes (L-threose and dialdo-D-ribofuranose) employing nitrones **4** and **5** as their iminium derivatives and furan as a carboxylate group equivalent.

Polyoxin J **1** is one of the components of a wide class of peptidyl pyrimidine nucleoside antibiotics¹ which are attracting increasing interest as antifungal compounds² because of their ability to inhibit fungal cell wall chitin biosynthesis.³ Two total syntheses of **1** have been reported with a twenty year gap between them,^{4,5} both methods involving coupling of the key components of the molecule, 5-*O*-carbamoylpolyoxamic acid **2** and thymine polyoxin C **3** (Scheme 1). The efficient synthesis of protected derivatives of these compounds is *per se* an important issue which although addressed in various instances over the years,⁶ still appears worth investigation. With the total synthesis of **1** and the development of a general synthetic method of the various components of the polyoxin complex as our goals, we studied new syntheses of **2** and **3** employing nitrones **4** and **5** derived from sugar aldehydes. A strategy for aldehyde homologation to the α -amino aldehyde and acid *via* addition of 2-thiazolyl-7 (masked formyl) and 2-furyllithium⁸ (masked carboxyl) to nitrones has been recently developed in our laboratories and its successful implementation to the synthesis of **2** and **3** has been described.^{9,10} Here we report an improved reaction sequence leading to the nucleoside **3** and the coupling with an activated derivative of **2** to complete the total synthesis of polyoxin J **1**.


The diastereoselective addition (d.s. 82% of correct isomer) of 2-furyllithium to the ribofuranosyl nitrone **5** precomplexed with Et₂AlCl to give the hydroxylamine **6** (72%) has been previously described¹⁰ (Scheme 2). Given the low yields of the reactions leading to **3**, particularly the anomeric acetolysis because of the competitive endocyclic mode of glycosyl cleavage,¹¹ compound **6** was elaborated as follows. The *N*-hydroxy group was removed *N*-benzylamino group was pro-

ected as the trifluoroacetamido derivative by treatment with trifluoroacetic anhydride in the presence of pyridine. In this way, amide **7** was obtained in 56% yield.[†] After deisopropylidenation of **7** by acid hydrolysis, treatment of the resulting furanose with acetic anhydride in the presence of pyridine gave the triacetate **8** (78%) as a mixture of α - and β -anomers (24:76). The carboxylic group was liberated from the furan ring of **8** by treatment with RuCl₃–NaIO₄. Without isolation, the carboxylic acid was converted by diazomethane into the methyl ribofuranosyl glycinate derivative **9** in 60% overall yield. This product was subjected to Vorbrüggen condensation¹² with 5-methyl-2,4-bis(trimethylsilyloxy)pyrimidine to exclusively provide the β -nucleoside **10** (70%). *N*-Debenzylation of **10** by hydrogenolysis over Pd(OH)₂ gave **11** from which all the remaining protective groups were removed by treatment with lithium hydroxide in THF. The thymine polyoxin C **3** was thus isolated in 67% yield and 10.6% overall yield from the nitrone **5**. Synthetic **3** prepared by the above and earlier reaction sequence¹⁰ showed identical characteristics which were in agreement with literature values.[‡]

Successful coupling of **3** with a protected derivative of **2** was performed by the *N,N*-dicyclohexylcarbodiimide–*N*-hydroxy-succinimide (DCC–HOSu) active ester method⁴ in Me₂SO and



Scheme 2 Reagents and conditions: i, 2-Lithiofuran, Et₂AlCl, THF–Et₂O (1:1), –80 °C, 1 h; ii, Zn, Cu(AcO)₂, AcOH, 70 °C, 2 h then (CF₃CO)₂O, pyridine–CH₂Cl₂ (1:1), room temp., 3 h; iii, AcOH–HCl–H₂O (80:1:19), 70 °C, 4 h then Ac₂O, DMAP, pyridine, room temp., 2 h; iv, RuCl₃, NaIO₄, MeCN–CCl₄–H₂O (2:2:3), room temp., 15 min. then CH₂N₂, Et₂O, 0 °C, 1 h; v, 5-methyl-2,4-bis(trimethylsilyloxy)pyrimidine, Me₃SiOSO₂CF₃, CH₂Cl₂, reflux, 2 h; vi, H₂, Pd(OH)₂–C 20%, MeOH, room temp., 6 h; vii, LiOH, THF–H₂O (6:1), 0 °C, 1 h



Scheme 3 Reagents and conditions: i, *N*-hydroxysuccinimide, DCC, EtOAc, 0 °C, room temp., 8 h; ii, thymine polyoxin C 3, Pr_2EtN , Me_2SO , room temp., 24 h; iii, TFA–MeOH– H_2O (1 : 1 : 2), 0 °C, 2 h

N,N-diisopropylethylamine as base¹³ (Scheme 3). Thus, the polyoxamic acid derivative **12** obtained as described earlier (21.7% from a protected derivative of L-threose),⁹ by treatment with DCC–HOSu was converted into the active ester **13** which was then condensed with **3** to give the dipeptide **14** (58%).[‡] Removal of the *N*-Boc and *O*-isopropylidene protecting groups from **14** upon acid hydrolysis afforded polyoxin J **1** in 80% yield (46.4% overall from **3** and **12**). The physical properties of compound **1** were in good agreement with the literature values of both synthetic and natural polyoxin J.[‡] The efficient coupling of unprotected **3** with the relatively hindered α -amino acid **12** is noteworthy since a similar reaction employed for the assemblage of the components of nikkomycin Z gave very poor results.¹⁴

This new synthesis of polyoxin J **1** from protected derivatives of L-threose and dialdo-D-ribofuranose[§] provides an illustration of the synthetic utility of the aminohomologation strategy and indicates a viable route for the synthesis of other components of the polyoxin¹ and nikkomycin¹⁴ families.

Acknowledgement is made to the Consiglio Nazionale delle Ricerche (Roma) and the Ministerio de Educacion y Ciencia (DGICYT, Madrid, Project PB94-0598) for support of this research. We also thank Dr X. Huguet (CID-CSIC, Barcelona) for mass spectral determinations.

Received, 25th July 1995; Com. 5/04932F

Footnotes

[†] All new compounds as well as the synthetic polyoxin J **1** and thymine polyoxin C **3** exhibited consistent ¹H and ¹³C NMR spectra.

[‡] Physical and spectroscopic data for compound **3**: mp 170 (soft), 192 °C, $[\alpha]_D^{25} +8.8$ (c 0.09, H_2O), HRMS (FAB, H_2O –glycerol): Calc. for $\text{C}_{11}\text{H}_{15}\text{N}_3\text{O}_7$ ($\text{M} + \text{H}^+$): 302.0988, Found: 302.0987. Lit.,¹¹ mp 182–185 and 190–194 °C, $[\alpha]_D^{25} +8.0$ (c 0.37, H_2O). For compound **14**: mp 110° (soft), 172 °C, $[\alpha]_D^{25} -17.2$ (c 0.34, MeOH), LRMS (FAB, MeOH–glycerol): 632 ($\text{M} + \text{H}^+$). For compound **1**: mp 200 °C (dec), $[\alpha]_D^{25} +30.3$ (c 0.10, H_2O), LRMS (FAB, MeOH–NBA): 492 ($\text{M} + \text{H}^+$). For synthetic **1**: Lit.,⁵ mp 200–210 °C (dec), $[\alpha]_D^{25} +35.0$ (c 0.8, H_2O), lit.,⁴ $[\alpha]_D^{25} +33$ (c 0.75, H_2O); for authentic **1**: lit.,⁵ mp 198–208 °C (dec), lit.,⁴ $[\alpha]_D^{25} +31$ (solvent not given).

[§] 4-*O*-Benzyl-2,3-*O*-isopropylidene-L-threose (T. Mukaiyama, K. Suzuki, T. Yamada and F. Tabusa, *Tetrahedron*, 1990, **46**, 265) is the starting material for the synthesis of **12** while methyl 2,3-*O*-isopropylidene- β -D-ribofuranose (A. M. Sepulchre, G. Vass and S. D. Gero, *Tetrahedron Lett.*, 1973, 3619) is used for the synthesis of **3**.

References

- 1 K. Isono, K. Asahi and S. Suzuki, *J. Am. Chem. Soc.*, 1969, **91**, 7490.
- 2 K. Isono and S. Suzuki, *Heterocycles*, 1979, **13**, 333; P. Shenbagamurthi, H. A. Smith, J. M. Becker, A. S. Steinfeld and F. Naider, *J. Med. Chem.*, 1983, **26**, 1518; G. Emmer, N. S. Ryder and M. A. Grassberger, *J. Med. Chem.*, 1985, **28**, 278; A. B. Cooper, J. Desai, R. G. Lovey, A. K. Saksena, V. M. Girijavallabhan, A. K. Ganguly, D. Loebenberg, R. Parmegiani and A. Cacciapuoti, *Bioorg. Med. Chem. Lett.*, 1993, **3**, 1079.
- 3 M. Mori, K. Kakiki and T. Misato, *Agric. Biol. Chem.*, 1974, **38**, 691.
- 4 H. Kuzuhara, H. Ohri and S. Emoto, *Tetrahedron Lett.*, 1973, 5055.
- 5 N. Chida, K. Koizumi, Y. Kitada, C. Yokoyama and S. Ogawa, *J. Chem. Soc., Chem. Commun.*, 1994, 111.
- 6 For recent syntheses of polyoxamic acid derivatives and the nucleoside portion of polyoxins, see ref. 5 and P. Garner, in *Studies in Natural Products Chemistry*, vol. 1, *Stereoselective Synthesis, Part A*, ed. Atta-Ur Rahman, Elsevier, Amsterdam, 1988, p. 397.
- 7 A. Dondoni, S. Franco, F. Junquera, F. L. Merchán, P. Merino, T. Tejero and V. Bertolasi, *Chemistry, A European Journal*, 1995, in the press.
- 8 A. Dondoni, F. Junquera, F. L. Merchán, P. Merino and T. Tejero, *Synthesis*, 1994, 1450.
- 9 A. Dondoni, S. Franco, F. L. Merchán, P. Merino and T. Tejero, *Tetrahedron Lett.*, 1993, **34**, 5479.
- 10 A. Dondoni, F. Junquera, F. L. Merchán, P. Merino and T. Tejero, *Tetrahedron Lett.*, 1994, **35**, 9439.
- 11 P. Garner and J. M. Park, *J. Org. Chem.*, 1990, **55**, 3772.
- 12 H. Vorbrüggen, K. Krolikiewicz and B. Bennua, *Chem. Ber.*, 1981, **114**, 1234.
- 13 E. Krainer, J. M. Becker and F. Naider, *J. Med. Chem.*, 1991, **34**, 174.
- 14 A. K. Saksena, R. G. Lovey, V. M. Girijavallabhan, H. Guzik and A. K. Ganguly, *Tetrahedron Lett.*, 1993, **34**, 3267.