The First Alkyl(silyl)palladium Complexes: Formation by Oxidative Addition of Silacyclobutanes to Palladium **Complexes, Reductive Elimination, and Other Reactivities Relevant to Catalysis**

Yoshifumi Tanaka,[†] Hiroshi Yamashita,[‡] Shigeru Shimada,[‡] and Masato Tanaka^{*,†,‡}

Department of Chemistry, University of Tsukuba, Tsukuba, Ibaraki 305, Japan, and National Institute of Materials and Chemical Research, Tsukuba, Ibaraki 305, Japan

Received March 6, 1997[®]

Summary: 1,1-Diphenyl- and 1,1-dimethylsilacyclobutanes reacted with $Me_2Pd(dmpe)$ (dmpe = 1,2-bis(dimethylphosphino)ethane) or Pd(PhCH=CH₂)(dmpe) to give 2,2-diphenyl- and 2,2-dimethyl-1-pallada-2-silacyclopentane complexes, the diphenyl complex being characterized by X-ray analysis. Treatment of the diphenyl complex with acetylenes or 1,2-disilacyclopentane induced reductive elimination to regenerate the parent 1,1diphenylsilacyclobutane. A dihydrosilane or a silacyclobutane reacted with the diphenyl complex to afford a 1,3-bis(hydrosilyl)propane or a 1,5-disilacyclooctane, respectively.

Although silicon-carbon bonds usually are unreactive toward transition-metal complexes, those of silacyclobutanes (1) are exceptions. They are reactive due to their ring strain.¹ Quite a few reactions of **1** that are catalyzed by transition-metal complexes have been reported, such as ring-opening polymerization,² dimerization,³ cross-dimerization with disilanes,⁴ and cycloaddition reactions⁵ with acetylenes and allenes. 1-Metalla-2silacylopentanes are believed to be involved as intermediates in these catalytic reactions. Indeed, 1-ferra-⁶ and 1-platina-2-silacyclopentane³ complexes have been isolated. In a broader view, the chemistry of 1-metalla-2-silacyclopentanes is important in its own right; these complexes provide a rare opportunity to study the reactivities of alkyl(silyl)metal species, a very important class of catalytic intermediates, which are not always sufficiently stable to allow their manipulation. This paper reports the chemistry of 1-pallada-2-silacyclopentane complexes, which are the first alkyl(silyl)palladium species.

In one example Me₂Pd(dmpe) (2a, 0.09 mmol; dmpe = 1,2-bis(dimethylphosphino)ethane) and 1,1-diphenylsilacyclobutane (1a, 0.27 mmol) were mixed in C_6D_6 (0.3 mL), and the solution was heated in a sealed NMR tube at 60 °C for 7 h. Monitoring of the reaction by ¹H, ¹³C, ²⁹Si, and ³¹P NMR spectroscopy revealed the formation of a 1-pallada-2-silacyclopentane complex (3a, 90% NMR yield)^{$\overline{7}$} and allylmethyldiphenylsilane (**4a**, 80% NMR yield) (eq 1). The ¹H NMR spectrum displayed a very



weak signal at δ 0.23, suggesting that methane had been generated. The structure of 4a was confirmed by GC-MS of the reaction mixture. The mixture was evaporated in vacuo, and the residue was recrystallized from toluene-hexane to give **3a** as pale yellow crystals (32.5 mg, 75%). The structure of 3a was unambiguously confirmed by X-ray diffraction, thus verifying that oxidative addition of the Si-C bond in the ring system had taken place. As the ORTEP⁸ drawing (Figure 1) shows,⁹ 3a is a square-planar complex, the deviation from planarity being very small.¹⁰ The Pd-Si bond distance (2.341(2) Å) is within the range of those reported in the literature (2.33-2.43 Å).¹¹ The Pd-P bond trans to the silicon (2.351(2) Å) is significantly longer than the other Pd-P bond trans to the carbon (2.291(2) Å). This is due to the strong *trans* influence

[†] University of Tsukuba

[‡] National Institute of Materials and Chemical Research

[®] Abstract published in Advance ACS Abstracts, June 15, 1997.

⁽¹⁾ The strain energies of silacyclobutane, silacyclopentane, and (i) The stand energies of shacyclobulate, shacyclopetate, and stand energies of shacyclohexane are 102.5, 20.1, and 13.0 kJ mol⁻¹, respectively. See: Gordon, M. S.; Boatz, J. A.; Walsh, R. J. Phys. Chem. **1989**, 93, 1584.
(2) (a) Weyenberg, D. R.; Nelson, L. E. J. Org. Chem. **1965**, 30, 2618.
(b) Nametkin, N. S.; Ushakov, N. V.; Vdovin, V. M. Vysokomol. Soedin., Social and the form of the form of the form. Ser. A 1971, 13, 29; Chem. Abstr. 1971, 74, 88325w. (c) Cundy, C. S. Eaborn, C.; Lappert, M. F. J. Organomet. Chem. 1972, 44, 291 and references cited therein. (d) Poletaev, V. A.; Vdovin, V. M.; Nametkin, N. S. Dokl. Akad. Nauk SSSR 1973, 208, 1112; Chem. Abstr. 1973, 79, 19191r. (e) Finkel'shtein, E. Sh.; Ushakov, N. V.; Pritula, N. A.; Andreev, E. A.; Plate, N. A. Izv. Akad. Nauk SSSR, Ser. Khim. 1992, 223; Chem. Abstr. 1992, 116, 256191f. (f) Ushakov, N. V.; Yarysheva, A. Yu.; Tal'roze, R. V.; Finkel'shtein, E. Sh.; Plate, N. A. Dokl. Akad. A. 10.; Fairoze, R. V.; Finkersmein, E. Sh.; Fiate, N. A. Doki. Akad. Nauk 1992, 325, 964; Chem. Abstr. 1993, 118, 102788k. (g) Bialecka-Florjanczyk, E.; Ganicz, T.; Stanczyk, W.; Sledzinska, I. Polimery (Warsaw) 1993, 38, 424; Chem. Abstr. 1994, 121, 36351w. (h) Liao, C. X.; Chen, M. W.; Sun, L.; Weber, W. P. J. Inorg. Organomet. Polym. **1993**, *3*, 231

 ⁽³⁾ Yamashita, H.; Tanaka, M.; Honda, K. J. Am. Chem. Soc. 1995, 117, 8873.

⁽⁴⁾ Reddy, N. P.; Hayashi, T.; Tanaka, M. Chem. Commun. 1996, 1865

⁽⁵⁾ Sakurai, H.; Imai, T. Chem. Lett. 1975, 891. Takeyama, Y.; Nozaki, K.; Matsumoto, K.; Oshima, K.; Utimoto, K. Bull. Chem. Soc. Jpn. 1991, 64, 1461.

⁽⁶⁾ Cundy, C. S.; Lappert, M. F. J. Chem. Soc., Chem. Commun. 1972, 445.

⁽⁷⁾ Compounds ${\bf 3a,b}$ showed satisfactory NMR and/or analytical data (see the Supporting Information). 1 H, 29 Si, and 31 P NMR spectral data (C₆D₆) for **3a** are as follows: ¹H NMR δ 0.64 (d, J = 7.3 Hz, 6H, 1.80 (f, J = 6.6 Hz, ZH, CH₂SJ), 2.39–2.53 (m, ZH, CH₂), 2.56–2.68 (m, 2H, CH₂), 7.15–7.33 and 7.92–7.97 (each m, 3H and 2H, C₆H₅); ²⁹Si NMR δ 47.1 (dd, $J_{PSi} = 15.9$ and 174.6 Hz); ³¹P NMR δ 10.6 (d, $J_{PP} = 9.6$ Hz, $J_{PSi} = 174.6$ Hz), 14.2 (d, $J_{PP} = 9.6$ Hz). (8) Johnson, C. K. ORTEP, a FORTRAN Thermal-Ellipsoid Plot Program for Crystal Structure Illustrations; Report ORNL-3794; Oak Ridge National Laboratory: Oak Ridge, TN, 1970.



Figure 1. ORTEP drawing of **3a**. C(6) is disordered, resulting in large thermal parameters. Selected bond lengths (Å) and angles (deg): Pd(1)–Si(2), 2.341(2); Pd(1)–P(3), 2.351(2); Pd(1)–P(4), 2.291(2); Pd(1)–C(7), 2.129(7): Si(2)–Pd(1)–P(3), 174.0(1); Si(2)–Pd(1)–P(4), 100.1(1); Si(2)–Pd(1)–C(7), 80.0(2); P(3)–Pd(1)–P(4), 85.0(1); P(3)–Pd(1)–C(7), 95.2(2); P(4)–Pd(1)–C(7), 175.9(2).

of a silyl ligand as compared with an alkyl ligand. Complex **3a** was unstable to air in solution but only slightly decomposed in air in the crystalline state.

1,1-Dimethylsilacyclobutane (1b) reacted similarly with **2a** at 60 °C over 100 h to generate another fivemembered alkyl(silyl)palladium complex (**3b**)⁷ in 60% yield along with allyltrimethylsilane (**4b**, 60%). However, before total consumption of **1b**, slight decomposition of **3b** started to form Pd(dmpe)₂¹² to a small extent, indicating that **3b** is thermally less stable than **3a**. Complex **3b** could not be isolated in a pure form. The thermal instability of **3b** presumably is associated with reductive elimination to re-form **1b**.¹³

As illustrated in Scheme 1, the foregoing reactions forming compounds **3** can be best explained by a sequence comprising (i) metathesis of Me₂Pd(dmpe) with the Si–C bond of **1** to generate methyl[3-(methyldior-ganylsilyl)propyl]palladium species, (ii) β -hydride elimi-

Scheme 1



nation to result in the formation of **4**, (iii) extrusion of a methane molecule to generate Pd(0) species, and (iv) reaction of the resulting Pd(0) species with a second molecule of **1** to give **3**. Accordingly, a Pd(0) complex having a more labile ligand, such as an olefin complex, is expected to react more readily with **1** to form the fivemembered-ring complex. Indeed, as monitored by ¹H, ¹³C, and ²⁹Si NMR spectroscopy, Pd(η^{2-} PhCH=CH₂)-(dmpe) (**2b**, 0.025 mmol) reacted with **1a** or **1b** (1 equiv) in C₆D₆ (0.25 mL) even at room temperature to give, within 10 min, **3a** or **3b** in quantitative yield (eq 2). Even though the formation of **3b** in the solution was quantitative, its isolation was not successful.¹⁴



Palladium complexes are able to catalyze the reactions of silacyclobutanes with acetylenes to give silacyclohexenes and/or vinyl(allyl)silanes, and the catalysis is envisioned to proceed *via* 1-pallada-2-silacyclopentane intermediates.⁵ However, when **3a** (0.06 mmol) was allowed to react with diphenylacetylene (**5a**, 1.5 equiv) in C₆D₆ (0.3 mL) at room temperature for 30 min in a sealed NMR tube, neither the corresponding silacyclohexene nor vinyl(allyl)silane was formed. NMR spectroscopic analysis of the resulting mixture revealed that **1a** was formed in ~12% yield along with a (diphenylacetylene)palladium dmpe complex (~13%).¹⁵ Contin-

⁽⁹⁾ Crystal data for **3a**: colorless transparent prism, $0.40 \times 0.25 \times 0.15 \text{ mm}^3$; C₂₁H₃₂P₂PdSi; FW = 479.0; monoclinic, space group P₂₁/a, a = 15.939(3) Å, b = 10.083(3) Å, c = 14.834(3) Å, $\beta = 105.08(2)^\circ$, V = 2302.0(9) Å³, Z = 4; $D_{calcd} = 1.38$ g cm⁻³; μ (Mo K α) = 9.85 cm⁻¹; Mac Science MXC18 diffractometer; ambient temperature, Mo K α radiation ($\lambda = 0.710$ 73 Å); $3.0^\circ < 2\theta < 55.0^\circ$, $\omega - 2\theta$ scan; 258 parameters; 4700 observed reflections ($F > 3.0\sigma(F)$). The structure was solved by direct methods and refined by a full-matrix least-squares procedure to yield the final residuals of R = 0.044 and $R_w = 0.075$. The non-hydrogen atoms were refined anisotropically. The hydrogen atoms were then included in the calculations but not refined. All calculations were performed using the Crystan GM crystallographic software package. (10) The high thermal parameters for C(6) suggest that C(6) is

disordered.

^{(11) (}a) Pan, Y.; Mague, J. T.; Fink, M. J. Organometallics **1992**, *11*, 3495. (b) Murakami, M.; Yoshida, T.; Ito, Y. Organometallics **1994**, *13*, 2900. (c) Suginome, M.; Oike, H.; Ito, Y. Organometallics **1994**, *13*, 4148. (d) Murakami, M.; Yoshida, T.; Kawanami, S.; Ito, Y. J. Am. Chem. Soc. **1995**, *117*, 6408. (e) Suginome, M.; Oike, H.; Shuff, P. H.; Ito, Y. Organometallics **1996**, *15*, 2170. (f) Shimada, S.; Tanaka, M.; Shiro, M. Angew. Chem., Int. Ed. Engl. **1996**, *35*, 1856.

⁽¹²⁾ Broadwood-Strong, G. T. L.; Chaloner, P. A.; Hitchcock, P. B. Polyhedron 1993, 12, 721.

⁽¹³⁾ Me₂Pd(PPh₂Me)₂ also reacted with **1a** (2 equiv) in C₆D₆. at 60 °C for 1 h to give **4a** (80% based on Pd), Pd(PPh₂Me)₄, and a black precipitate (presumably metallic palladium). However, the corresponding cyclic complex was not found by NMR in the reaction mixture, suggesting its thermal instability. Me₂Pd(PMe₃)₂ behaved very similarly to give **4a** (90% based on Pd) when treated with **1a** at 60 °C for 30 min.

⁽¹⁴⁾ The deterioration of the complex during the evaporation (0.1 Torr/room temperature) of the solvent appeared to be induced by removal of **1b** extruded by spontaneous reductive elimination, suggesting the reversibility of the oxidative-addition–reductive-elimination processes under the conditions. In the resulting mixture were found Pd(dmpe)₂ by ¹H NMR and a black precipitate (presumably metallic palladium).

^{(15) &}lt;sup>1</sup>H NMR: δ 0.95–1.17 (m, 16H, PMe and PCH₂CH₂P), 6.92– 8.04 (m, 10H, C₆H₅), ³¹P NMR: δ –7.7. These NMR data are consistent with those of an authentic sample of (PhC=CPh)Pd(dmpe) generated by the reaction of Pd(η^3 -CH₂=CHCH₂)(η^5 -C₅H₅), diphenylacetylene, and dmpe in C₆D₆. See: Krause, J.; Bonrath, W.; Pörschke, K. R. Organometallics **1992**, *11*, 1158. Since the ¹H NMR signals for PMe and PCH₂CH₂P of (PhC=CPh)Pd(dmpe) were distinctly separated from those for PCH₂CH₂P of **3a** (0.66–0.95 ppm), the yield of (PhC=CPh)-Pd(dmpe) was readily evaluated by integration.

 Table 1. Reactions of 1-Pallada-1-silacyclopentane

 Complex 3a with Acetylenes 5^a

RC≡CR′ (5)	temp (°C)/time (h) b	yield of 1a (%)
PhC≡CPh (5a)	60/0.5	66 (12)
PhC≡CH (5b)	60/3	30 (4)
PhC≡CMe (5c)	60/3	16 (0)
ⁿ PrC≡C ⁿ Pr (5d)	80/12	10 (0)
	100/12	30
CH≡CCOOEt (5e)	room temp/5.5	73 (20)
MeOOCC≡CCOOMe (5f)	room temp/5.5	80 (54)

^{*a*} Complex **3a** (0.06 mmol) was treated in a sealed NMR tube with **5** (0.09 mmol) in C₆D₆ (0.3 mL), first at room temperature for 0.5 h, and then under the conditions given in the table. ^{*b*} Reaction conditions after the beginning 0.5 h. ^{*c*} ¹H NMR yield. Figure in parentheses indicate yields after the beginning 0.5 h at room temperature.

ued reaction at 60 °C for 30 min increased the yields of **1a** and the complex to 66 and 67% respectively, but the yields remained unchanged thereafter (eq 3). The reac-



tion was very clean; the starting **3a** was still found (34%) in the reaction mixture, and no other byproducts were formed in substantial amounts. The results strongly suggest that the incomplete conversion to **1a**, *i.e.*, 66% yield of **1a** at the 30 min reaction time and thereafter, is due to attainment of an equilibrium under the conditions.

Other acetylenes reacted similarly with 3a to give 1a (Table 1). Very interestingly, however, the reaction rate and the yield very much depended on the structure of the acetylenes. For instance, 4-octyne (5d) was very reluctant to induce reductive elimination; 1a was not found by ¹H NMR at all below 60 °C, and heating at 80 °C for 12 h resulted in only a 10% yield of 1a. The yield remained as low as 30% even after heating to 100 °C over 12 h. On the other hand, treatment of 3a with dimethyl acetylenedicarboxylate (5f) at room temperature for 30 min already gave 1a in 54% yield. The yield increased over 6 h to 80% and remained unchanged for an additional 42 h. These results, inclusive of those with other acetylenes shown in Table 1, clearly indicate that electron-withdrawing groups bound to the acetylenic carbon increase the reactivity in the reductive elimination both kinetically and thermodynamically. Similar observations have been reported for a cis-alkyl(silyl)bis(phosphine)platinum complex.¹⁶ A mechanism that involves dissociation of a phosphine ligand trans to the silyl group and coordination of an acetylene to the coordination site prior to the extrusion of an alkylated silane has been proposed as a major pathway. The similarity between the platinum and our palladium cases tempts us to consider essentially the same mechanism. However, taking into account the difficulty of dissociation of one of the phosphorus atoms of the dmpe ligand, it may be premature to extend further discussion along these lines.¹⁷ Five-coordinate species proposed in a similar π -acid-induced reductive elimination from a dialkyl(bipyridyl)nickel¹⁸ complex also has to be considered.

Besides the reaction with acetylenes, 1-metalla-2silacyclopentane complexes are envisioned to be involved in several catalytic reactions of **1** (*vide supra*).^{2–5} In this context, the reactivities of complex **3a** with those substrates relevant to the catalyses were briefly examined. Methylphenylsilane (**6**, 0.24 mmol) reacted with **3a** (0.06 mmol) in C_6D_6 (0.3 mL) in a sealed NMR tube even at room temperature, and the 1,3-bis(hydrosilyl)-propane compound **7** was formed in 70% yield after 24 h (eq 4). This result is in good agreement with the ring-opening reaction of silacyclobutanes with hydrosilanes.^{2a}

$$Me_{2}$$

$$Pd + PhMeSiH_{2} \rightarrow HPh_{2}Si \rightarrow SiPhMeH (4)$$

$$Me_{2} Ph_{2} \qquad 6 (4 eq) \qquad HPh_{2}Si \rightarrow SiPhMeH (4)$$

$$3a$$

The reaction of **3a** (0.06 mmol) with **1b** (0.24 mmol) in C_6D_6 (0.3 mL) in a sealed NMR tube also proceeded at 100 °C to afford a 1,5-disilacyclooctane (**8**, 44%) in 30 h (eq 5). This also provides a good model for the dimerization of silacyclobutanes.³



We have reported palladium-catalyzed selective Si-C/Si-Si cross-metathesis between a silacyclobutane and a 1,2-disilacyclopentane to form a cross-dimer.⁴ When 1,1,2,2-tetramethyl-1,2-disilacyclopentane (0.066 mmol) was added to **3a** (0.06 mmol) in C_6D_6 (0.3 mL) in a sealed NMR tube, nearly quantitative formation of a bis-(silyl)complex (**9**) was observed after 24 h at room temperature, together with **1a** (97%) arising from reductive elimination. Thus, the reaction was unable to provide a good model for the catalysis.



Acknowledgment. We are grateful to the Japan Science and Technology Corporation (JST) for partial financial support through the CREST program. Y.T. thanks Prof. Wataru Ando for helpful discussions.

Supporting Information Available: Text giving procedures to generate authentic samples of $Pd(dmpe)_2$ and $(PhC \equiv CPh)Pd(dmpe)$ and characterization data for **3a**, **3b**, **4a**, **4b**, **7**, **8**, and **9** and tables giving full details of the crystal structure analysis for **3a** (8 pages). Ordering information is given on any current masthead page.

OM970186J

⁽¹⁶⁾ Ozawa, F.; Hikida, T.; Hayashi, T. J. Am. Chem. Soc. 1994, 116, 2844.

⁽¹⁷⁾ On the basis of our observation,¹¹ reductive elimination from a T-shaped intermediate generated *via* dissociation of one of the phosphorus atoms of the dmpe ligand is also plausible. Such a mechanism is more generally accepted for the reductive elimination from *cis*-dialkylpalladium species. See: Yamamoto, A. *Organotransition Metal Chemistry: Fundamental Concepts and Applications*; Wiley-Interscience: New York, 1986; p 240. (b) Crabtree, R. H. *The Organometallic Chemistry of the Transition Metals*, 2nd ed.; Wiley-Interscience: New York, 1988; p 151.

⁽¹⁸⁾ Yamamoto, T.; Yamamoto, A.; Ikeda, S. J. Am. Chem. Soc. 1971, 93, 3350.