

plexes will vary greatly depending on the ratio of iodine to iodide in the complex; moreover, it may be possible to determine this ratio quite unam-

biguously by the method of continuous variations, if a system can be found in which equilibrium exists. AMES, IOWA

## COMMUNICATIONS TO THE EDITOR

### THE STRUCTURE OF DIOSCORINE


Sir:

Dioscorine,  $C_{13}H_{19}O_2N$ , a water-soluble alkaloid, has been isolated from *Dioscorea hirsuta* Blume<sup>1,2</sup> and *Dioscorea hispida* Dennst.<sup>3,4</sup> The early structural investigations<sup>2</sup> were continued by Pinder<sup>3-9</sup> who has proposed three structures, including I, for dioscorine. No facts have been presented from which the structure of the nitrogenous nucleus and the location of the lactone ring present in the alkaloid could be deduced conclusively. In this communication we wish to present findings which establish structure I for dioscorine.

Reduction of the alkaloid with lithium aluminum hydride gave a diol (II), m.p. 121° (picrate, m.p. 164–165°, lit.<sup>8</sup> 159–160°), which by ozonization was converted to glycolaldehyde and a hydroxyketone (III), m.p. 25°, infrared peaks at 3540  $cm^{-1}$  (OH); 1705  $cm^{-1}$  (C=O); 1405  $cm^{-1}$  (COCH<sub>2</sub>) and 1360  $cm^{-1}$  (COCH<sub>3</sub>) (picrate, m.p. 120–122°). On exposure to 0.1 *N* sodium hydroxide, III underwent retroaldol cleavage to acetone and a ketone (IV), infrared peak at 1723  $cm^{-1}$  (CS<sub>2</sub>), 1730  $cm^{-1}$  (CCl<sub>4</sub>);  $[\alpha]^{25}_D +17^\circ$  (*c* 1.43 in H<sub>2</sub>O) (di-*p*-toluoyl-D-tartrate, m.p. 149–151°). Treatment of IV with ethanedithiol and then desulfurization produced tropone (V), picrate, m.p. 280–288° (dec.); chloroplatinate, m.p. 208–210°, identical with authentic samples.<sup>10</sup> The ketone IV is therefore tropan-2-one or tropan-6-one. We initially preferred<sup>11</sup> the 6-substituted structure, mainly because of the position of the carbonyl band in the infrared of IV. Reduction with sodium borohydride or with hydrogen over Adams catalyst yielded an alcohol (VI), m.p. 70° (picrate, m.p. 262–263°; di-*p*-toluoyl-D-tartrate, m.p. 165–166°; di-*p*-toluoyl-L-tartrate, m.p. 164–165°) which had an infrared spectrum entirely different from that of a mixture of 6 $\alpha$ - and 6 $\beta$ -hydroxytropone, m.p. 65–81° (*cf.* ref. 9) prepared by desulfurization of 6 $\beta$ -hydroxytropan-3-one thioketal. The three derivatives of the alcohol from dioscorine (I), however, were identical with those of 2 $\alpha$ -hydroxy-

tropone (VI).<sup>12</sup> Similarly the di-*p*-toluoyl-D-tartrates of IV from cocaine and dioscorine were identical. The nuclear magnetic resonance spectrum of dioscorine (in CHCl<sub>3</sub>) has no peaks above 1200 c.p.s.<sup>13</sup> and unactivated methyls are therefore absent.

The configuration at C<sub>2</sub> then was determined: The saturated diol prepared by catalytic reduction of II had a  $pK_a$  of 8.81 (Cellosolve) like 2 $\alpha$ -hydroxytropone (VI),  $pK_a$  8.42, while 2 $\beta$ -hydroxytropone (VII) had  $pK_a$  9.62. The observed differences in basicity are similar to those of quinine and epiquinine<sup>14</sup> and demonstrate that the tertiary hydroxyl in II has the  $\alpha$ -configuration which is in



agreement with the previously reported infrared evidence.<sup>8</sup> 2 $\beta$ -Hydroxytropone (VII), m.p.  $\sim -10^\circ$ ,  $n^{20}_D$  1.4862 (picrate, m.p. 263–265°) was prepared by lithium aluminum hydride reduction of 2 $\beta$ ,3 $\beta$ -epoxytropone (IX) (picrate, m.p. 248–254°) which in turn was available by oxidation of tropidine (VIII) with trifluoroacetic acid in acetonitrile.<sup>15</sup> The conversion of both cocaine<sup>12</sup> and dioscorine to the same tropan-2-one (IV) demonstrates identical absolute configuration<sup>16</sup> in the two alkaloids.

We are indebted to Eli Lilly and Co. for financial support and to Prof. E. Schlittler for plant material.

DEPARTMENT OF CHEMISTRY  
MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
CAMBRIDGE, MASSACHUSETTS

D. E. AYER  
G. BÜCHI  
P. REYNOLDS-WARNHOFF  
DWAIN M. WHITE

RECEIVED OCTOBER 6, 1958

- (1) H. W. Schütte, *Chem. Zentr.*, **68**, II, 130 (1897).
- (2) M. K. Gorter, *Rec. trav. chim.*, **30**, 161 (1911).
- (3) A. R. Pinder, *Nature*, **168**, 1090 (1951).
- (4) A. R. Pinder, *J. Chem. Soc.*, 2236 (1952).
- (5) A. R. Pinder, *ibid.*, 1825 (1953).
- (6) A. R. Pinder, *ibid.*, 1577 (1956).
- (7) A. R. Pinder, *Chemistry and Industry*, 1240 (1957).
- (8) A. R. Pinder, *Tetrahedron*, **1**, 301 (1957).
- (9) J. B. Jones and A. R. Pinder, *Chemistry and Industry*, 1000 (1958).
- (10) K. Hess, *Ber.*, **51**, 1001 (1918); R. Willstätter and F. Igler, *ibid.*, **33**, 1170 (1900).
- (11) G. Büchi, D. E. Ayer and D. M. White, XVIth Intern. Congr. of Pure and Applied Chemistry, Paris, July, 1957.

- (12) M. R. Bell and S. Archer, accompanying paper. We wish to thank Dr. S. Archer for a sample of this substance.
- (13) Relative to an arbitrarily assigned peak of 1000 c.p.s. for the aromatic proton of toluene.
- (14) V. Prelog and O. Häfliger, *Helv. Chim. Acta*, **33**, 2021 (1950).
- (15) Method of G. Fodor, J. Toth, I. Koczor, P. Dobo and I. Vincze, *Chemistry and Industry*, 764 (1956).
- (16) E. Hardegger and H. Ott, *Helv. Chim. Acta*, **38**, 312 (1955).