

apical ethylidyne group,  $\equiv\text{CCH}_3$ , is symmetrically connected to the  $\text{H}_3\text{Fe}_3(\text{CO})_9$  triangular fragment with an average C-Fe distance of 1.947 Å. The C-C bond distance in the ethylidyne group is 1.466 Å. Other bond distances and angles are normal.

The tricobalt analogue of I,  $\text{Co}_3(\text{CO})_9\text{CCH}_3$ , a member of a class of compounds that have been extensively studied,<sup>5</sup> has a smaller M-C<sub>apical</sub>-M angle (81°) and a significantly shorter M-M distance (2.467 Å).<sup>6</sup> The mixed alkylidyne cluster analogue,  $\text{HfFeCo}_2(\text{CO})_9\text{CCH}_3$ , has also been prepared and characterized.<sup>7</sup> In addition, the ruthenium<sup>3,8</sup> and osmium<sup>8,9</sup> analogues of I have been observed. The X-ray diffraction results on the former<sup>10</sup> reveal the same cluster structure as that determined here for I. The role of the apical carbon<sup>11</sup> and the role of the bridging hydrogens in bonding<sup>12</sup> are two features of cluster bonding that may be fruitfully approached by a comparative study of the electronic structures of these closely related compounds. Also, because ethylidyne has been proposed as the stable species formed from  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$  chemisorption on metal surfaces,<sup>13</sup> the properties of I are pertinent to the metal cluster-metal surface analogy<sup>14</sup> as applied to iron.

The aliphatic  $\equiv\text{CCH}_3$  moiety results from the reduction of CO originally bound to iron. This has been demonstrated by preparing I from  $^{13}\text{C}$  enriched  $\text{Fe}(\text{CO})_5$  (about 30%). The CO's and  $\text{CH}_3$  group of the product I exhibited roughly equal  $^{13}\text{C}$  enrichments as measured by  $^{13}\text{C}$  NMR. Although the specific mechanism for the formation of I is not presently known, there is ample precedent in the literature for the type of reaction that must take place.<sup>15</sup> For example, the reaction of iron bound carbonyl with hydride has been shown to yield formyl complexes,<sup>16</sup> and the reaction of boranes with iron acetyl complexes has been shown to yield alkyliron species.<sup>17</sup> Particularly pertinent is the demonstrated tautomeric interconversion between methyl and hydrido methylene ligands in a trisium carbonyl cluster and the ultimate conversion of both species to  $\text{Os}_3(\mu_2\text{-H})_3(\text{CO})_9(\mu_3\text{-CH})$ .<sup>18</sup> The conversion of iron bound carbonyl to hydrocarbons by hydride has been demonstrated, e.g.,  $\text{Cp}_2\text{Fe}_2(\text{CO})_4$ ,<sup>19</sup> and I, II, and III must be considered as possible intermediates in such reactions.

The isolation of compound I is of significant interest for other reasons as well. It is the first example of the formation of a member of this class of well-known trinuclear clusters from a mononuclear metal carbonyl. When the general decrease in thermal stability of hydrido derivatives in going from Os to Ru

to Fe is considered,<sup>20</sup> the evident stability of I is further proof of the intrinsic stability of alkylidyne trinuclear metal systems.

Details of the structure and other related investigations will be presented in a subsequent publication.

**Acknowledgment.** We thank the National Science Foundation (Grant CHE78-11600) for the support of this work and the University of Notre Dame Computer Center for providing computing time. We also thank Professor W. R. Scheidt and Dr. K. Haller for aid in the crystal structure determination and Professor J. A. Labinger for helpful discussions.


## $\pi$ - and $\sigma$ -Acetoxy Radicals

P. S. Skell\* and D. D. May

Department of Chemistry, The Pennsylvania State University  
University Park, Pennsylvania 16802

Received June 30, 1980



We report here the intermediacy of two types of acetoxy radicals in chain reactions, one as a hydrogen abstractor of low selectivity in alkane halogenation and the other undergoing decarboxylation (Hunsdiecker reaction) exclusively. This behavior is analogous to that shown by succinimidyl radicals,<sup>1</sup> and it is possible that the acetoxys may also be represented as  $\sigma$  and  $\pi$  radicals.



All previous work on acetoxy radical recognized cage decarboxylation ( $k = 1.6 \times 10^9 \text{ s}^{-1}$ )<sup>2</sup> as the major pathway, with indications that cage-wall trapping by cyclohexene solvent is the only reaction which competes with decarboxylation; no authenticated noncage reactions were recognized.<sup>3</sup> We observe acetoxy reactions outside the cage which we attribute to  $\pi$ -acetoxy and we attribute all previous work to  $\sigma$ -acetoxy. Acetoxy radical is generated in chain reactions of acetyl hypobromite ( $\text{MeCO}_2\text{Br}$ ).

Until recently acetyl hypobromite had not been available as the pure substance. We confirm its reported isolation<sup>4</sup> and have improved on the method by using Freon 11 in place of  $\text{CCl}_4$ , obtaining pure crystalline material in 50-60% yield.

In the absence of H donors, irradiation of solutions of acetyl hypobromite in Freon 11 at  $-78^\circ\text{C}$  result in quantitative conversion to methyl bromide and carbon dioxide (Hunsdiecker). These are chain reactions, both in the presence of  $10^{-3} \text{ M}$   $\text{Br}_2$  or



in solutions containing 0.1 M vinylidene chloride as  $\text{Br}_2$  scavenger.<sup>5</sup>

Brominations of alkanes were observed in oxygen-free irradiated solutions of acetyl hypobromite containing  $10^{-3} \text{ M}$   $\text{Br}_2$ . Neo-

(5) Seyferth, D. *Adv. Organomet. Chem.* **1976**, *14*, 98.

(6) Sutton, P. W.; Dahl, L. F. *J. Am. Chem. Soc.* **1967**, *89*, 261.

(7) Epstein, R. A.; Withers, H. W.; Geoffroy, G. L. *Inorg. Chem.* **1979**, *18*, 942.

(8) Keister, J. B.; Horling, T. L. *Inorg. Chem.* **1980**, *19*, 2304.

(9) Deeming, A. J.; Underhill, M. *J. Chem. Soc., Chem. Commun.* **1973**, 277. Deeming, A. J.; Hasso, S.; Underhill, M.; Canty, A. J.; Johnson, B. F. G.; Jackson, W. G.; Lewis, J.; Matheson, T. W. *Ibid.* **1974**, 807. Yesinowski, J. P.; Bailey, D. J. *Organomet. Chem.* **1974**, *65*, C27.

(10) Sheldrick, G. M.; Yesinowski, J. P. *J. Chem. Soc., Dalton Trans.* **1975**, 873.

(11) It has been suggested recently on the basis of qualitative calculations that the triangular metal fragment in the cobalt analogue is held together primarily by bonding with the apical carbon. Evans, J. *J. Chem. Soc., Dalton Trans.* **1980**, 1005.

(12) The effect of bridging hydrogens on metal-metal distances is a subject of contemporary interest. See, for example: Bau, R.; Teller, R. G.; Kirtley, S. W.; Koetzle, T. F. *Acc. Chem. Res.* **1979**, *12*, 176.

(13) Kesmodel, L. L.; Dubois, L. H.; Somorjai, G. A. *Chem. Phys. Lett.* **1978**, *56*, 267.

(14) Muetterties, E. L.; Rhodin, T. N.; Band, E.; Brucker, C. F.; Pretzer, W. R. *Chem. Rev.* **1979**, *79*, 91.

(15) The general principles of the activation of bound CO by Lewis acids have been discussed recently. Shriver, D. F.; Whitmire, K.; Stimson, R. E. "Abstracts"; 2nd Chemical Congress of North American Continent, August 1980; American Chemical Society: Washington, DC, 1980; Abstract INOR-3. In the reaction we report that a borane probably coordinates to the carbonyl oxygen and the hydride attacks the carbonyl carbon.

(16) Masters, C.; van der Woude, C.; van Doorn, J. A. *J. Am. Chem. Soc.* **1979**, *101*, 1633. Gladysz, J. A.; Tam, W. *Ibid.* **1978**, *100*, 2546. Casey, C. P.; Neumann, S. M. **1978**, *100*, 2544. Collman, J. P.; Winter, S. R. *Ibid.* **1973**, *95*, 4089.

(17) van Doorn, J. A.; Masters, C.; Volger, H. C. *J. Organomet. Chem.* **1976**, *105*, 245.

(18) Calvert, R. B.; Shapley, J. R. *J. Am. Chem. Soc.* **1977**, *99*, 5226.

(19) Wong, A.; Atwood, J. D. *J. Organomet. Chem.* **1980**, *199*, C9.

(20) Kesz, H. D.; Saillant, R. B. *Chem. Rev.* **1972**, *72*, 231.

Table I. Photoinitiated Halogenations of 1:1 Neopentane-Methylene Chloride Mixtures

halogenating agent	yields <sup>a</sup>				account-ability <sup>c</sup>
	CO <sub>2</sub>	neo-C <sub>5</sub> H <sub>11</sub> X	CHCl <sub>2</sub> X	selectivity <sup>b</sup>	
AcOBr/Br <sub>2</sub> <sup>e</sup>	77 <sup>d</sup>	7	14	12	98
AcOCl/Cl <sub>2</sub> <sup>e</sup>	66	14	28	12	96
AcOBr, Cl <sub>2</sub> C=CH <sub>2</sub> <sup>f</sup>	100	0	0		100
AcOCl, Cl <sub>2</sub> C=CH <sub>2</sub> <sup>f</sup>	98	0	0		98

<sup>a</sup> Yields determined by GC and expressed as percentages based on starting hypohalite. <sup>b</sup> Selectivity calculated from the molar ratio (CHCl<sub>2</sub>X/neo-C<sub>5</sub>H<sub>11</sub>X) times (12 neopentane hydrogens/2 CH<sub>2</sub>Cl<sub>2</sub> hydrogens) to account for the statistical factor. <sup>c</sup> (mmol of CO<sub>2</sub> + mmol of halogenated product)/mmol of AcOX)100. <sup>d</sup> Yield of methyl bromide was 75%; yield of acetic acid was 25%. <sup>e</sup> 10<sup>-3</sup> M. <sup>f</sup> 0.1 M.

pentane/methylene chloride mixtures were used. Typically these reactions (tungsten lamp irradiation, 2-3 h for 100% conversion, -78 °C, CCl<sub>3</sub>F solvent, 0.2 M AcOBr, 10<sup>-3</sup> M Br<sub>2</sub>, 1 M neo-C<sub>5</sub>H<sub>12</sub>, and 1 M CH<sub>2</sub>Cl<sub>2</sub>) give CO<sub>2</sub> (77% yield), bromination products (21%), and acetic acid (25%). This appears to be the first report of a noncage reaction of acetoxy radical. The ratio of bromodichloromethane to neopentyl bromide is 2 ± 0.3.

In the absence of acetyl hypobromite, attempts at photobromination failed under the above conditions using various concentrations of Br<sub>2</sub>; there was no bromination.

The same reaction conditions using acetyl hypochlorite give similar results in the presence of 10<sup>-3</sup> M chlorine—66% CO<sub>2</sub> and 40% chlorination products. The ratio of chloroform to neopentyl chloride is 2 ± 0.3. Chlorine alone under these conditions produces neopentyl chloride exclusively; the reaction of Cl<sub>2</sub> with CH<sub>2</sub>Cl<sub>2</sub> is not a significant pathway in this competition.






AcOBr/Br<sub>2</sub> under the above reaction conditions brominates 1-bromobutane—68% CO<sub>2</sub> and 32% halogenation products, equal amounts of 1,1-dibromobutane (11%), 1,2-dibromobutane (11%), and 1,3-dibromobutane (11%) are obtained; 1,4-dibromobutane is not produced. Bromine atom chains, if they operated at these low temperatures, would produce almost exclusively 1,2-dibromobutane (by extrapolation from higher temperatures).<sup>6a</sup> Little discrimination between C-2 and C-3 is a characteristic of radicals such as succinimidyli,<sup>1</sup> Cl<sub>2</sub>,<sup>6b</sup> and *t*-BuO<sup>•</sup>.<sup>6c</sup>

It is remarkable that in the above experiments there is no halogenation of substrates if free halogen is scavenged. Experiments in which vinylidene chloride (0.1 M) was employed as scavenger for the halogens, all other experimental details being the same, resulted in quantitative Hunsdiecker reactions, without detectable attack on neopentane, methylene chloride, or 1-bromobutane.

The possibility that chain carriers other than acetoxy are participating must be considered: Bromine atoms do not react at -78 °C. Chlorine atoms are excluded because they show a totally different selectivity. Methyl radicals are the intermediates in the Hunsdiecker, and yet there is no halogenation of substrate. This leaves acetoxy radicals as the probable intermediate, a conclusion which is supported by observing the same selectivity for both AcOBr and AcOCl.

A reasonable hypothesis is that two different types of acetoxy are produced, one, produced in the presence of halogens, which abstracts hydrogen atoms beyond the cage with low selectivity and the other, produced in halogen-scavenged systems, which undergoes decarboxylation too rapidly for noncage encounters. The latter type of behavior had been predicted for the <sup>2</sup>A<sub>1</sub> σ state on the grounds that <sup>2</sup>A<sub>1</sub> correlates with ground-state CO<sub>2</sub>.<sup>7</sup> The acetoxy which accounts for hydrogen atom abstraction can be

assigned the <sup>2</sup>B<sub>2</sub> σ state or the <sup>2</sup>A<sub>2</sub> π state;<sup>7-9</sup> neither of these correlates with ground-state CO<sub>2</sub>, and thus it is possible for them to escape the cage and engage in relatively nondiscriminate hydrogen atom abstractions. For convenience, the notation π is given to the hydrogen atom abstracting species and σ to the other acetoxy (<sup>2</sup>A<sub>1</sub>). The σ state (<sup>2</sup>A<sub>1</sub>) would result from attack of CH<sub>3</sub>• on CH<sub>3</sub>CO<sub>2</sub>X and the π state (hydrogen atom abstracting intermediate) from attack of halogen atom on CH<sub>3</sub>CO<sub>2</sub>X.



The calculations concerned with carboxylate radicals lead to no single conclusion about the ordering of the states;<sup>7-9</sup> they have been done at too simple a level. It is hoped that suitable calculations will become available in the future. We expect that experiments which are in progress will provide an experimental basis for energy sorting the acetoxy states.

**Acknowledgment.** Support of this research by the National Science Foundation under Grant CHE-7810049 is gratefully acknowledged.

## Chiral Aggregation Phenomena. 2. Evidence for Partial "Two-Dimensional Resolution" in a Chiral Monolayer

Edward M. Arnett\* and Orlean Thompson\*

Department of Chemistry, University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Received May 1, 1980

We wish to report the results of a novel attempt to resolve a racemic monolayer film of a chiral surfactant spread on an aqueous subphase by seeding with crystals of one of the enantiomers. Although the investigation of monolayers at the gas-liquid interface is a well-studied and currently active field,<sup>1</sup> there have been relatively few reports of stereochemical phenomena in films cast from chiral surfactants.<sup>2</sup> In the course of comparing the surface properties of racemic *N*-α-methylbenzylstearamide with those of its two pure enantiomers,<sup>3</sup> we noted a considerably higher surface free energy for the racemic films on aqueous sulfuric acid subphases than for comparable films of the enantiomers. This was manifested both in terms of higher surface pressures at

\* Address correspondence to this author at the Department of Chemistry, Mail Gross Laboratories, Duke University, Durham, N.C. 27706.

\* These results and experimental conditions are described more fully in the doctoral thesis of Orlean Thompson, Department of Chemistry, University of Pittsburgh, 1980.

(1) Several excellent texts which review this field are: (a) Gaines, G. L. "Insoluble Monolayers at Liquid-Gas Interfaces", Wiley: New York, 1966. (b) Adam, N. K. "The Physics and Chemistry of Surfaces", 3rd ed.; Oxford University Press: London, 1941. (c) Davies, J. T.; Rideal, E. K. "Interfacial Phenomena", 2nd ed.; Academic Press: New York, 1963. (d) Adamson, A. W. "Physical Chemistry of Surfaces", 3rd ed.; Wiley-Interscience: New York, 1976. (e) Cadenhead, D. A. In "Chemistry and Physics of Interfaces, II"; Ross, S., Ed.; American Chemical Society: Washington, D.C., 1971, p 28. (f) Gershfeld, N. L. *Annu. Rev. Phys. Chem.* **1976**, *27*, 349-368. This is an excellent recent review of the field, and subjects some commonly held assumptions to close scrutiny. (g) Goddard, E. D. *Adv. Chem. Ser.* **1975**, *No. 144*, 1.

(2) For a review of previously published studies of chiral monolayers, see: Stewart, M. V.; Arnett, E. M. *Top. Stereochem.* **1981**, *13*, 1.

(3) Arnett, E. M.; Chao, J.; Kinzig, B.; Stewart, M.; Thompson, O. J. *Am. Chem. Soc.* **1978**, *100*, 5575-5576.

(6) (a) Skell, P. S.; Shea, K. J. *Am. Chem. Soc.* **1972**, *94*, 6551. (b) Thaler, W. A. *Ibid.* **1963**, *85*, 2607. (c) Walling, C.; Jacknow, B. B. *Ibid.* **1960**, *81*, 6113.

(7) Koenig, T.; Wielesek, R. A.; Huntingdon, J. G. *Tetrahedron Lett.* **1974**, *26*, 2283.

(8) Kikuchi, O. *Tetrahedron Lett.* **1977**, *28*, 2421.

(9) McBride, J. M.; Merrill, R. A. *J. Am. Chem. Soc.* **1980**, *102*, 1723.