

ALKYLATION OF ETHYL BROMOMALONATE BY ALKYLCOBALOXIMES *

M. VEBER, K. N. V. DUONG, F. GAUDEMER and A. GAUDEMER

*Laboratoire de Chimie de Coordination Bioorganique, CNRS (LA 255) Université Paris-Sud,
 Centre d'Orsay, 91405 Orsay (France)*

(Received April 4th 1979)




Summary

Allylcobaloximes react under very mild conditions with ethyl bromomalonate to yield allyl-substituted ethylmalonates in good yield. In the case of crotyl-, 3,3 dimethyl-, allyl- and cinnamyl-cobaloximes, the substitution occurs with total rearrangement of the allyl groups. Similar rearrangements are observed during the reactions of propargyl- and allenylcobaloximes with $\text{BrCH}(\text{CO}_2\text{Et})_2$ yielding allenyl- and propargyl-malonic esters respectively.

Introduction

Alkylcobaloximes $\text{R}-\text{Co}(\text{DmgH})_2\text{Py}$ ** have recently been shown to react with organic compounds which are considered to be good radical precursors: polyhaloalkanes [1], diphenyl disulphide [2], diphenyl diselenide [2] and alkylsulphonyl chlorides [3]. These reactions give the products $\text{R}-\text{CCl}_3$, $\text{R}-\text{SPh}$, $\text{R}-\text{SePh}$ and $\text{R}-\text{SO}_2-\text{R}'$ respectively, but their mechanisms are not fully understood. It has been proposed that they proceed by a chain radical mechanism (e.g. Scheme 1 for the diphenyldisulphide reaction [2]).

Scheme 1



The inversion of configuration observed during the reaction of diphenyldi-

* Dedicated to Prof. Henri Normant on the occasion of his 72nd birthday

** Alkylcobaloximes $\text{RCo}(\text{DmgH})_2\text{Py}$ = bis(dimethylglyoximate) alkylpyridinecobalt (III)

sulphide with optically active 2-octylcobaloxime and the rate increase produced by visible light irradiation are in agreement with this mechanism

In the present paper, we describe the results obtained in a study of the reactions between various alkylcobaloximes and ethyl bromomalonate this compound can be easily reduced to the free malonyl radical by reducing organometallic complexes such as $\text{Fe}(\text{CO})_5$ [4] or non (II) porphyrins [5] and an a priori reaction mechanism leading to the alkylmalonic esters may be envisaged (Scheme 2).

Scheme 2

Initiation: $\text{R}-\text{Co}(\text{DmgH})_2\text{Py} \rightarrow \text{Co}^-(\text{DmgH})_2\text{Py} + \text{Organic products}$

Propagation: $\text{Co}^-(\text{DmgH})_2\text{Py} + \text{BrCH}(\text{CO}_2\text{Et})_2 \rightarrow \text{BrCo}(\text{DmgH})_2\text{Py} + \text{CH}(\text{CO}_2\text{Et})_2$

$$\text{CH}(\text{CO}_2\text{Et})_2 + \text{R}-\text{Co}(\text{DmgH})_2\text{Py} \rightarrow \text{R}-\text{CH}(\text{CO}_2\text{Et})_2 + \text{Co}(\text{DmgH})_2\text{Py}$$

In the case of allylcobaloximes the reaction was expected to give allyl-substituted malonates with high regioselectivity, by analogy with the reaction with BrCCl_3 [1]

Results and discussion


Reactions of allylcobaloximes with ethyl bromomalonate

Allylcobaloximes I to V react with ethyl bromomalonate in CHCl_3 at, or just above, room temperature to give ethyl allylmalonates VI to X (Scheme 3). Structures of the esters were assigned from their spectroscopic data, and the NMR spectra of esters VIII, IX and X in particular revealed the characteristic multiplets of a vinyl group $\text{CH}=\text{CH}_2$. In the course of the reactions leading to these compounds there is a complete rearrangement of the allyl group initially present in the cobaloxime. The allylmalonic esters are always accompanied by variable amounts of ethyl malonate but neither allylmalonic esters having an unrearranged allyl group nor products of diallylation could be detected. These results demonstrate the total regioselectivity of the reaction and are to be compared with those of Johnson et al. [1] for polyhaloalkanes.

Crude yields of ethyl allylmalonates ranged between 50 and 80% and a study of the influence of temperature and of bromomalonate concentration was carried out with 3,3-dimethylallylcobaloxime IV in order to define the best experimental conditions. An increase in the reaction temperature (Table 1) from 20 to 60°C resulted in a decrease in the yield of ester IX. On the other hand, an increase of the ratio ethyl bromomalonate/complex IV gave an increase in the yield of ethyl malonate with respect to ester IX (Table 2). Thus the best yields of the latter compound were obtained by running the reaction at room temperature with equal concentrations of ethyl bromomalonate and allylcobaloximes.

Irradiation of the reaction mixture with visible light from a tungsten lamp resulted in an increase of the reaction rate. For example compound IV is completely converted into a mixture of ester IX (69%) and ethyl malonate after 17

SCHEME 3



I, VI $R_1 = R_2 = R_3 = H$, II, VII $R_1 = R_2 = H$, $R_3 = CH_3$, III VIII $R_1 = R_3 = H$, $R_2 = CH_3$,
 IV, IX $R_1 = R_2 = CH_3$, $R_3 = H$, V, X $R_1 = R_3 = H$, $R_2 = C_6H_5$





TABLE 1

INFLUENCE OF TEMPERATURE ON THE YIELD OF ESTER IX IN THE REACTION OF COMPLEX IV (0.2 mmol) WITH ETHYL BROMOMALONATE (0.3 mmol) IN CHLOROFORM

Temp, °C)	Reaction time	Ester IX (%)	Ethyl malonate (%)
20	2 days	83	17
40	6 hours	75	25
60	30 min	70	30

TABLE 2

INFLUENCE OF THE CONCENTRATION OF ETHYL BROMOMALONATE ON THE YIELD OF ESTER IX IN THE REACTION WITH COMPLEX IV (0.2 mmol) IN CHLOROFORM AT 60°C

Concn. of ethyl bromomalonate (mmol)	Ester IX (%)	Ethyl malonate (%)
0.2	70	30
0.4	68	32
0.6	66	33
0.8	64	35

One of the reaction products was biomocobaloxime ($\text{B1Co(DmgH)}_2\text{Py}$) (yield 36%) in agreement with the proposed mechanism.

This mechanism rationalises most of the features of these reactions but does not account for the high regioselectivity observed with complexes III, IV and V. A similar pathway can be envisaged for the reaction of allenyl- or propargylcobaloximes. However, we have no experimental evidence for the formation, during thermolysis or photolysis of these compounds, of cobaloxime (II), the initiating entity of the chain mechanism, whereas such evidence does exist in the case of allylcobaloximes [9].

Conclusion

The results reported herein demonstrate the synthetic value of these reactions. Several methods have been proposed to prepare allylmalonates which are important intermediates in the synthesis of barbiturates. Allylation of sodium malonate by allyl halides has been widely used [10] and more recently other reactions, which involve transition metal salts or complexes, have been reported: e.g. allylation of ethyl malonate by alkenes in the presence of Mn^{III} (or Co^{III}) and Cu^{II} salts [11], and nucleophilic attack of sodium malonate on π -allyl complexes of Ni^{II} [12] or Pd^{II} [13]. These reactions give rise, in most cases, to mixtures in which the product having the most substituted double bond is predominant. The reactions of allylcobaloximes with ethyl bromomalonate provide an alternative method of synthesis of allylmalonic esters and its interest lies in the possibility of preparing compounds in which the malonyl group is bound to secondary or tertiary carbon atoms.

Experimental

Chemicals and instrumentation

Allyl-, allenyl- and propargylcobaloximes were known compounds which were prepared by treating pyridinecobaloxime (I) with allylic or propargylic halides from commercial sources. Ethyl bromomalonate (Fluka), containing 6% ethyl malonate, was used without further purification. Infrared spectra of reaction products were recorded neat with a Beckman Acculab 6 spectrophotometer. ^1H NMR spectra at 90 MHz were measured with a Perkin-Elmer R-32 spectrometer in CDCl_3 solution containing 1% TMS. Mass spectra were obtained with an AEI MS 50 spectrometer (ICSN - Gif-sur-Yvette, France).

General procedure for the reactions between ethyl bromomalonate and alkylcobaloximes

In a typical reaction, 1 mmol of alkylcobaloxime and 1.5 mmol of ethyl bromomalonate were added to 6 ml of degassed CHCl_3 in the dark. The mixture was left at room temperature for 48 h under nitrogen and then passed through a column of Silica gel CC7. Elution by CH_2Cl_2 allowed a crude separation of the organic products: unreacted ethyl bromomalonate, ethyl malonate and ethyl allylmalonate from bromopyridinecobaloxime. The reaction yields were deduced from the integration curve of the NMR spectra of the mixtures of organic products. Pure allyl-, propargyl- and allenyl-malonates were obtained

either by chromatography on an alumina (activity 1) column (eluant hexane and hexane/ethyl acetate 95/5) or by preparative thin-layer chromatography on Silica gel (Merck F-254) (eluant hexane/ethyl acetate 1/1)

Photochemical reactions

0.2 mmol of alkylcobaloxime and 0.3 mmol of ethyl bromomalonate were dissolved in 0.7 ml degassed CDCl_3 in an NMR tube filled with N_2 and closed by a serum cap. The tube was irradiated in a Pyrex vessel cooled at 0°C , by two 300 W tungsten lamps. The reactions were followed by measuring the NMR spectrum of the mixture at regular intervals.

Ethyl allylmalonate (VI) (yield 50%) was identified by comparison of its NMR spectrum with that of an authentic sample and by its mass spectrum m/e 200 (M^+), 155, 127.

Ethyl (2-methyl) allylmalonate (VII) (yield 71%). IR (cm^{-1}) $\nu(\text{CH})$ 3040, $\nu(\text{C=O})$ 1735, $\nu(\text{C=C})$ 1650; NMR δ 4.79, 4.72 (2s, 2H, C=CH_2), 4.18 (q, 4H, O-CH_2), 3.53 (t, 1H, $\text{CH}(\text{CO}_2\text{C}_2\text{H}_5)_2$), 2.62 (d, 2H, $\text{CH}_2\text{-CH}$), 1.75 (s, 3H, $\text{CH}_3\text{-C=}$) and 1.27 ppm (t, 6H, $\text{O-CH}_2\text{-CH}_3$), mass spectrum m/e 214 (M^+), 159, 141, 55.

Ethyl (1-methyl) allylmalonate (VIII) (yield 60%). IR (cm^{-1}) $\nu(\text{CH})$ 3010, $\nu(\text{C=O})$ 1730, $\nu(\text{C=C})$ 1650, NMR δ 5.83 (m, 1H, CH=C), 5.10 (m, 2H, $\text{CH}_2=\text{C}$), 4.22 (q, 4H, O-CH_2), 3.28 (d, 1H, $\text{CH}(\text{CO}_2\text{C}_2\text{H}_5)_2$), 2.95 (m, 1H, $\text{CH}(\text{CH}_3)$), 1.28 (t, 6H, $\text{O-CH}_2\text{-CH}_3$) and 1.12 ppm (d, 3H, $\text{CH}_3(\text{CH})$), mass spectrum m/e : 214 (M^+), 169, 160, 141, 55.

Ethyl (1,1-dimethyl) allylmalonate (IX) (yield 75%). IR (cm^{-1}) $\nu(\text{CH})$ 3035, $\nu(\text{C=O})$ 1740, $\nu(\text{C=C})$ 1635; NMR δ 6.08 (q, 1H, CH=C), 5.00 (m, 2H, $\text{CH}_2=\text{C}$), 4.16 (q, 4H, O-CH_2), 3.30 (s, 1H, $\text{CH}(\text{CO}_2\text{C}_2\text{H}_5)_2$), 1.26 (t, 6H, $\text{O-CH}_2\text{-CH}_3$) and 1.26 ppm (s, 6H, $(\text{CH}_3)_2\text{C}$); mass spectrum m/e 228 (M^+), 183, 160, 155, 69.

Ethyl (1-phenyl) allylmalonate (X) (yield 52%). IR (cm^{-1}) $\nu(\text{CH})$ 3010, $\nu(\text{C=O})$ 1725, $\nu(\text{C=C})$ 1500, 1600, 1635, NMR δ 7.22 (s, 5H, C_6H_5), 6.00 (m, 1H, CH=C), 5.10 (m, 2H, $\text{CH}_2=\text{C}$), 4.21 (q, 4H, O-CH_2), 3.85 (m, 2H, -CH-CH-) and 1.27 ppm (t, 3H, $\text{O-CH}_2\text{-CH}_3$), mass spectrum m/e 276 (M^+), 203, 117, 91.

Ethyl (3-methyl) allenylmalonate (XII) (yield 70%). IR (cm^{-1}) $\nu(\text{C=O})$ 1740, $\nu(\text{C=C=C})$ 1960; NMR δ 4.70 (m, 8 lines, 2H, $\text{CH}_2=\text{C}$, J 1.6 Hz, J 3 Hz), 4.14 (q, 4H, O-CH_2), 4.01 (t, 1H, $\text{CH}(\text{CO}_2\text{C}_2\text{H}_5)_2$, J 1.6 Hz), 1.82 (t, 3H, $\text{CH}_3\text{-C}$; J 3 Hz) and 1.30 ppm (t, 6H, $\text{O-CH}_2\text{-CH}_3$), mass spectrum m/e 212 (M^+), 167, 139, 110, 65.

Ethyl (1,1-dimethyl) propargylmalonate (XIV) (yield 30%). IR (cm^{-1}) $\nu(\text{CH})$ 3105, $\nu(\text{C=O})$ 1745; NMR δ 4.22 (q, 4H, O-CH_2), 3.42 (s, 1H, $\text{CH}(\text{CO}_2\text{C}_2\text{H}_5)_2$), 2.19 (s, 1H, $\text{HC}\equiv\text{C}$), 1.46 (s, 6H, $(\text{CH}_3)_2\text{C}$) and 1.29 ppm (t, 6H, $\text{O-CH}_2\text{-CH}_3$), mass spectrum m/e : 226 (M^+), 181, 160, 153, 67.

Acknowledgment

We thank Dr. M. D. Johnson for numerous fruitful discussions. Acknowledgment is made to the Délégation Générale à la Recherche Scientifique et Technique (Grant no. 77 7 0761) for support of this research.

References

- 1 A L Crease B D Gupta M D Johnson and S Moorhouse J Chem Soc Perkin Trans II in press
- 2 K N V Duong A Gaudemer B D Gupta and M D Johnson J Chem Soc Perkin II in press
- 3 A E Crease B D Gupta M D Johnson E Bialkowska K N-V Duong and A Gaudemer J Chem Soc Perkin I in press
- 4 R K Freidlina and F K Velichko Synthesis (1978) 145
- 5 R S Wade and C E Castro J Amer Chem Soc 95 (1973) 226
- 6 J P Collman J N Cawse and J W Kaug Inorg Chem 8 (1969) 2574 C J Cooksey D Dodd C Gatford M D Johnson G J Lewis and D M Titchmarsh J Chem Soc Perkin II (1972) 655
- 7 P D Bartlett and C Ruckardt J Amer Chem Soc 82 (1960) 1756
- 8 K U Ingold Free Radicals Vol I 1973 p 37
- 9 E Bialkowska K N V Duong and A Gaudemer, unpublished results
- 10 A E Cope H L Holmes and H O House Org React 9 (1957) 107
- 11 G I Nikishin M G Vinogradov and T M Fedorova J Chem Soc Chem Commun (1973) 693
- 12 R Baker A H Cook and M J Crimmin J Chem Soc Chem Commun (1975) 727
- 13 J Tsuji Bull Chem Soc Japan 46 (1973) 1897 B M Trost L Weber P E Strege T J Fullerton and T J Ditsche J Amer Chem Soc 100 (1978) 3407 3416 and 3426
- 14 C Fontaine K N V Duong C Merienne A Gaudemer and C Giannotti J Organometal Chem 38 (1972) 167