


This article was downloaded by: [New York University]

On: 10 October 2014, At: 01:41

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Chiral Bis(oxazoline)-copper Catalyzed Enantioselective Imidation of Sulfides

Hiroya Takada , Yoshiaki Nishibayashi , Kouichi Ohe & Sakae Uemura

^a Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering , Kyoto University , Sakyo-ku, Kyoto 606-01, Japan

^b Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering , Kyoto University , Sakyo-ku, Kyoto 606-01, Japan

^c Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering , Kyoto University , Sakyo-ku, Kyoto 606-01, Japan

^d Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering , Kyoto University , Sakyo-ku, Kyoto 606-01, Japan

Published online: 17 Mar 2008.

To cite this article: Hiroya Takada , Yoshiaki Nishibayashi , Kouichi Ohe & Sakae Uemura (1997) Chiral Bis(oxazoline)-copper Catalyzed Enantioselective Imidation of Sulfides, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 120:1, 363-364, DOI: [10.1080/10426509708545550](https://doi.org/10.1080/10426509708545550)

To link to this article: <http://dx.doi.org/10.1080/10426509708545550>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and

should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Chiral Bis(oxazoline)-copper Catalyzed Enantioselective Imidation of Sulfides

HIROYA TAKADA, YOSHIAKI NISHIBAYASHI, KOUICHI OHE
AND SAKAE UEMURA

*Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering,
Kyoto University, Sakyo-ku, Kyoto 606-01, Japan*

Prochiral sulfides reacted with $\text{PhI}=\text{NTs}$ in the presence of a catalytic amount of $\text{Cu}(\text{I})$ salt together with a chiral 4,4'-disubstituted bis(oxazoline) ligand to afford the corresponding chiral sulfimides.

KEYWORDS: sulfide, sulfimide, sufenamides, bis(oxazoline) ligand

INTRODUCTION

Quite recently, chiral sulfimides were demonstrated to be useful in organic synthesis as methylenedioxy transfer reagents to prochiral carbonyl groups, leading to optically active epoxides.¹ Despite the fact that other optically active organosulfur compounds such as sulfonium ylides and sulfoxides have many asymmetric synthetic applications, the chemistry of their nitrogen analogues, sulfimides, has been much less investigated.^{2,3} We now report that the asymmetric reaction of various sulfides **1** with $\text{PhI}=\text{NTs}$ in the presence of $\text{Cu}(\text{I})$ catalyst derived from CuOTf and the chiral 4,4'-disubstituted bis(oxazoline) ligand **3** produces the corresponding chiral sulfimides **2**.⁴

RESULTS AND DISCUSSION

Our initial attempts were to explore and exploit a new imidation method of sulfides **1** with $\text{PhI}=\text{NTs}$ using $\text{Cu}(\text{I})$ salt as catalyst, eqn. (1). The synthesis of sulfimides was carried out by treatment of suitable sulfides with 1 equiv $\text{PhI}=\text{NTs}$ in MeCN as solvent in the presence of 5 mol% of $\text{Cu}(\text{I})$ triflate (CuOTf). Representative results are summarized in Table I. Treatment of methyl *p*-tolyl sulfide with $\text{PhI}=\text{NTs}$ at 25 °C for 26 h afforded the corresponding sulfimide in 83 % isolated yield (entry 1). Similarly, a variety of sulfides could be converted smoothly to the corresponding sulfimides.

In addition, using a chiral bis(oxazoline) ligand we applied the direct imidation of various sulfides into optically active sulfimides. Thus, in the imidation of benzyl 1-





TABLE II
Synthesis of chiral sulfimides.

1		2	
R ¹	R ²	yield / %	ee / %
1-Nap	Bn	75	71
Ph	Bn	78	64
PhCH ₂ CH ₂	Bn	63	22
4-NO ₂ C ₆ H ₄ CH ₂	Ph	37	25
Ph	<i>i</i> -Pr	44	23

TABLE I
Catalytic synthesis of sulfimides.


1		2
R ¹	R ²	yield / %
p-Tol	Me	83
Ph	Ph	79
Ph	Bn	82
PhCH ₂ CH ₂	Bn	50
4-MeOC ₆ H ₄	Bn	70
1-Nap	Bn	54
4-NO ₂ C ₆ H ₄ CH ₂	Ph	72

TABLE III
Synthesis of chiral sulfenamides.

4		6	
R ³	R ⁴	yield / %	ee / %
Ph	Ph	40	27
2-NO ₂ C ₆ H ₄	Bn	37	43
1-Nap	Ph	80	58
Ph	Me	30	25
1-Nap	Me	35	50

naphthyl sulfide, the best enantiomeric excess (71 % ee) was obtained when the reaction was performed in toluene in the presence of 6 mol% chiral ligand 3 and 5 mol% CuOTf at 25 °C for 48 h to afford the corresponding chiral sulfimide in 75 % isolated yield (Table II). Other results are also shown in the table.

When the reaction was applied to allylic sulfides 4, the expected aziridination to the double bond did not occur at all and, instead, the chiral allylic sulfenamides 6 were obtained selectively in good yields, Table III. This fact shows that the nitrogen attack occurred only at the sulfur atom to give the chiral allylic sulfimide intermediates 5 and the [2,3] sigmatropic rearrangement with chirality transfer followed, eqn. (2) (Table III).



REFERENCES

- 1 C. P. Baird and P. C. Taylor, *J. Chem. Soc., Chem. Commun.*, 893 (1995).
- 2 D. J. Cram, J. Day, D. R. Rayner, D. M. von Schriltz, D. J. Duchamp and D. C. Garwood, *J. Am. Chem. Soc.*, **92**, 7369 (1970).
- 3 R. Annunziata, M. Cinquini, S. Colonna and F. Cozzi, *J. Chem. Soc., Perkin 1*, 3118 (1981).
- 4 H. Takada, Y. Nishibayashi, K. Ohe and S. Uemura, *J. Chem. Soc., Chem. Commun.*, 931 (1996).