

2-Aralkoxyadenosines: Potent and Selective Agonists at the Coronary Artery A₂ Adenosine Receptor

Masayuki Ueeda, Robert D. Thompson, Luis H. Arroyo, and Ray A. Olsson*


Department of Internal Medicine, University of South Florida, 12901 Bruce B. Downs Boulevard, Tampa, Florida 33612.
Received August 27, 1990

A Langendorff guinea pig heart preparation served for the assay of agonist potency of a series of 26 2-aralkoxyadenosines at the A₁ and A₂ receptors of, respectively, the atrioventricular node (conduction block) and coronary arteries (vasodilation). All of the analogues are weak agonists at the A₁ receptor, requiring concentrations >9 μM to cause second degree heart block. At the A₂ receptor 2-phenethoxyadenosine is the most potent of the 2-phenylalkyladenosines. The activity of ring-substituted (F, Cl, CH₃, and OCH₃) 2-phenethoxyadenosines increases ortho < meta < para. The EC₅₀s of coronary vasoactivity of several para-substituted analogues are in the subnanomolar range. The most potent analogue, 2-[2-(4-methylphenyl)ethoxy]adenosine 19, has an EC₅₀ for coronary vasodilation of 190 pM and an A₁/A₂ selectivity ratio of 44000. Aryl groups such as thienyl, indoloyl, or naphthyl also support A₂ agonist activity. Although 2-oxoadenosine is 3 times more vasoactive than 2-aminoadenosine, the activities of the phenyl derivatives are markedly different; 2-phenoxyadenosine is 23 times weaker than 2-(phenylamino)adenosine (CV-1808).

A previous report¹ describes the synthesis of a series of 2-alkoxyadenosines and bioassays of their agonist potencies at the A₁ and A₂ adenosine receptors (A₁AR and A₂AR) in the AV node and coronary arteries, respectively, of guinea pig hearts. Analysis of the structure-activity relationships (SAR) of these 2-alkoxyadenosines yielded provisional models of the C-2 regions of the A₁AR and A₂AR (Figure 1). Features common to both receptors include (a) an X subregion that accommodates the oxygen atom that links the alkyl substituent to purine C-2 and (b) an alkyl subregion of very limited bulk tolerance that seems to interact negatively with short *n*-alkyl substituents and even more poorly with wide *sec*-alkyl groups. A prominent feature of the A₂AR is a hydrophobic subregion that accommodates cycloalkyl and bicycloalkyl groups. The interaction of such groups with the hydrophobic subregion overcomes the negative influence of interaction with the alkyl subregion and can promote strong agonist activity. By contrast, such substituents do not promote the agonist activity of adenosine at the A₁AR. Such a result implies that either the A₁AR lacks a hydrophobic subregion or, alternatively, that binding to such a subregion does not promote activity.

The present study extends the mapping of the C-2 regions of the A₁AR and A₂AR by examining the SAR of several types of 2-aralkoxyadenosines. The only previous study of 2-(ar)alkoxyadenosines is that of Marumoto et al.,² who report the synthesis of 2-phenoxyadenosine, three of its ring-substituted derivatives, and 2-phenoxyethoxyadenosine. In the open-chest-dog preparation used for bioassays, all were weaker coronary vasodilators than adenosine. The same authors also evaluated the 2-benzyl, 2-benzylamino, 2-phenethylamino and 2-benzylthio derivatives of adenosine and found that these analogues, too, were moderately to markedly less active than adenosine. Because this investigation antedated the discovery of A₁AR and A₂AR,^{3,4} the authors did not evaluate selectivity. Subsequent comparisons of the activity of 2-(phenylamino)adenosine (CV-1808) showed that this analogue is 5-20-fold selective for the A₂AR.^{5,6} Recent work shows

Scheme I



that the 2-phenethylamino derivative of *N*-ethyladenosin-5'-uronamide (NECA) and several of its ring-substituted analogues are both potent and selective agonists at the A₂AR.⁷ Although the number of analogues included in that report is not large, their high potency and selectivity urges further studies of the C-2 region of the A₂AR such as those described below.

Results and Discussion

Chemistry. The reaction of the 2',3'-*O*-isopropylidene or the 2',3'-*O*-ethoxymethylidene ketal of 2-chloroadenosine (1) with either sodium phenolate or an lithium alkoxide^{1,2} furnished all the analogues described below (Scheme I). The alcohols were either commercially available or were obtained through the reduction of the corresponding acyl chloride with LiAlH₄. For most analogues, workup consisted of the removal of unreacted alcohol by flash chromatography, hydrolysis of the blocking group, and purification by preparative reverse-phase HPLC. Analogues 12 and 13, the 3- and 4-chloro derivatives of 2-phenethoxyadenosine, are exceptions, crystallizing spontaneously after the deblocking step. Owing to the susceptibility to acid hydrolysis of the glycosylic bond of the 2-(ar)alkoxyadenosines,¹ substantial losses of product occurred during the deblocking step. The discovery near the end of this project of a satisfactory procedure for deblocking, namely, refluxing in a solution of 1-2% formic acid and 50% acetic acid in water *with close monitoring by HPLC*, improved yields.


We previously found that alkoxides containing a basic nitrogen atom did not displace the chlorine atom from blocked 2-chloroadenosine.¹ Such was also the case with the pyridyl ethoxides, which contain a basic nitrogen. The nitrogen of indole is not basic and, accordingly, the lithium

- Ueeda, M.; Thompson, R. D.; Arroyo, L. H.; Olsson, R. A. *J. Med. Chem.*, previous paper in this issue.
- Marumoto, R.; Yoshioka, Y.; Miyashita, O.; Shima, S.; Imai, K.; Kawazoe, K.; Honjo, M. *Chem. Pharm. Bull.* 1975, 23, 759-774.
- Londos, C.; Cooper, D. M. F.; Wolff, J. *Proc. Natl. Acad. Sci. U.S.A.* 1980, 77, 2551-2554.
- Van Calker, D.; Muller, M.; Hamprecht, B. *J. Neurochem.* 1979, 33, 999-1005.
- Bruns, R. F.; Lu, G. H.; Pugsley, T. A. *Mol. Pharmacol.* 1986, 29, 331-346.

(6) Ukena, D.; Olsson, R. A.; Daly, J. W. *Can. J. Physiol. Pharmacol.* 1987, 65, 365-376.

(7) Hutchinson, A. J.; Williams, M.; de Jesus, R.; Yokoyama, R.; Oei, H. H.; Ghai, G. R.; Webb, R. L.; Zogamas, H. C.; Stone, G. A.; Jarvis, M. F. *J. Med. Chem.* 1990, 33, 1919-1924.

Table I. Physical and Analytical Data for Analogues 1-26



no.	R	formula	anal.	mp, °C	purific ^a	% yield	UV; λ _{max} (ε)
2	C ₆ H ₅	C ₁₆ H ₁₇ N ₅ O ₅	C, H, N	124	A	22	267 (13 200), 276 sh (9 600)
3	C ₆ H ₅ CH ₂	C ₁₇ H ₁₉ N ₅ O ₅	C, H, N	172-175	A	42	252 sh (9 900), 267 (12 600)
4	C ₆ H ₅ (CH ₂) ₂	C ₁₈ H ₂₁ N ₅ O ₅	C, H, N	95-97	B, 60	65	254 sh (8 000), 267 (10 000)
5	C ₆ H ₅ (CH ₂) ₃	C ₁₉ H ₂₃ N ₅ O ₅	C, H, N	100-102	C, 50/60	20	254 sh (9 300), 268 (11 700)
6	C ₆ H ₅ (CH ₂) ₄	C ₂₀ H ₂₅ N ₅ O ₅	C, H, N	93-96	C, 50/60	23	254 sh (9 600), 268 (12 000)
7	C ₆ H ₅ (CH ₂) ₅	C ₂₁ H ₂₇ N ₅ O ₅	C, H, N	102-104	C, 60/70	19	254 sh (10 000), 268 (12 700)
8	2-FC ₆ H ₅ (CH ₂) ₂	C ₁₈ H ₂₀ FN ₅ O ₅	C, H, N, F	124-126	C, 30/50	29	257 sh (10 000), 267 (12 300)
9	3-FC ₆ H ₄ (CH ₂) ₂	C ₁₈ H ₂₀ FN ₅ O ₅	C, H, N, F	149-154	C, 50/80	6	254 sh (10 200), 267 (12 600)
10	4-FC ₆ H ₄ (CH ₂) ₂	C ₁₈ H ₂₀ FN ₅ O ₅	C, H, N, F	148-150	C, 50/70	36	253 sh (11 300), 267 (14 600)
11	2-ClC ₆ H ₄ (CH ₂) ₂	C ₁₈ H ₂₀ ClN ₅ O ₅	C, H, N, Cl	98-100	C, 45/65	34	255 sh (10 000), 268 (12 700)
12	3-ClC ₆ H ₄ (CH ₂) ₂	C ₁₈ H ₂₀ ClN ₅ O ₅	C, H, N, Cl	118-120	D	25	255 sh (9 200), 268 (11 600)
13	4-ClC ₆ H ₄ (CH ₂) ₂	C ₁₈ H ₂₀ ClN ₅ O ₅	C, H, N, Cl	159	D	16	254 sh (9 400), 267 (12 000)
14	2-CH ₃ OC ₆ H ₄ (CH ₂) ₂	C ₁₉ H ₂₃ N ₅ O ₆	C, H, N	126-130	C, 40/70	38	255 sh (11 300), 269 (15 100)
15	3-CH ₃ OC ₆ H ₄ (CH ₂) ₂	C ₁₉ H ₂₃ N ₅ O ₆	C, H, N	103-105	C, 40/65	37	255 sh (11 100), 269 (15 000)
16	4-CH ₃ OC ₆ H ₄ (CH ₂) ₂	C ₁₉ H ₂₃ N ₅ O ₆	C, H, N	145-146	C, 40/70	48	255 sh (11 200), 268 (14 800)
17	2-CH ₃ C ₆ H ₄ (CH ₂) ₂	C ₁₉ H ₂₃ N ₅ O ₅	C, H, N	166-168	C, 40/70	20	254 sh (10 900), 268 (13 800)
18	3-CH ₃ C ₆ H ₄ (CH ₂) ₂	C ₁₉ H ₂₃ N ₅ O ₅	C, H, N	111-113	C, 45/65	30	255 sh (9 400), 268 (11 800)
19	4-CH ₃ C ₆ H ₄ (CH ₂) ₂	C ₁₉ H ₂₃ N ₅ O ₅	C, H, N	97-100	C, 50/70	38	256 sh (8 600), 267 (9 700)
20	2-thienyl-(CH ₂) ₂	C ₁₆ H ₁₉ N ₅ O ₅ S	C, H, N, S	104-106	C, 35/50	23	236 sh (12 600), 267 (11 300)
21	3-thienyl-(CH ₂) ₂	C ₁₆ H ₁₉ N ₅ O ₅ S	C, H, N, S	99-102	C, 35/50	22	241 sh (11 400), 268 (12 100)
22	3-indolyl-(CH ₂) ₂	C ₂₀ H ₂₂ N ₆ O ₅	C, H, N	138-140	C, 40/70	37	255 sh (11 000), 268 (14 500)
23	1-naphthyl-(CH ₂) ₂	C ₂₂ H ₂₃ N ₅ O ₅	C, H, N	125-130	C, 50/80	7	271 (15 200), 291 sh (4 500)
24	2-naphthyl-(CH ₂) ₂	C ₂₂ H ₂₃ N ₅ O ₅	C, H, N	104-108	B, 40	30	254 sh (11 000), 268 (14 500)
25	3,4-(CH ₃ O) ₂ C ₆ H ₃ (CH ₂) ₂	C ₂₀ H ₂₅ N ₅ O ₇	C, H, N	202-204	C, 30/50	37	255 sh (8 800), 269 (12 200)
26	3,4,5-(CH ₃ O) ₃ C ₆ H ₂ (CH ₂) ₂	C ₂₁ H ₂₇ N ₅ O ₈	C, H, N	110-112	C, 30/50	5	255 sh (9 200), 269 (11 900)

^a Purification methods included A, flash chromatography on silica gel eluted with CH₃OH/CHCl₃, 15:85 v/v; B, reverse-phase HPLC, isocratic elution with indicated percent of CH₃OH in H₂O; C, reverse-phase HPLC, gradient elution at initial/final percent of CH₃OH in H₂O; and D, crystallization from CH₃OH-H₂O.




Figure 1. Diagrammatic representation of the C-2 region of adenosine receptors.

salt of 2-indol-3-ylethanol reacted satisfactorily. The thienyl ethoxides, too, gave useful yields of product.

Table I lists the properties of novel analogues.

Cardiovascular Activity. In the assay system used here,¹ prolongation of the stimulus-QRS interval of the electrocardiogram of a paced guinea pig heart reflects potency at the atrioventricular node⁸ and coronary vasodilation potency at the A₂AR in coronary artery,⁹ respectively. Table II lists the results of such assays and also measurements of *k'*, a hydrophobicity index. All the analogues appear to be full agonists at both receptors, causing second degree heart block and maximum coronary vasodilation.

Analogues 2-7, 2-phenoxyadenosine and its lower alkyl 2-phenylalkoxy congeners, probe the importance of the length of the alkyl chain on activity. At the A₁AR the activities of 2-7 are low, the EC₅₀s of stimulus-QRS interval prolongation ranging between 4.6 and 47 μM. Activity is unrelated to alkyl chain length. At the A₂AR

mediating coronary vasodilation, the activity of 2 is low; the EC₅₀ is 0.42 μM, 8 times higher than that of adenosine and 23 times higher than that of 2-(phenylamino)adenosine (CV-1808), its nitrogen isostere.¹ Such a result is baffling. The coronary vasoactivity of 2-oxoadenosine (isoguanosine) is 3 times higher than that of 2-aminoadenosine, yet the potency ranking of the phenyl derivatives is reversed. The activity of 2-(benzyloxy)adenosine (3) is the same as that of 2, but further increasing the length of the alkyl chain by one more methylene residue greatly increases activity. Phenethoxy analogue 4 is the most potent member of this series; the EC₅₀ of coronary vasoactivity is 2.8 nM, an increase in potency of nearly 150 times over that of 2 or 3. The EC₅₀s of 5-7 are somewhat lower than that of 3 but still in the low nanomolar range.

Analogues 8-19 explore the effects of ring substituents on the activity of 3. The substituents are fluorine (8-10), chlorine (11-13), methoxy (14-16), and methyl (17-19). At the A₁AR, ring substitution has a negligible effect on activity, the EC₅₀s of stimulus-QRS interval prolongation differing from that of 4 by 2.5-fold or less. At the A₂AR, by contrast, the ring position and hydrophobicity of a substituent strongly influences activity. With each kind of substituent, activity increases ortho < meta < para. At each ring position a methoxy group contributes least and a methyl group the most to activity; among the para substituted analogues the activity ranking, OCH₃ < F < Cl < CH₃, parallels the hydrophobicity of each substituent as reflected in the hydrophobicity index, *k'*. Thus, hydrophobicity appears to explain the potency ranking of the para substituted 2-phenethoxyadenosines. Analogues 10, 13, and 19 are active in the subnanomolar range. The most active of the four, 2-[2-(4-methylphenyl)ethoxy]adenosine (19), has an EC₅₀ for coronary vasodilation of 190 pM. Owing to the relatively constant activity of 8-19 at the

(8) Clemo, H. F.; Belardinelli, L. *Circ. Res.* 1986, 59, 427-436.

(9) Kusachi, S.; Thompson, R. D.; Olsson, R. A. *J. Pharmacol. Exp. Ther.* 1983, 227, 316-321.

Table II. Cardiac and Coronary Activity of 2-Aralkoxyadenosines

no.	substit	N ^a	-log EC ₅₀ , M		A ₁ /A ₂ ^d	k'
			stim-QRS ^b	coronary ^c		
	adenosine	14	5.47 ± 0.07	7.29 ± 0.06	80 ± 11	
			C ₆ H ₅ (CH ₂) _n			
	R =					
2	C ₆ H ₅	4	4.33 ± 0.09	6.38 ± 0.06	130 ± 40	0.31
3	C ₆ H ₅ CH ₂	4	5.20 ± 0.06	6.38 ± 0.09	17 ± 4.3	0.61
4	C ₆ H ₅ (CH ₂) ₂	6	4.70 ± 0.09	8.55 ± 0.04	8200 ± 2200	0.90
5	C ₆ H ₅ (CH ₂) ₃	4	4.70 ± 0.04	7.21 ± 0.08	330 ± 33	1.36
6	C ₆ H ₅ (CH ₂) ₄	4	5.15 ± 0.03	8.02 ± 0.04	760 ± 85	2.20
7	C ₆ H ₅ (CH ₂) ₅	4	5.34 ± 0.07	8.20 ± 0.05	730 ± 61	3.63
			R = XC ₆ H ₄ (CH ₂) ₂			
	X =					
8	2-F	4	4.45 ± 0.05	8.58 ± 0.03	14000 ± 2200	0.94
9	3-F	4	4.46 ± 0.09	8.72 ± 0.10	18000 ± 2300	0.83
10	4-F	6	4.59 ± 0.10	9.06 ± 0.08	38000 ± 12000	0.86
11	2-Cl	4	4.66 ± 0.03	8.26 ± 0.07	4000 ± 500	1.55
12	3-Cl	4	4.79 ± 0.10	8.52 ± 0.07	6400 ± 1600	1.63
13	4-Cl	4	4.84 ± 0.04	9.41 ± 0.04	39000 ± 6500	1.64
14	2-CH ₃ O	4	4.45 ± 0.03	7.50 ± 0.05	1100 ± 150	1.04
15	3-CH ₃ O	4	4.79 ± 0.10	8.59 ± 0.10	8200 ± 3300	0.87
16	4-CH ₃ O	4	4.71 ± 0.08	8.85 ± 0.06	14000 ± 1300	0.82
17	2-CH ₃	4	4.60 ± 0.11	8.42 ± 0.14	9100 ± 4400	1.39
18	3-CH ₃	4	4.89 ± 0.07	8.80 ± 0.06	8400 ± 1200	1.48
19	4-CH ₃	4	5.13 ± 0.04	9.72 ± 0.11	44000 ± 9700	1.56
			R = Ar(CH ₂) ₂			
	Ar =					
20	2-thienyl	4	4.93 ± 0.10	8.43 ± 0.03	3500 ± 750	0.67
21	3-thienyl	4	4.73 ± 0.06	8.47 ± 0.06	5700 ± 700	0.67
22	3-indolyl	4	4.85 ± 0.09	8.01 ± 0.03	1600 ± 480	0.68
23	1-naphthyl	4	5.08 ± 0.05	8.30 ± 0.05	1700 ± 290	2.45
24	2-naphthyl	4	4.94 ± 0.07	9.30 ± 0.05	23000 ± 3200	2.45
25	3,4-(CH ₃ O) ₂ C ₆ H ₃	5	4.37 ± 0.02	8.31 ± 0.10	7400 ± 1200	0.53
26	3,4,5-(CH ₃ O) ₃ C ₆ H ₂	4	4.33 ± 0.08	7.66 ± 0.06	2300 ± 520	0.51

^a Number of assays. ^b Prolongation of the stimulus-QRS interval, an index of agonist activity at the A₁AR. ^c Coronary vasodilation, an index of agonist activity at the A₂AR. ^d Selectivity ratio, stim-QRS + coronary. ^e In the series of analogues 2-7, *n* increases consecutively 0, 1, 2...5.

A₁AR, the A₁/A₂ activity ratio varies directly with activity at the A₂AR. The A₁/A₂ activity ratio of 19 is 44 000, that of 10 is 38 000, that of 13 is 39 000, and that of 16 is 14 000. The potencies and selectivities of these analogues compare favorably with those of *N*-ethyl-2-[[2-[4-(2-carboxyethyl)phenyl]ethyl]amino]adenosine-5'-uronamide (CGS 21,680), currently the standard A₂AR agonist.¹⁰ In the same guinea pig heart Langendorff preparation as used in the present study, CGS 21,680 exhibits an EC₅₀ for coronary vasodilation of 0.74 ± 0.08 nM and an A₁/A₂ activity ratio of 33 000 ± 2300.¹

Analogues 20-24 are congeners of 4 that contain aryl groups other than benzene. All are very poor A₁AR agonists. As A₂AR agonists the two thiophene congeners, 20 and 21, are essentially equipotent with 4 and the larger but more polar 3-indolylethyl analogue 22 is only 3 times less potent. The rather high vasoactivity of 24 is evidence that the hydrophobic subregion is substantially larger than a cyclohexane or benzene ring. The two naphthyl congeners, 23 and 24, differ in A₂AR agonist activity by a factor of 10. The more potent 2-naphthyl isomer, 24, has an EC₅₀ for coronary vasoactivity of 0.50 nM. Studies with molecular models suggest that bulk tolerance might be

somewhat limited in the part of the receptor occupied by ortho and, to a lesser extent, meta substituents on the benzene ring of 4.

3,4-Dimethoxyphenyl and 3,4,5-trimethoxyphenyl analogues 25 and 26 together with monomethoxy derivative 16 explore the effects of polarity and steric bulk on activity at the A₂AR. The potency ranking 16 > 25 > 26 parallels the hydrophobicity indices. Studies of molecular models show that it is possible to superimpose the methoxy groups of 25 on the distal ring of the naphthyl group of 23. Accordingly, hydrophobicity, rather than steric hindrance, seems a likely explanation for the difference in activity between 16 and 26. It is possible that the second meta substituent of 26 could be sterically hindering, but the rather mild loss of activity suggests that the greater polarity of this analogue is probably a better explanation.

Antagonism by Alkylxanthines

Antagonism by alkylxanthines such as theophylline¹¹ is one criterion used to show that adenosine acts through adenosine receptors. The unselective antagonist 8-(*p*-sulfophenyl)theophylline¹² (8-PST) at a dose of 50 μM competitively antagonized the activity of 4 and 10 at both

(10) Watson, S.; Abbott, A. *Trends Pharmacol. Sci.* 1989, Suppl., 3.

(11) Sattin, A.; Rall, T. W. *Mol. Pharmacol.* 1970, 6, 13-23.

(12) Daly, J. W.; Padgett, W.; Shamim, M. T.; Butts-Lamb, P.; Waters, J. J. *Med. Chem.* 1985, 28, 487-492.




Figure 2. Schild plots characterizing the antagonism by 8-(*p*-sulphophenyl)theophylline of the actions of 19 at the A_1 AR of AV node (above) and at the A_2 AR of coronary artery (below). Abbreviations: dr, dose ratio, the quotient of the EC_{50} of agonist in the presence of antagonist divided by the EC_{50} in the absence of antagonist; Stim-QRS, stimulus-QRS interval, an index of activity at the A_1 AR of AV node; coronary flow, an index of activity at the A_2 AR in coronary artery.

receptors. At the A_1 AR, 8-PST shifted the dose-response curves of 4 and 10 rightward 3.2- and 3.1-fold, respectively; at the A_2 AR the corresponding rightward shifts were 225- and 40-fold. In addition to these studies at a single concentration of agonist, we estimated the pK_B of 8-PST vs 19 (Figure 2). Such estimates of pK_B , 5.27 at the A_1 AR and 6.45 at the A_2 AR, are similar to estimates in the literature.^{6,13}

Receptor Models. The uniformly low activity of the 2-aralkoxyadenosines at the A_1 AR and the lack of correlations between structure and activity support a major conclusion reached by analysis of the SAR of the 2-aralkoxyadenosines, namely, that the A_1 AR lacks a hydrophobic subregion or, alternatively, that the interaction of 2-substituents with this subregion does not contribute to activity.¹ That study also showed that the 2-isoalkoxyadenosines and 2-(cyclohexylalkoxy)adenosines exhibit A_1 AR activity that parallels the number of methylene residues in the alkyl chain. It was pointed out that such a result was not necessarily evidence for interaction with the C-2 region. Instead, it could reflect the interaction of such long-chain alkyl groups with the N-6 region of the A_1 AR. Indeed, branched-chain and cycloalkyl substituents at N-6 greatly enhance the potency of adenosine at the A_1 AR^{14,15} but aromatic groups in such substituents will lower activity.^{14,16,17} The A_1 AR potency of 2-7 did not

correlate with alkyl chain length, possibly a reflection of the negative influence of the phenyl group on interactions of these C-2 substituents with the N-6 region.

The activities of the 2-aralkoxyadenosines at the A_2 AR confirm and refine the model of the A_2 AR developed from the SAR of the 2-alkoxyadenosines. At the coronary receptor, the activities of the para-substituted 2-phenethoxyadenosines are inversely proportional to the polarity of the nucleosides, a result consistent with the existence of a hydrophobic subregion in the coronary A_2 AR. The high activity of 2-[2-(2-naphthyl)ethoxy]adenosine (24) is evidence that the hydrophobic subregion is larger than indicated by the previously reported¹ activity of 2-[2-(cyclohexyl)ethoxy]adenosine. Additionally, the 10-fold difference in the activities of the 1- and 2-naphthyl isomers suggests that there may be local areas of limited bulk tolerance within the hydrophobic subregion. The low activity of 2-(benzyloxy)adenosine supports the notion that the alkyl region adversely affects activity.

In summary, this study of the agonist activities of 2-aralkoxyadenosines confirms the provisional models of the C-2 regions of the A_1 AR and A_2 AR described previously. Several of these analogues are very potent and highly selective A_2 AR agonists.

Experimental Section

Flash chromatography on 60- μ m silica gel eluted with 2% CH_3OH in $CHCl_3$ purified the blocked nucleosides. Deblocking entailed boiling in a solution of 2% formic acid and 50% acetic acid with close monitoring by HPLC. The purification of 2-26 employed a Rainin Autoprep fitted with a 1 \times 25 cm column of C-18 silica, eluted with CH_3OH -water in either the isocratic or gradient mode. Melting points are uncorrected. A Varian EM 360L spectrometer yielded proton NMR spectra of nucleoside solutions in $DMSO-d_6$, which were consistent with the assigned structures. The ultraviolet spectra were determined on a Beckman DU64 spectrophotometer. MHW Laboratories, Tucson, AZ, performed the elemental analysis, which differed from the calculated composition by <0.4%. Product accounted for >99% of the UV-absorbing material in samples submitted for bioassay. The retention time of a nucleoside on a reverse-phase HPLC column served for the calculation¹⁸ of a hydrophobicity index, k' , by the formula $k' = (t - t_0)/t_0$, where t is the retention time of the solute and t_0 is the transit time of the solvent. In the present experiments the mobile phase consisted of a mixture containing 35% 10 mM $NaHPO_4$, pH 7.0, and 65% CH_3OH .

Preparation of 2-Arylalkoxyadenosines from Primary Alcohols. 2-[2-(4-Methylphenyl)ethoxy]adenosine [6-Amino-2-[2-(4-methylphenyl)ethoxy]-9- β -D-ribofuranosyl-9H-purine]. A solution of 2-(4-methylphenyl)ethanol (4.6 g, 33.6 mmol) in 70 mL of dry 1,2-dimethoxyethane was cooled to 10 $^\circ$ C in an ice bath. To this solution was added 1.6 M *n*-butyllithium (19.9 mL, 31.9 mmol), the solution was stirred for 15 min, and then dry 2-chloro-2',3'-*O*-(ethoxymethylidene)adenosine (3.0 g, 8.4 mmol) was added in one portion. After 5 days at reflux, HPLC showed that <5% of the starting material was present. The solvents were removed in vacuo and a solution of the residue in 70 mL water was extracted with ethyl acetate (4 \times 50 mL). The combined extracts were dried over $MgSO_4$ and evaporated in vacuo to a dark syrup for purification by flash chromatography. Fractions containing product were concentrated and dissolved in acetic acid (25 mL), water (25 mL), and 98% formic acid (1 mL). After boiling of the solution for 2 h, HPLC showed that the blocked nucleoside had disappeared. Solid sodium bicarbonate was carefully added until the pH was basic. Removing the solvents in vacuo and the purifying product by preparative reverse-phase HPLC, as described in Table I, yielded 1.23 g (38%) of a white solid.

(13) Jacobson, K. A.; De La Cruz, R.; Schulick, R.; Kiriasis, L.; Padgett, W.; Pfliederer, W.; Kirk, K. L.; Neumeyer, J. L.; Daly, J. W. *Biochem. Pharmacol.* **1988**, *37*, 3653-3661.

(14) Daly, J. W.; Padgett, W.; Thompson, R. D.; Kusachi, S.; Bugni, W. J.; Olsson, R. A. *Biochem. Pharmacol.* **1986**, *35*, 2467-2481.

(15) Moos, W. H.; Szotek, D. S.; Bruns, R. F. *J. Med. Chem.* **1985**, *28*, 1383-1384.

(16) Paton, D. M.; Olsson, R. A.; Thompson, R. D. *Naunyn Schmiedeberg Arch. Pharmacol.* **1986**, *333*, 313-322.

(17) Dunwiddie, T. V.; Worth, T. S.; Olsson, R. A. *Naunyn Schmiedeberg Arch. Pharmacol.* **1986**, *334*, 77-85.

(18) Brent, D. A.; Sabathka, J. J.; Minick, D. J.; Henry, D. W. *J. Med. Chem.* **1983**, *26*, 1014-1020.

Bioassay. A Langendorff guinea pig heart preparation paced at 260 beats/min via the left atrium served for assays of A₁AR and A₂AR agonist activity. The perfusion buffer consisted of (mM) NaCl (120), NaHCO₃ (27), KCl (3.7), KH₂PO (1.3), MgSO₄ (0.64), CaCl₂ (1.3), pyruvate (2), and glucose (5). The buffer was saturated with 95% O₂-5% CO₂, equilibrated at 37 °C in a heat exchanger and delivered at a pressure equivalent to 55 mmHg. Continuous drainage of the left ventricle by means of a catheter inserted across the mitral valve insured that this cardiac chamber did no external work. An electrode in the right ventricle monitored the electrocardiogram. Time collections of cardiac effluent in a graduated cylinder during the steady-state phase of the flow responses to analogue administration measured total coronary flow, which was also monitored by an in-line electromagnetic flowmeter in the aortic perfusion cannula. The rate of the nucleoside infusion was increased stepwise until the appearance of second degree heart block. The quotient of the ratio of nucleoside infusion (mol/min) divided by coronary flow rate (L/min) equals agonist concentration in the perfusate. The EC₅₀ of prolongation of the stimulus-QRS interval, the concentration of agonist needed

to prolong the interval by 50% of the maximum response,⁸ reflects activity at the A₁AR. Logit transformation of the coronary flow data and solution of the regression of logit (coronary flow) on log [analogue] for logit = 0 yielded an estimate of EC₅₀ of coronary vasodilation, an index of A₂AR activity. Table II reports the mean ± SEM of the -log EC₅₀ values from assays in four or more hearts. The quotient of the EC₅₀ of stimulus-QRS prolongation divided by the EC₅₀ of coronary vasodilation provided an index of selectivity. Values of the index <1 indicate selectivity for the A₁AR and values >1 selectivity for the A₂AR. Table II reports the mean ± SEM of the A₁/A₂ activity ratios of individual experiments.

Registry No. 1 (R' = Me₂), 24639-06-3; 1 (R' = OEt), 56720-43-5; 2, 50257-82-4; 3, 131865-78-6; 4, 131865-79-7; 5, 131865-80-0; 6, 131865-81-1; 7, 131865-82-2; 8, 131865-83-3; 9, 131865-84-4; 10, 131865-85-5; 11, 131865-86-6; 12, 131865-87-7; 13, 131865-88-8; 14, 131865-89-9; 15, 131865-90-2; 16, 131865-91-3; 17, 131865-92-4; 18, 131865-93-5; 19, 131865-94-6; 20, 131865-95-7; 21, 131865-96-8; 22, 131865-97-9; 23, 131865-98-0; 24, 131865-99-1; 25, 131866-00-7; 26, 131866-01-8.

Inhibition of Human Placental Aromatase by Novel Homologated 19-Oxiranyl and 19-Thiiranyl Steroids


Wayne E. Childers,[†] James V. Silverton,[‡] James T. Kellis, Jr.,[§] Larry E. Vickery,[§] and Cecil H. Robinson*[†]

Department of Pharmacology and Molecular Sciences, The Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, Laboratory of Chemistry, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland 20892, and The Department of Physiology and Biophysics, University of California, Irvine, California 92717. Received August 18, 1989

Novel homologated 19-oxiranyl- and 9-thiiranyl-androst-4-ene-3,17-diones (8a,b and 9a,b, respectively) have been synthesized. The configuration and conformation of compound 8a have been established by X-ray crystallographic analysis. All four compounds have been shown to be competitive inhibitors of human placental aromatase. The thiiranes were more potent inhibitors than the corresponding oxiranes, and the 2'S isomers (8b and 9b) were better inhibitors than the 2'R (8a and 9a) diastereomers in each series. Spectroscopic studies with purified human placental aromatase suggest that the oxiranyl oxygen and thiiranyl sulfur of 2'S compounds 8b and 9b coordinate to the enzyme's heme iron.


Aromatase is a cytochrome P-450 enzyme complex which is responsible for the important transformation of androgens (1) to estrogens (2) (Scheme I). Inhibitors of aromatase may be valuable as therapeutic agents in the treatment of estrogen-sensitive breast tumors and as possible antifertility agents, and a number of competitive inhibitors has been reported in recent years.^{1,2}

Previous work from our laboratories showed that 10β-oxiranyl- and 10β-thiiranylestro-4-ene-3,17-diones (3 and 4, respectively) were potent competitive inhibitors of human placental aromatase.³⁻⁵ Furthermore, stereoselectivity was observed, with the 19R diastereomers (3a, 4a)



being 10-75 times more potent inhibitors than the 19S isomers (3b, 4b). Spectroscopic studies with purified enzyme demonstrated that in the case of the 19R isomers, the oxiranyl and thiiranyl heteroatom coordinated with the enzyme's heme iron. This coordination, combined with

Scheme I



the inherent binding selectivity of the steroid nucleus, endowed these inhibitors with high specificity.

X-ray crystallographic analysis of the above 10β-oxiranes showed³ that the oxirane heteroatom in the more potent

[†]The Johns Hopkins University School of Medicine.

[‡]National Institutes of Health.

[§]University of California.

- (1) Johnston, J. O.; Metcalf, B. W. *Novel Approaches to Cancer Therapy*; Sunkara, P., Ed.; Academic Press: New York, 1984; Chapter 9, p 307.
- (2) Brodie, A. M. H. *Biochem. Pharmacol.* 1985, 34, 3213.
- (3) Shih, M.-J.; Carrell, M. L.; Carrell, H. L.; Wright, C. L.; Johnston, J. O.; Robinson, C. H. *J. Chem. Soc. Chem. Commun.* 1987, 213.
- (4) Childers, W. E.; Robinson, C. H. *J. Chem. Soc. Chem. Commun.* 1987, 320.
- (5) Kellis, J. T.; Childers, W. E.; Robinson, C. H.; Vickery, L. E. *J. Biol. Chem.* 1987, 262, 4421.