

Preliminary communication

Synthesis of organocobaloximes: modification of the organic group in organocobaloximes

B.D. Gupta^{*}, Vandana Dixit

Department of Chemistry, I.I.T., Kanpur 208016, U.P., India

Received 22 July 1996; revised 26 September 1996

Abstract

5-Substituted heteroaromaticmethyl cobaloximes are synthesized by a very simple reaction of arenesulfonyl chloride, ArSOCl ($\text{Ar} = \text{Ph}$, C_6Cl_5 and $2,4\text{-(NO}_2)_2\text{C}_6\text{H}_3$) with heteroaromaticmethyl cobaloximes, $\text{RCo}^{\text{III}}(\text{dmgH})_2\text{Py}$ [$\text{R} = 2\text{- and } 3\text{-thienylmethyl, furfuryl}$] in the dark at 0°C . As organocobaloximes are susceptible to Co–C bond cleavage by the electrophiles and free radicals, the ring substitution is quite remarkable.

Keywords: Organocobaloxime; Organocobalt

1. Introduction

The initial studies on organocobaloximes, $\text{RCo}(\text{dmgH})_2\text{Py}$, were focused largely on its role as a model for vitamin B_{12} coenzyme [1–10]. Recent work, however, has shown that its chemistry is so interesting that it has acquired an independent research field [11–16]. These have been used as templates in many organic syntheses [17] and as catalysts in chemical processes [18]. The key feature of these complexes is the weak Co–C bond and, since small structural changes in the molecule have a profound effect on the reactivity of the Co–C bond [9], the synthesis of new complexes continues to interest chemists. Many synthetic routes have been reported and many new ones are still appearing in the literature [10]. Generally, one starts with a Co(I) , Co(II) or Co(III) complex to synthesize organocobaloximes. The use of organocobaloximes as precursors for the synthesis of new organocobaloximes, however, does not find an adequate place in the literature. The known examples, though few, have generally employed simple reactions like hydrolysis (both acidic

as well as basic) [19,20], acetal hydrolysis [21], cyclisation reactions [22–24], transalkylations [25], $\sigma\text{--}\pi$ rearrangements [26–28], cycloaddition [29,30] and Diels–Alder reaction [31]. Solid state photoisomerisation of optically active alkyl cobaloximes has also been reported [32].

In this paper we report the synthesis of 5-substituted heteroaromaticmethyl cobaloximes by a simple reaction of the arene sulfonyl chloride, ArSOCl ($\text{Ar} = \text{Ph}$, C_6Cl_5 and $2,4\text{-(NO}_2)_2\text{C}_6\text{H}_3$) with heteroaromaticmethyl cobaloximes.

2. Experimental

The synthesis of organocobaloximes (1–3) has been reported earlier [15]. Benzene sulfonyl chloride (A) was prepared [33] by the chlorination of diphenyldisulfide with sulfuryl chloride at ambient temperature (b.p. $46\text{--}48^\circ\text{C}/4\text{ mm}$, lit. $49^\circ\text{C}/2\text{ mm}$, $^1\text{H NMR } \delta$: (CCl_4) 7.4 (s, Ar). Pentachlorobenzene sulfonyl chloride (B) was prepared [34] by the chlorination of pentachlorobenzene thiol with chlorine gas (m.p. 100°C , lit. 104°C). The thiol was prepared [35] from hexachlorobenzene, sodium sulfide and sulfur under refluxing conditions. 2,4-Dinitrobenzene sulfonyl chloride (C) was bought from Fluka and used as such.

^{*} Corresponding author.

3. Results and discussion

Organocobaloximes (1–3) react with arene sulfonyl chloride, ArSCI (A, B or C) [see Scheme 1] in 1:1 molar ratio in the dark (reaction flask covered with aluminum foil) at 0 °C in dichloromethane. The progress of the reaction is monitored by TLC on silica gel using ethyl acetate. The reaction is generally complete within 3 h. The mixture is then concentrated and poured into solvent ether. The precipitated solid is washed with ether and is further purified by column chromatography on silica gel using a mixture of dichloromethane and ethyl acetate. The combined ether extract after evaporation does not afford any characterizable organic product. Similar reactions when carried out under photochemical conditions (irradiation by 2 × 200 W tungsten lamps) at 0 °C give a complicated mixture of products, both organic as well as organometallic, in each reaction (Table 1).

The exclusive formation of the ring-substituted organocobaloximes suggests that we are dealing with a unique class of sigma bonded organo cobalt complexes

having an activated arene ring which is susceptible to attack by the electrophilic ArS⁺ species. This is justified in view of our earlier observations that the methylomethyl group –CH₂Co(dmgH)₂Py is highly activating, and its activation is even more than that of the methoxy group [16]. As organocobaloximes are susceptible to Co–C cleavage by the electrophiles [16,36–39] and free radicals [11–16], the substitution into the aromatic ring is quite remarkable. This is a rare observation in organocobaloxime chemistry.

One-electron oxidation potential values of the ring-substituted products are comparable to those of the parent cobaloximes and are irreversible in nature.


The ring-substituted organocobaloximes undergo very facile oxygen insertion into the Co–C bond under photochemical conditions (*hν*, 0 °C/2 h) (Scheme 1, footnote).

We also observe a remarkable difference in the reactivity of 3- and 4-methoxybenzyl cobaloximes [15] with arene sulfonyl chloride. For example, the cleavage of the Co–C bond occurs in the 4-OMe case with 4-OMeC₆H₄CH₂SPh as one of the many products: ¹H

$\text{RCo(dmgH)}_2\text{Py} + \text{ArSCI} \xrightarrow[1-3\text{h}/\text{CH}_2\text{Cl}_2]{\text{dark}/0^\circ\text{C}} \text{R}'\text{S-Co(dmgH)}_2\text{Py}$			
R = ^a RCo ^(III)	ArSCI (ZCl) Z =	Product (Product no)	Yield (%)
 (1)	A, B, C	Z = 	
		Z = A (4)	64
		B (5)	60
		C (6)	55
 (2)	A, B, C	Z = 	
		Z = A (7)	50
		B (8) [§]	70
		C (9)	28#
 (3)	A, B, C	Z = 	
		Z = A (10)	52
		B (11)	58
		C (12)	27#

A = PhS-, B = C₆H₄S-, C = 2,4 (NO₂)₂ C₆H₃S-

Incomplete even after 5h, ^a (3) on reaction with Cl₂ or Br₂ at 0 °C in dark forms the 5-halo substituted organocobaloximes whereas (1) and (2) lead to the cleavage of the Co–C bond [15]. [§] (8) with molecular oxygen under photochemical conditions forms



Scheme 1.


Table 1

Characteristics of the organocobaloximes (1–3) and ring-substituted products (4–12)

Product No.	¹ H NMR (chemical shifts) δ				UV-vis $\lambda_{\max}$ (nm) (CH ₂ OH)	CV values ^a $E_{1/2}$ vs. SCE (V)	Anal. Found (Calc.) (%) ^b			
	dmgH	CH ₂	Heteroaromatic	Py/aromatic			C	H	N	S
1	2.05 (s)	3.00 (s)	6.65, 7.00 (dd)	7.20, 7.65, 8.50	385, 281, 240	0.950	c			
2	2.00 (s)	2.40 (s)	6.00, 7.40 (dd)	7.30, 7.75, 8.60	383, 284, 239	0.931	c			
3	2.00 (s)	2.85 (s)	6.65, 7.20 (dd)	7.30, 7.70, 8.50	359, 277, 239	0.950	c			
4	2.10 (s)	2.06 (s)	6.02–6.40 (dd)	7.20, 7.60, 8.40	379, 240, 208	0.825	50.12 (50.26)	4.82 (4.88)	12.18 (12.21)	11.10 (11.16)
5	2.00 (s)	2.50 (s)	5.96–6.40 (dd)	7.20, 7.50, 8.35	382, 217	0.834	38.58 (38.63)	3.01 (3.08)	9.32 (9.38)	8.55 (8.58)
6	2.06 (s)	2.50 (s)	5.95–6.50 (dd)	7.20, 7.60, 8.40 7.66 (s), 7.40 (s) 8.20 (d), 8.90 (d) 9.00 (d)	299, 239, 207	0.915	43.35 (43.43)	3.85 (3.92)	14.68 (14.78)	9.60 (9.65)
7	1.96 (s)	2.75 (s)	6.50–6.80 (dd)	7.20, 7.40, 8.35 7.04–7.10 (m)	380, 286, 243 208	0.899	51.62 (51.70)	5.91 (5.02)	12.08 (12.13)	5.68 (5.74)
8	1.90 (s)	2.80 (s)	6.60–6.90 (dd)	7.20, 7.60, 8.40	385, 290, 219	0.846	39.41 (39.47)	3.14 (3.15)	7.50 (7.59)	4.32 (4.38)
9	2.18 (s)	3.00 (s)	6.80–7.08 (dd)	7.34, 7.80, 8.52 7.60 (s), 7.40 (s) 8.30 (d), 8.35 (d) 9.10 (d)	308, 241, 206	1.071	44.82 (44.85)	3.95 (4.04)	15.10 (15.26)	4.88 (4.98)
10	1.96 (s)	2.76 (s)	6.70–6.80 (dd)	7.20, 7.50, 8.40 7.20 (m)	367, 244, 209	0.884	50.28 (50.26)	4.80 (4.88)	12.18 (12.21)	11.10 (11.16)
11	2.01 (s)	2.95 (s)	6.65–6.85 (dd)	7.20, 7.50, 7.85	380, 237, 218	0.958	38.59 (38.63)	3.02 (3.08)	9.35 (9.38)	8.50 (8.58)
12	2.01 (s)	2.56 (s)	6.7–7.0 (m)	7.60 (s), 7.40 (s) 7.70 (m), 8.10 (d) 8.25 (d), 9.10 (d)	306, 242, 206	1.01	43.30 (43.43)	3.88 (3.92)	14.79 (14.78)	9.64 (9.65)

^a Oxidation potential.^b (5) Cl, 23.72 (23.80); (8) Cl, 24.21 (24.33); (11) Cl, 23.70 (23.80).^c Taken from Ref. [15].

NMR (CDCl₃) δ : [3.86 (s, -CH₂S), 7.10, 6.50–7.00 (dd, Ar), 3.70 (s, OMe-)]; m.p. 75 °C, $\lambda_{\max}$ (MeOH) 265, 231, whereas ring substitution is the facile process in 3-methoxybenzyl cobaloxime.



Yield 40%. ¹H NMR (CDCl₃) δ : 1.98 (s, dmgH), 2.98 (s, -CH₂), 3.72 (s, -OMe), 6.60–7.20 (m, Ar and Py), 7.48, 8.40 (m, Py); $\lambda_{\max}$ (MeOH) 344, 246, 212.

Acknowledgements

We thank the DST, New Delhi, for funding this project.

References

- [1] G.N. Schrauzer and J. Kohnle, *Chem. Ber.*, **97** (1964) 3056.
- [2] G.N. Schrauzer, L.P. Lee and J.W. Sibert, *J. Am. Chem. Soc.*, **92** (1970) 2997.
- [3] G. Costa, G. Mestroni and L. Stefani, *J. Organomet. Chem.*, **7** (1967) 193; G. Costa, G. Mestroni and G. Pellizzer, *J. Organomet. Chem.*, **11** (1968) 333.
- [4] J. Halpern, *Science*, **227** (1985) 869.
- [5] G. Costa, *Coord. Chem. Rev.*, **8** (1972) 63.
- [6] D. Dodd and M.D. Johnson, *J. Organomet. Chem.*, **52** (1973) 1.
- [7] D. Dolphin (ed.), *B₁₂*, Vol. 1, Wiley Interscience, New York, 1982.
- [8] P.J. Toscano and L.G. Marzilli, *Progr. Inorg. Chem.*, **31** (1984) 104.
- [9] N. Bresciani-Pahor, M. Forcolin, L.G. Marzilli, L. Randaccio, M.F. Summers and P.J. Toscano, *Coord. Chem. Rev.*, **63** (1985) 1; L. Randaccio, N. Bresciani-Pahor, E. Zangrando and L.G. Marzilli, *Chem. Rev.*, (1989) 225.
- [10] B.D. Gupta and S. Roy, *Inorg. Chim. Acta*, **146** (1988) 209.
- [11] B. Giese, in *Radicals in Organic Synthesis: Formation of C–C bonds*, Pergamon Press, Oxford, 1986.
- [12] R. Scheffold, G. Rytzel and L. Walder, in S. Scheffold (ed.), *Transition Metals in Organic Synthesis*, Vol. 3, Wiley, Chichester, 1983.
- [13] M.D. Johnson, *Acc. Chem. Res.*, **16** (1983) 343.

- [14] B.D. Gupta, M. Roy, S. Roy, M. Kumar and I. Das, *J. Chem. Soc., Perkin Trans. II*, (1990) 537; S. Roy, I. Das, K. Bhanuprakash and B.D. Gupta, *Tetrahedron*, (1994) 1847.
- [15] B.D. Gupta, M. Kumar and S. Roy, *Inorg. Chem.*, 28 (1989) 11.
- [16] G. Pattenden, *Chem. Soc. Rev.*, 17 (1988) 361.
- [17] E. Langer (ed.), *Methoden der Organische Chemie (Houben-Weyl)*, Vietie Auflage, 13/9b, George Thieme, Stuttgart, 1984, pp. 1–284; A.K. Ghosh and Y. Chen, *Tetrahedron Lett.*, 36 (1995) 869; S. Sukamoto, Y. Kondo and S. Igarashi, *Heterocycles*, 41 (1995) 1771; Y. Takahashi, Y. Honda and T. Tsuchiya, *Carbohydr. Res.*, 270 (1995) 77; B.P. Branchaud, R.M. Slade and S.K. Janisse, *Tetrahedron Lett.*, 34 (1993) 7885.
- [18] A. Rockenbauer, M. Eyor, M. Kwiecinski and S. Tyrlik, *Inorg. Chim. Acta*, 58 (1982) 237; M. Noguchi and S. Kambara, *J. Polym. Sci. B*, (1963) 553; S. Namethi and L. Simandi, *J. Mol. Catal.*, 14 (1982) 87; L.I. Simandi, T.M. Bama, L. Koeze and A. Rockenbauer, *Tetrahedron Lett.*, 34 (1993) 317.
- [19] K.L. Brown, A.W. Awrtrey and R. LeGates, *J. Am. Chem. Soc.*, 100 (1978) 823.
- [20] G.N. Schrauzer and R.J. Windgassen, *J. Am. Chem. Soc.*, 89 (1967) 1999.
- [21] B.T. Golding, T.J. Kemp, C.S. Sell, P.J. Sellers and W.P. Watson, *J. Chem. Soc., Perkin Trans. II*, (1978) 839.
- [22] E.A. Parfenov, T.G. Chervyakova, M.G. Edelev and A.M. Yurkevich, *Zh. Obshch. Khim.*, 44 (1974) 2362.
- [23] P. Bougeard, C.J. Cooksey, M.D. Johnson, M.J. Lewis, S. Mitchell and P.A. Owens, *J. Organomet. Chem.*, 288 (1985) 349.
- [24] E.A. Parfenov and T.G. Chervyakova, *Zh. Obshch. Khim.*, 45 (1975) 1200, 842.
- [25] E.A. Parfenov, T.G. Chervyakova and M.G. Edelev, *Zh. Obshch. Khim.*, 47 (1977) 834.
- [26] M.P. Atkins, B.T. Golding and P.J. Sellers, *Chem. Commun.*, (1978) 954.
- [27] M.P. Atkins, B.T. Golding, A. Bury, M.D. Johnson and P.J. Sellers, *J. Am. Chem. Soc.*, 102 (1980) 3630.
- [28] A. Bury, M.R. Ashcroft and M.D. Johnson, *J. Am. Chem. Soc.*, 100 (1978) 3217.
- [29] D. Dodd, M.D. Johnson, I.P. Steeples and B.L. McKenzie, *J. Am. Chem. Soc.*, 98 (1976) 6399.
- [30] C.J. Cooksey, D. Dodd, M.D. Johnson and B.L. Lockman, *J. Chem. Soc., Dalton Trans.*, (1978) 1814.
- [31] M. Tada and T. Shimizu, *Bull. Chem. Soc. Jpn.*, 65 (1992) 1252.
- [32] Y. Ohgo, Y. Arai and S. Takeuchi, *Chem. Lett.*, (1991) 455; Y. Takenaka, Y. Ohashi, T. Tamura, A. Uchida, Y. Sarada, Y. Ohgo and S. Buba, *Acta Crystallogr.*, B49 (1993) 272; Y. Ohgo, Y. Arai, M. Hagiwara, S. Tekeuchi, H. Kogo, A. Seune, H. Uekusu and Y. Ohashi, *Chem. Lett.*, (1994) 751; Y. Ohashi, T. Nemoto and Y. Takenaka, *Mol. Cryst. Sci. Technol.*, A242 (1994) 371.
- [33] W.H. Mueller and P.E. Butler, *J. Am. Chem. Soc.*, 90 (1965) 2075.
- [34] R.E. Putnam and W.H. Sharply, *J. Am. Chem. Soc.*, 79 (1957) 6526.
- [35] A. Wein, W.N.J. Renner, H.K. Steincker and K. Wedmayer, *Ger. D.E. 314,649*; *Chem. Abstr.*, 102: 456150.
- [36] M.D. Johnson, in F.R. Hartley and S. Patai (eds.), *The Nature and Cleavage of Metal–Carbon Bonds*, Wiley Interscience, New York, 1985.
- [37] K.L. Brown, in D. Dolphin (ed.), *B₁₂*, Vol. 1, Wiley Interscience, New York, 1982, p. 245.
- [38] J. Halpern, in D. Dolphin (ed.), *B₁₂*, Vol. 1, Wiley Interscience, New York, 1982, p. 501.
- [39] M. Rosenblum, *Acc. Chem. Res.*, 7 (1974) 122.