

The Synthesis of Biphenomycin A¹


Ulrich Schmidt,* Volker Leitenberger, Regina Meyer and Helmut Griesser

Institut für Organische Chemie und Isotopenforschung der Universität Stuttgart, Pfaffenwaldring 55, 7000 Stuttgart 80, Germany

Biphenomycin A, a highly potent antibiotic against Gram-positive, β -lactam-resistant bacteria, which was previously isolated from culture filtrates of *Streptomyces griseorubiginosus* No. 43708, has now been synthesized.

Biphenomycins A **1a** and B **1b** are cyclic tripeptides exhibiting high antibiotic activities against Gram-positive, β -lactam-resistant bacteria and have previously been isolated from culture filtrates of *Streptomyces griseorubiginosus* No. 43708 in the ratio of 10 : 1 (**1a** : **1b**).²⁻⁴ These cyclopeptides, together with those of the vancomycin group, are unique in that they


contain a biphenyl structural moiety. We have reported on the total synthesis of biphenomycin B⁵ which contains the two non-proteinogenic amino acids (2*S*,4*R*)-hydroxyornithine and (*S*,*S*)-diisotyrosine. In place of the latter, the main metabolite biphenomycin A contains the corresponding hydroxy derivative **2** with an additional stereogenic centre *i.e.* the biphenyl



Abbreviations: Bn = benzyl; Boc = *tert*-butoxycarbonyl; TMSE = trimethylsilylethyl; Z = benzyloxycarbonyl

Scheme 1 Reagents and conditions: i, 3, Sn(OTf)₂, *N*-ethylpiperidine, tetrahydrofuran (THF), -78 °C, 1.5 h, **4**, 2 h, -78 °C, 75%; ii, MeOMgBr, MeOH, 0 °C, 3 min, 90%; iii, (Boc)₂O, dimethylaminopyridine (DMAP), CH₂Cl₂, room temp., 20 min, 90%; iv, H₂O₂, formic acid, CH₂Cl₂, 0 °C, 30 min, 90%; v, THF, (Ph₃P)₄Pd, reflux, 1 h, 12 mol dm⁻³ Na₂CO₃, reflux 24 h, 50%; vi, CsCO₃, MeOH, room temp., 2 h, 80%; vii, *N*-benzyloxycarbonyl(dimethoxyphosphoryl) glycine trimethylsilylethyl ester, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), CH₂Cl₂, -78 °C, 15 h, 80%; viii, [Rh(1.5-cod)(dipamp)]⁺ BF₄⁻, H₂, MeOH, room temp., 72 h, quant.; ix, 6 mol dm⁻³ HCl, dioxane, room temp., 30 min, quant.

unit formally represents the dimer from (*S*)-*o*-hydroxyphenylalanine and (2*S*,3*R*)-*o*-hydroxyphenylserine. In this communication we describe the total synthesis of biphenomycin A. The key step for the construction of the biphenyl unit is the Pd⁰-catalysed coupling⁶ of an arylboronic acid containing an unprotected aldehyde function with a protected 2-hydroxy-5-



Scheme 2 Reagents and conditions: i, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC), hydroxybenzotriazole, CH₂Cl₂, -10 to 20 °C, 14 h, 80%; ii, AcOH-H₂O (9:1), 50 °C, 2 h, 70%; iii, Bu₄NF, dimethylformamide, room temp., 30 min; iv, C₆F₅OH, EDC, CH₂Cl₂, -15 to 20 °C, 14 h, 95%; v, 6 mol dm⁻³ HCl, dioxane, room temp., 30 min, evaporation, quant.; vi, CHCl₃-NaHCO₃ (aq.), 20 °C, 5 min, 70%; vii, 1 mol dm⁻³ LiOH, dioxane, room temp., 30 min, evaporation, quant.; viii, trimethylsilyl bromide-thioanisole (1:1), CF₃CO₂H, 2 h, room temp., 60%

bromophenylserine. In the course of the synthesis it was necessary to mask two phenolic hydroxy groups, two amino groups, two hydroxy groups and one carboxy group with functions that are compatible with the protecting groups of the α-amino group of hydroxyornithine and the one carboxy group of the hydroxy diisotyrosine which needs to be cleaved during the peptide construction and ring closure steps.

The synthesis of a protected (*R*)-hydroxy-(*S,S*)-diisotyrosine **11** is illustrated in Scheme 1. The aryl bromide **6** represents a completely protected 2-hydroxy-5-bromophenylserine and can be constructed *via* **5** in excellent enantioselectivity and diastereoselectivity by the Evans method.⁷ The Pd⁰-catalysed coupling of **6**, with the boronic acid **7** containing an unprotected aldehyde function gives rise to the biphenyl derivative **8**. After opening of the oxazolidinone, the highly diastereoselective⁸ construction of the *Z*-dehydroamino acid derivative **10** is achieved by condensation of **9** with an appropriate phosphorylglycine⁹ and subsequent completely diastereoselective hydrogenation using the [Rh(1.5-cod)(dipamp)]⁺ BF₄⁻ hydrogenation catalyst {cod = cycloocta-1,5-diene; dipamp = (*R,R*)-1,2-bis[2-(2-methoxyphenyl)-phenylphosphino]ethane}¹⁰ gives rise to the biphenyl unit **11**. The further steps are illustrated in Scheme 2. The construction of the linear substrate **13** for the ring closure reaction is achieved starting from the hydroxyornithine derivative **12**.¹¹

in analogy to the synthesis of biphenomycin B. The only exception is that the free carboxy group is now protected as a methyl ester (in place of a benzyl ester in the biphenomycin B synthesis). The ring closure reaction of the ω -aminopentafluorophenyl ester in the two phase system CHCl_3 - NaHCO_3 (aq.) occurs within 5 min to give a 70% yield without dilution.¹² After saponification of the methyl ester **16**, the four benzylic protecting functions are cleaved by the method of Yajima¹³ using trimethylsilyl bromide-thioanisole in trifluoroacetic acid. The synthesized product is separated by reversed phase chromatography and lyophilized. It is identical in every respect with the natural compound.

Support of this research work by BASF AG, the Fonds der Chemischen Industrie, the Deutsche Forschungsgemeinschaft and the Land Baden-Württemberg is gratefully acknowledged. We thank Dr M. Shirahashi, Fujisawa Pharmaceutical Co. Ltd, for a sample of natural biphenomycin A and Professor Dr G. Jung and Dr J. Metzger, Universität Tübingen, for mass spectra and Dr P. Fischer for NMR spectra.

Received, 16th March 1992; Com. 2/01370C

References

- For Part 82 of the series Amino Acids and Peptides see U. Schmidt and J. Schmidt, *J. Chem. Soc., Chem. Commun.*, 1992, 529.
- Isolation and structure: M. Ezaki, M. Iwami, M. Yamashita, S. Hashimoto, T. Komori, K. Umehara, Y. Mine, M. Kohsaka, H. Aoki and H. Imanaka, *J. Antibiot.*, 1985, **38**, 1453; I. Uchida, N. Shigematsu, M. Ezaki, M. Hashimoto, H. Aoki and H. Imanaka, *J. Antibiot.*, 1985, **38**, 1462; I. Uchida, M. Ezaki, N. Shigematsu and M. Hashimoto, *J. Org. Chem.*, 1985, **50**, 1342.
- Configuration: R. Kannan and D. H. Williams, *J. Org. Chem.*, 1987, **52**, 5435.
- Configuration and conformation: J. C. Hempel and F. K. Brown, *J. Am. Chem. Soc.*, 1989, **111**, 7323; F. K. Brown, J. C. Hempel, J. S. Dixon, S. Amato, L. Mueller and P. W. Jeffs, *J. Am. Chem. Soc.*, 1989, **111**, 7328.
- U. Schmidt, R. Meyer, V. Leitenberger, A. Lieberknecht and H. Griesser, *J. Chem. Soc., Chem. Commun.*, 1991, 275.
- N. Miyaara, T. Yanagi and A. Suzuki, *Synth. Commun.*, 1981, **11**, 513.
- D. A. Evans and A. E. Weber, *J. Am. Chem. Soc.*, 1986, **108**, 6757.
- U. Schmidt, H. Griesser, V. Leitenberger, R. Mangold, R. Meyer and B. Riedl, *Synthesis*, 1992, 487.
- U. Schmidt, A. Lieberknecht and J. Wild, *Angew. Chem.*, 1982, **94**, 797; *Angew. Chem., Int. Ed. Engl.*, 1982, **21**, 770; *Angew. Chem. Suppl.*, 1982, 1682; U. Schmidt, A. Lieberknecht and J. Wild, *Synthesis*, 1984, 53.
- W. S. Knowles, M. J. Sabacky, B. D. Vineyard and D. J. Weinkauff, *J. Am. Chem. Soc.*, 1975, **97**, 2567; B. D. Vineyard, W. S. Knowles, M. J. Sabacky, G. L. Bachman and D. J. Weinkauff, *J. Am. Chem. Soc.*, 1977, **99**, 5946.
- U. Schmidt, R. Meyer, V. Leitenberger, F. Stäbler and A. Lieberknecht, *Synthesis*, 1991, 409.
- U. Schmidt, A. Lieberknecht, H. Griesser and J. Talbiersky, *J. Org. Chem.*, 1982, **47**, 3261; in a two-phase system: U. Schmidt, R. Utz, A. Lieberknecht, H. Griesser, B. Potzulli, J. Bahr, K. Wagner and P. Fischer, *Synthesis*, 1987, 236; U. Schmidt, M. Kroner and H. Griesser, *Tetrahedron Lett.*, 1988, **29**, 4407.
- H. Yajima, N. Fujii, S. Funakashi, T. Watanabe, E. Murayama and A. Otaka, *Tetrahedron*, 1988, **44**, 805.