

1,4-Dialkoxy 1,3-dienes and bis(dienes)–stereocontrolled synthesis and reactivity

Anne Guillaum, Jacques Maddaluno and Lucette Duhamel*

Unité de Recherche Associée au CNRS 464, Faculté des Sciences et Techniques and IRCOF, Université de Rouen, 76821 - Mont St Aignan, France

γ -Aryloxy and -alkoxy α,β -unsaturated dimethyl acetals are direct precursors to (1*Z*, 3*E*)-1,4-dialkoxy 1,3-dienes and bis(dienes) which are fine partners in cycloaddition reactions with classical dienophiles.


Although the Diels–Alder reaction remains a source of constant attention, a restricted number of functionalized dienes have found a place among the commonly used building blocks, *e.g.* Danishefsky's diene. Only a few 1,4-dialkoxydienes have been prepared before¹ and are reported to be especially fragile. Here we report a new stereocontrolled synthesis of 1,4-dialkoxy 1,3-dienes and bis(dienes) together with a few model cycloaddition reactions involving these compounds.

Halogen substitution on chloroacetal **1** by sodium phenate directly yields the corresponding ether **2a** with preservation of the original *E/Z* ratio, Scheme 1.^{2c–d} Secondary sodium alcoholates can also be employed, such as sodium menthylate, but require harsher conditions to provide **2b**. Similarly, energetic stirring of a THF solution of acetal **1** with aqueous sodium hydroxide provides directly and in excellent yield bis(acetal) **3**. The 2*E*,2'*E*/2*E*,2'*Z* ratio is roughly 60:40.[†]


These acetals can be deprotonated in THF using strong bases such as *n*- or *tert*-butyllithium[‡] or potassium bis(trimethylsilyl)amide KHMDS. Amide **2a** selectively gives the corresponding 1,4-dienol diether **4a** whatever the base.^{2d§} By contrast **2b** yields **4b** (100%) when treated with BuLi and **5b** (100%) with KHMDS, Table 1. The 1*Z*,3*E* to 1*E*,3*E* ratio in **4** is *ca.* 90:10, a striking result with respect to the high 1*E*,3*E* selectivity obtained in identical conditions with corresponding arylthio dimethyl acetals (**2a** with ArS instead of PhO).² Similarly, bis(acetal) **3** provides either bis(dienes) **6** or a mixture of bis(dienes) **6** and **7**, Scheme 2, of which the ratio

depends on the base employed.[¶] The yields are good to excellent.


It was then interesting to assess the synthetic potential of these compounds in classical [4 + 2] cycloaddition reactions. Our method provides dienes **4–7** of sufficient purity and stereohomogeneity to be used as crude starting materials, Scheme 3. We first condensed **4a** with *N*-methyl maleimide (NMM). Adduct **8** was obtained with total *endo* selectivity. As expected, the 1*Z*,3*E* configuration of the starting diene leads to 3,6-*trans* disubstituted cyclohexenyl structures. The conformation of the bicyclic compounds is of the 'convex-boat type',



Scheme 2 Reagents and conditions: i, BuLi -40°C for 15 min. or KHMDS at room temp. in THF for 1 h., 1 equiv. per acetal, yield 88–99%




Scheme 3 Reagents and conditions: i, 1 equiv. *N*-methylmaleimide in toluene at 110°C for 6 h; ii, neat methyl acrylate at 80°C for 5 d or 3 equiv. methyl acrylate in MeOH for 1 d under 12 kbar, yield 71–73%



Scheme 1 Reagents and conditions: i, 4 equiv. PhONa, $\text{H}_2\text{O}/\text{THF}$ at room temp. for 2 d, yield 98% or 1 equiv. sodium menthylate at 70°C in THF/hexamethylphosphoramide (HMPT) for 1 d, yield 54%; ii, 3 equiv. NaOH, $\text{H}_2\text{O}/\text{THF}$ at 70°C with Bu_4NI for 1 d, yield 93%

Table 1 Deprotonation of acetals 2–3

Acetal	R	Base	'Endo'	'Exo'
2a	Ph	<i>n</i> - or <i>tert</i> -BuLi or KHMDS	100 (4a)	0
2b	Menth	BuLi	100 (4b)	0
2b	Menth	KHMDS	0	100 (5b)
3	Bis	<i>n</i> - or <i>tert</i> -BuLi	100 (6)	0
3	Bis	KHMDS	50 (6)	50 (7)



Scheme 4 Reagents and conditions: i, 1 equiv. *N*-methylmaleimide in THF at room temp. for 1 d; ii, 1 equiv. *N*-methylmaleimide in THF at 40°C in the same pot for 2 d, yield 59%; iii, 1 equiv. tetracyanoethylene in THF at -40°C in the same pot for 15 min, yield 57%

the methoxy group occupying an axial position. Both menthoxy ethers **4b** and **5b** also react with NMM with 42 and 32% de, respectively. The regioselectivity has been tested on **4a** with respect to methyl acrylate under thermal or high-pressure (12 kbar) activation conditions. The single *endo* adduct **9** is recovered in both cases, Scheme 3.

The reactivity of bis(diene) **6** has then been considered, Scheme 4. It adds onto NMM at room temperature in THF to provide intermediate monoadduct **10**, which on slight warming (40 °C) gives the double adduct **11**. Compound **10** also reacts with other dienophiles to yield mixed adducts. For instance using NMM and then tetracyanoethylene at -40 °C gives **12** in a one pot reaction. The second addition takes place with some diastereoselectivity with respect to the first. In the cases represented here, the de of **11** is almost zero while it reaches 80% in the case of **12**.

The '*exo*' bis(diene) **7** adds easily to NMM at room temperature to provide almost instantaneously double adduct **13**, Scheme 5. Advantage has been taken of this very high reactivity to prepare bis(heterocycle) **14**, a potential precursor of pseudo-disaccharides, applying **7** in a hetero-Diels-Alder

reaction with diethyl 2-oxomalonate at room temperature in THF. Yield of **14** is 87% after flash chromatography.

In conclusion, these results indicate that 1,4-dialkoxydienes **4** and **6** are easily accessible from acetals **2–3**. The configuration of the dienes thus prepared is largely in favour of the *ZE*-isomer. Depending on the substituent, '*exo*' (**5b** and **7**) dienes may also be obtained. The reactivity of all these compounds in [4 + 2] cycloaddition reactions is satisfactory under thermal and high-pressure conditions.

Footnotes

† Corresponding to the statistical combination of two *E/Z* = 80:20 units of **1** which is an industrial intermediate (Rhône-Poulenc Chimie).

‡ As first described for crotonaldehyde and senecialdehyde diethyl acetal.^{3,4}


§ '*Exo*' diene **5a** is not obtained in these basic conditions but can be prepared as a mixture with **5a** by Me₃SiOTf catalysis as described for saturated acetals.⁵

¶ For **6**, 1*Z*,3*E*, 1'*Z*,3'*E*:1*E*,3*E*, 1'*Z*,3'*E* ratio = 90:10 while '*exo*' **7** is obtained as a single 1*E*, 1'*E* isomer.

References

- (a) H. Meister, *Chem. Ber.*, 1963, **96** 1688; (b) J. W. Scheeren, A. T. M. Marcelis, R. W. Aben and R. J. Nivard, *J. R. Neth. Chem. Soc.*, 1975, **94**, 196; (c) R. K. Boeckman Jr, T. M. Dolak and K. O. Culos, *J. Am. Chem. Soc.*, 1978, **100**, 7098; (d) H. Hiranuma and S. I. Miller, *J. Org. Chem.*, 1982, **47**, 5083.
- (a) O. Gaonac'h, J. Maddaluno, J. Chauvin and L. Duhamel, *J. Org. Chem.*, 1991, **56**, 4045; (b) O. Gaonac'h, J. Maddaluno, G. Pié and L. Duhamel, *Tetrahedron Lett.*, 1992, **33**, 2473; (c) O. Gaonac'h, Ph.D. Thesis, University of Rouen, 1992; (d) J. Maddaluno, O. Gaonac'h, Y. Le Gallic and L. Duhamel, *Tetrahedron Lett.*, 1995, **36**, 8136; (e) J. Maddaluno, O. Gaonac'h, A. Marcual, L. Toupet and C. Giessner-Prettre, *J. Org. Chem.*, in the press.
- C. Mioskowski, S. Manna and J. R. Falck, *Tetrahedron Lett.*, 1984, **25**, 519.
- A. Deagostino, C. Prandi and P. Venturello, *Tetrahedron*, 1996, **52**, 1433 and references cited therein.
- P. G. Gassman, S. J. Burns and K. B. Pfister, *J. Org. Chem.*, 1993, **58**, 1449.

Received, 11th March 1996; Com. 6/01676F



Scheme 5 Reagents and conditions: i, 2 equiv. *N*-methylmaleimide in THF at room temp. yield 83%; ii, 2 equiv. diethyl 2-oxomalonate at room temp. in THF (X = CO₂Et), yield 87%