

Article

Discovery and Elucidation of Counteranion Dependence in Photoredox Catalysis

Elliot P Farney, Steven J Chapman, Wesley B. Swords,
Marco D. Torelli, Robert J Hamers, and Tehshik P. Yoon

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.9b01885 • Publication Date (Web): 21 Mar 2019

Downloaded from <http://pubs.acs.org> on March 21, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

is published by the American Chemical Society, 1155 Sixteenth Street N.W.,
Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society.
However, no copyright claim is made to original U.S. Government works, or works
produced by employees of any Commonwealth realm Crown government in the course
of their duties.

Discovery and Elucidation of Counteranion Dependence in Photoredox Catalysis

Elliot P. Farney, Steven J. Chapman, Wesley B. Swords, Marco D. Torelli, Robert J. Hamers, Tehshik P. Yoon*

Department of Chemistry, University of Wisconsin–Madison, 1101 University Avenue, Madison, Wisconsin 53706

ABSTRACT: Over the past decade, there has been a renewed interest in the use of transition metal polypyridyl complexes as photoredox catalysts for a variety of innovative synthetic applications. Many derivatives of these complexes are known, and the effect of ligand modifications on their efficacy as photoredox catalysts has been the subject of extensive, systematic investigation. However, the influence of the photocatalyst counteranion has received little attention, despite the fact that these complexes are generally cationic in nature. Herein, we demonstrate that counteranion effects exert a surprising, dramatic impact on the rate of a representative photocatalytic radical cation Diels–Alder reaction. A detailed analysis reveals that counteranion identity impacts multiple aspects of the reaction mechanism. Most notably, photocatalysts with more non-coordinating counteranions yield a more powerful triplet excited state oxidant and longer radical cation chain length. It is proposed that this counteranion effect arises from Coulombic ion-pair interactions between the counteranion and both the cationic photoredox catalyst and the radical cation intermediate, respectively. The comparatively slower rate of reaction with coordinating counteranions can be rescued by using hydrogen-bonding anion binders that attenuate deleterious ion-pairing interactions. These results demonstrate the importance of counteranion identity as a variable in the design and optimization of photoredox transformations and suggest a novel strategy for the optimization of organic reactions using this class of transition metal photocatalysts.


Introduction

Ruthenium(II) polypyridyl complexes have been among the most widely studied molecular photocatalysts for a variety of applications. The photophysical, electrochemical, and physical properties of this class of luminescent transition metal complexes have been extensively characterized. They generally exhibit strong absorbance in the visible spectrum, feature high intersystem crossing efficiency, and can participate in a diverse range of photoinduced electron- and energy-transfer processes. Because of these attractive features, Ru(II) photocatalysts were instrumental in the early development of solar fuels technologies;¹ in addition, some of the best light-harvesting sensitizers for dye-sensitized solar cells belong to this family of complexes.² Over the past decade, the recognition that Ru(bpy)₃²⁺ and its analogues are also useful photocatalysts for organic transformations has stimulated a renewal of interest in photochemical synthesis.³ Because of the exceptional utility of Ru(bpy)₃²⁺ in so many diverse applications, numerous structurally varied Ru(II) polypyridyl photocatalysts have been prepared, and the effects of ligand modifications on catalyst properties are well-understood (Figure 1).⁴

The effect of counteranion structure on the photoactivity of these cationic complexes, on the other hand, has not been subject to similar systematic study. In this paper, we document the discovery of the unexpected impact counteranion identity plays on the efficiency of a radical cation Diels–Alder cycloaddition, a representative photoredox transformation. We rationalize the observed rate increase as the consequence of (1) a change in the photocatalyst ground-state electrochemical properties, (2) a significant shift in its triplet-state energy, and (3) an increase in the efficiency of radical cation chain propagation. The results reported herein suggest that this counterion effect may be an unappreciated but important phenomenon in many photoredox reactions. Understanding the impact of this experimental variable, therefore, should benefit the growing

community of scholars interested in the use of these complexes as photoredox catalysts in organic chemistry.

Ligand effects well-characterized



Effect of counteranion structure poorly documented




Figure 1. Structurally varied Ru(II) photocatalysts.


Results and Discussion

Counterion Effects in Radical Cation Cycloadditions

Several years ago, we reported that visible light photoredox catalysis offered an efficient means to conduct radical cation Diels–Alder cycloadditions between a wide range of electron-rich

styrenes and diverse dienes.⁶ The highly electron-deficient $[\text{Ru}(\text{bpz})_3](\text{BAR}^{\text{F}}_4)_2$ complex⁷ proved to be a potent photocatalyst for this transformation, providing excellent rates and yields at ambient temperatures with as little as 0.5 mol% of photocatalyst. Our proposal for the mechanism of this reaction is briefly summarized in Scheme 1. Photoexcitation of $\text{Ru}(\text{bpz})_3^{2+}$ with visible light results in the efficient formation of a long-lived redox-active triplet state. The electron-deficient bpz ligands render the photoexcited catalyst a substantially stronger oxidant (+1.4 V vs SCE) than the parent $\text{Ru}(\text{bpy})_3^{2+}$ catalyst (+0.89 V vs SCE), enabling the one-electron photooxidation of anethole (**4**, +1.1 V vs SCE). The resulting alkene radical cation undergoes rapid [4+2] cycloaddition with diene **5** to afford product radical cation **6**⁺⁺.⁹ Formation of the neutral product can occur by one of two mechanisms: either radical chain-propagating oxidation of another equivalent of alkene **4** or by chain-terminating oxidation of the reduced $\text{Ru}(\text{bpz})_3^+$ catalyst.¹⁰ The latter process regenerates the photoactive $\text{Ru}(\text{II})$ state of the catalyst and closes the catalytic cycle.

Scheme 1. Proposed Photocatalytic Radical Cation Diels–Alder Cycloaddition Mechanism



Despite the efficiency and broad scope of this reaction, the $\text{Ru}(\text{bpz})_3^{2+}$ chromophore suffers from limited solubility in non-polar organic solvents. Empirical screening indicated that the reaction proceeds more rapidly in these solvents and resulted in the use of CH_2Cl_2 in optimized reaction conditions. Nevertheless, the solubility of the bpz complex in CH_2Cl_2 is modest, and reactions conducted at moderate catalyst loadings are often visibly heterogeneous. These constraints limited our ability to systematically investigate catalyst structure on the rate of photoredox reactions. We thus became interested in the use of strongly oxidizing photocatalysts with greater lipophilicity that might be freely soluble in low-dielectric solvents.

As a starting point for these studies, we prepared a series of photocatalysts based upon the $\text{Ru}(\text{btfmb})_3^{2+}$ chromophore (Figure 1, **3**; btfmb = 4,4'-bis(trifluoromethyl)-2,2'-bipyridyl). The photophysical and electrochemical properties of the homoleptic $[\text{Ru}(\text{btfmb})_3](\text{PF}_6)_2$ complex were previously investigated in acetonitrile by Furue and Kamachi.¹¹ Given the oxidizing potential reported for its excited state (+1.3 V vs SCE), we hypothesized that $\text{Ru}(\text{btfmb})_3^{2+}$ would be an effective photooxidative catalyst for electron-rich styrenes such as **4**. Moreover, we hoped that the lipophilic CF_3 substituents would improve the solubility of the photocatalyst in non-polar organic solvents compared to $\text{Ru}(\text{bpz})_3^{2+}$. In order to maximize the organic solubility of this chromophore, we prepared a

series of salts bearing a variety of lipophilic counteranions (**3a–f**) and assessed their activities in a model photoreaction.

The results of this initial catalyst screen for the radical cation Diels–Alder cycloaddition between anethole (**4**) and isoprene (**5**) are summarized in Table 1. In general, these various photocatalysts each promoted the reaction, but surprisingly, the rates of reaction varied dramatically depending on counteranion identity. While the cycloaddition of **4** and **5** is complete in 20 min using 1 mol% of $\text{BAR}^{\text{F}}_4^-$ catalyst **3a**, the reaction proceeds to only 55% yield after 24 h with the analogous triflate catalyst **3b**, 14% yield with the 3,5-bis(trifluoromethyl)benzenesulfonate ($\text{Ar}^{\text{F}}\text{SO}_3^-$) catalyst **3d**, and only 2% yield with the tosylate catalyst **3e**. No conversion was observed in this timeframe using the carboxylate complex **3f**. Thus, there appears to be a correlation between the rate of product formation and the non-coordinating nature of the catalyst counteranions.¹² Unaware of any previous report of similarly dramatic counterion effects on the rate of organic photoredox transformations, we elected to investigate the origins of this phenomenon.

Table 1. Counteranion Effect on the Rate of Radical Cation Diels–Alder Cycloaddition.^a

| entry | catalyst | time | yield 6 ^b | unreacted 4 ^b |
|-------|---|--------|-----------------------------|---------------------------------|
| 1 | 3a ($\text{BAR}^{\text{F}}_4^-$) | 20 min | 98% | 0% |
| 2 | 3b (TfO^-) | 24 h | 55% | 37% |
| 3 | 3d ($\text{Ar}^{\text{F}}\text{SO}_3^-$) | 24 h | 14% | 80% |
| 4 | 3e (TsO^-) | 24 h | 2% | 95% |
| 5 | 3f ($\text{Ar}^{\text{F}}\text{CO}_2^-$) | 24 h | 0% | 100% |

^a General conditions: anethole (0.06 mmol), isoprene (0.18 mmol), CH_2Cl_2 (0.08 M), $[\text{Ru}]$ (1 mol%). Each reaction was subjected to 3 freeze-pump-thaw cycles prior to irradiation from a 23 W CFL bulb. ^b ¹H NMR yields referenced to trimethyl(phenyl)silane internal standard.

To begin to understand this effect, we performed Stern–Volmer analyses of the relationship between the concentration of anethole (**4**) in CH_2Cl_2 and the photoluminescence intensity of the $\text{BAR}^{\text{F}}_4^-$, PF_6^- , and $\text{Ar}^{\text{F}}\text{SO}_3^-$ complexes of $\text{Ru}(\text{btfmb})_3^{2+}$. This investigation demonstrated that the degree of excited-state quenching between the photoexcited catalyst and the organic substrate (i.e., the Stern–Volmer constant, K_{sv}) decreased by two orders of magnitude from the least coordinating counteranion, $\text{BAR}^{\text{F}}_4^-$, to the most coordinating counteranion in this study, $\text{Ar}^{\text{F}}\text{SO}_3^-$ (Figure 2A). This is consistent with the markedly superior reactivity of the $\text{BAR}^{\text{F}}_4^-$ complex. The value of K_{sv} is dependent both upon the excited-state lifetime of the photocatalyst (τ) and the bimolecular electron-transfer rate constant (k_q); $K_{\text{sv}} = \tau k_q$. To deconvolute whether the large change in the Stern–Volmer constant arises primarily from a change in catalyst triplet lifetime or in the electron-transfer rate constant, we measured τ for each $\text{Ru}(\text{btfmb})_3^{2+}$ complex (Table 2). These results show that while the counteranion does have an influence on triplet lifetime, the effect is relatively small — approximately two-fold over the range of counteranions investigated. The impact on k_q , therefore, is much larger, spanning two orders of magnitude from $\text{Ar}^{\text{F}}\text{SO}_3^-$ **3d**




Figure 2. A. Stern–Volmer plots for excited-state quenching of catalysts **3a**–**c** in CH_2Cl_2 . B/C. Effect of counteranion identity on excited-state properties of $[\text{Ru}(\text{btfmb})_3](X)_2$, X is indicated in the legend. (B) Absorption (solid line) and photoluminescence (dashed line) spectra in MeCN. (C) Absorption (solid line) and photoluminescence (dashed line) spectra in CH_2Cl_2 .

to BARF_4^- **3a**. Thus, the unanticipated conclusion from these preliminary studies is that the identity of the photocatalyst counteranion can impact a photooxidative reaction by dramatically altering the intrinsic rate constant of bimolecular electron transfer to the photocatalyst excited state.

Table 2. Spectroscopic Properties and Stern–Volmer Quenching Constants^a

| entry | catalyst | λ_{abs} (nm) | λ_{em} (nm) | K_{SV} ($\times 10^2 \text{ M}^{-1}$) | τ (ns) | k_q ($\times 10^8 \text{ M}^{-1} \text{ s}^{-1}$) |
|-------|-----------|--------------------------------|-------------------------------|---|----------------|--|
| 1 | 3a | 453 | 573 | 17 | 520 | 33 |
| 2 | 3c | 458 | 605 | 4.2 | 860 | 4.9 |
| 3 | 3d | 458 | 618 | 0.61 | 950 | 0.64 |

^a Data acquired in CH_2Cl_2 .

This finding was surprising. While the importance of the bipyridyl ligand structure in the design and optimization of photocatalytic reactions is well appreciated,¹³ the effect of the catalyst counterion on photocatalytic reaction rates has received significantly less attention. Meyer and coworkers have examined ion-pairing effects on the photophysics of Ru(II) polypyridyl chromophores.¹⁴ These investigations show that addition of Cl^- to solutions of $[\text{Ru}(\text{bpy})_2(\text{deeb})](\text{PF}_6)_2$ in CH_2Cl_2 results in the formation of tight ion pairs and a concomitant decrease in triplet excited-state energy and lifetime.¹⁵ However, neither the impact of structurally complex organic counteranions on the photochemical properties of Ru(II) complexes, nor the effects of ion pairing on the rate of synthetic photocatalytic applications, have been systematically explored.¹⁶

Spectroscopic, Electrochemical, and Quantum Yield Studies

We have collected absorption, emission, and electrochemical data for a representative series of $\text{Ru}(\text{btfmb})_3^{2+}$ complexes bearing BARF_4^- , PF_6^- , Ar^fSO_3^- , and TsO^- counteranions. These data are depicted in Figure 2B and C. First, we obtained spectral data for these complexes in acetonitrile (Figure 2B). Both the absorption and emission spectra are superimposable in MeCN, consistent with the attenuated impact of ion pairing in high-dielectric solvents. On the other hand, Coulombic effects are more significant in non-polar solvents, which

are often ideal for applications in organic synthesis. Figure 2C shows absorption and emission spectra for the same series of catalysts in CH_2Cl_2 . The impact of counteranion identity on the absorption spectrum in this relatively non-polar solvent is modest, suggesting that if differences in Coulombic interactions exert any influence on the ground-state properties of the photocatalyst or on the singlet excited state, it is a small effect. In contrast, the photoluminescence spectra of the various complexes differ markedly. Most notably, the λ_{max} of photoluminescence varies by 52 nm from the least coordinating (BARF_4^- , 573 nm) to the most coordinating (TsO^- , 625 nm) counteranion, corresponding to a substantial energy difference of 4.2 kcal/mol (0.18 eV).

Thus, altering the identity of the counteranion produces an unexpectedly large change in the energy of the emissive triplet state of $\text{Ru}^*(\text{btfmb})_3^{2+}$. One would expect these changes to be reflected in excited-state redox potentials. To quantify this effect, we measured the one-electron reduction potential $E(\text{Ru}^{2+/+})$ of the $\text{Ru}(\text{btfmb})_3^{2+}$ complexes in CH_2Cl_2 . Each measurement was made using a matching $n\text{-Bu}_4\text{N}^+\text{X}^-$ salt as a supporting electrolyte in order to avoid complications arising from counteranion exchange. Counteranions of lower Lewis basicity resulted in significant anodic shifts in the ground-state potentials, with the largest and most significant effect observed for the BARF_4^- counteranion (Table 3). This effect can also be rationalized as a consequence of ion pairing where the one-electron reduction of the least electrostatically stabilized BARF_4^- complex is more energetically favorable than the tightly ion-paired tosylate complex. To calculate the excited-state redox potential, we made the commonly utilized assumption^{17,18} that the Gibbs free energy change for the S_0 to T_1 transition is represented by the energy of the corresponding photoluminescence maximum (ΔG_{ES}). The catalytically relevant first triplet excited-state reduction potential $E(\text{Ru}^{2+*/+})$ can then be approximated from the sum of ΔG_{ES} and $E(\text{Ru}^{2+/+})$. As the data in Table 3 show, these potentials span a range of 480 mV (11 kcal/mol), with the BARF_4^- complex having the most positive reduction potential of +1.52 V vs SCE. The conclusion from these studies, therefore, is that the degree of ion pairing has a synergistic effect on both the excited-state triplet energy and on the ground-state electrochemical potential, leading to a large net dependence of photooxidant strength on the identity of the catalyst counteranion. These results are consistent with the experimentally observed effect of counteranion identity on the radical cation Diels–Alder reaction described above. The most non-coordinating

counteranion (BAR^{F_4-}) results in the largest driving force for photoinduced electron transfer, consistent with a faster rate of photoinitiation and a shorter reaction time.


Table 3. Ground- and Excited-State Redox Potentials for $\text{Ru}(\text{btfmb})_3^{2+}$ in CH_2Cl_2 .^a

| entry | catalyst | ΔG_{ES} | $E(\text{Ru}^{2+/+})$ | $E(\text{Ru}^{2+*/+})$ |
|-------|-----------|------------------------|-----------------------|------------------------|
| 1 | 3a | 2.17 eV | −0.65 V | +1.52 V |
| 2 | 3c | 2.05 eV | −0.89 V | +1.16 V |
| 3 | 3d | 2.01 eV | −0.93 V | +1.08 V |
| 4 | 3e | 1.99 eV | −0.95 V | +1.04 V |

^aElectrochemical potentials were measured through cyclic voltammetry in a standard three-electrode set-up, scan rate = 100 mV/s. A 100 mM solution of a $n\text{-Bu}_4\text{N}^+\text{X}^-$ salt that matched the photocatalyst counteranion was used as a supporting electrolyte. Potentials were corrected to SCE through an external ferrocene reference.

In order to rationalize the impact of counteranion identity on the triplet excited-state energy of the $\text{Ru}(\text{btfmb})_3^{2+}$ chromophore, we propose an explanation based upon an empirical physical model for charge redistribution between the ground and electronically excited states of this canonical class of transition metal photocatalysts (Scheme 2). The ground state of the $\text{Ru}(\text{btfmb})_3^{2+}$ chromophore has D_3 symmetry and consequently cannot support a permanent dipole moment. On the other hand, the emissive states of $\text{Ru}(\text{II})^*$ tris(bipyridyl) complexes are understood to be metal-to-ligand charge-transfer (MLCT) triplets, and considerable experimental evidence supports the contention that the transferred electron is localized to a single ligand without significant delocalization across the other two ligands.¹⁹ Thus, electronically excited $\text{Ru}^*(\text{btfmb})_3^{2+}$ is best conceptualized as a C_2 -symmetric, charge-separated state with an oxidized $\text{Ru}(\text{III})$ core and a single reduced $\text{btfmb}^{\cdot-}$ ligand (Scheme 2). This lower-symmetry MLCT state would therefore be expected to have a very large dipole moment. Meyer has estimated the dipole moment of the triplet $\text{Ru}^*(\text{bpy})_3^{2+}$ state to be approximately 14 D.²⁰ If the photocatalyst exists largely in an ion-paired state in non-polar solvents, stabilizing charge–dipole interactions should have a larger effect on the triplet excited state than they do on the ground state. One would further expect that more strongly coordinating anions, which produce tighter ion pairs, would better stabilize the triplet excited state. Finally, a strong solvent dependence would be consistent with this model, as charge–dipole interactions are attenuated by increasing solvent dielectric.

Scheme 2. Representation of the ground state (no dipole) and triplet excited state (significant dipole) for $[\text{Ru}(\text{btfmb})_3](\text{X})_2$.



In the radical cation Diels–Alder reaction, the radical cation intermediates ($4^{+\cdot}$ and $6^{+\cdot}$) would also be expected to exist as ion pairs, and the most reasonable counteranion would be that introduced by the photocatalyst.¹⁶ We wondered if this ion-pairing interaction might also affect the dynamics of the product-forming cycloaddition and chain propagation steps as well as the photoinitiation step. To investigate this question, we utilized the same protocol we previously described for estimating the chain length in radical cation cycloadditions.¹⁰ First, we measured the reaction quantum yield with the BAR^{F_4-} and $\text{Ar}^{\text{F}}\text{SO}_3^-$ catalysts through chemical actinometry (see Supporting Information). The quantum yield using the BAR^{F_4-} catalyst **3a** (1 mol%) was measured to be $\Phi = 26$, comparable to the value we determined for the corresponding $[\text{Ru}(\text{bpz})_3](\text{BAR}^{\text{F}_4-})_2$ catalyst in previous studies.¹⁰ However, when $\text{Ar}^{\text{F}}\text{SO}_3^-$ complex **3d** was utilized as the photocatalyst, we measured a significantly decreased quantum yield of $\Phi = 0.35$. To correct for the differing efficiency of photoinitiation by photocatalysts with different excited-state oxidation potentials, we divided the measured quantum yield values by the quenching fraction. The quenching of **3a** by anethole was highly efficient ($Q > 0.99$), and thus the estimated average chain length is the same as the quantum yield ($CL = 26$). The quenching fraction for **3d** was lower ($Q = 0.83$), but the resulting average chain length was still calculated to be quite low ($CL = 0.42$). Thus, in addition to influencing the ground and excited states of the photocatalyst, the counteranion impacts the efficiency of the subsequent radical chain reaction.

These results have several significant implications. First, counteranion identity is a previously underappreciated variable in the optimization of organic photoredox reactions that has the potential to dramatically impact the success and efficiency of synthetically useful organic reactions. Second, counteranion effects impact multiple aspects of the photocatalytic mechanism, including the energy of the reactive triplet excited state, the rate of the electron-transfer photoinitiation event, and the dynamics of non-photochemical product-forming radical chain propagation events. Finally, because the strength of ion-pairing interactions is sensitive to solvent dielectric, these counterion effects are expected to be most important in the relatively non-polar organic solvents that are often optimal for synthetic applications. Such effects may be important in a much wider range of organic photoredox reactions than previously appreciated.


Hydrogen-Bonding Anion Binders as Co-Catalysts

The model proposed above suggests that the rate of photocatalytic Diels–Alder cycloaddition is strongly influenced by Coulombic interactions between the counteranion and both the cationic photocatalyst and the radical cation intermediates. As a further test of this model, we hypothesized that other strategies for disrupting ion pairing might be used to exert a similar effect. In particular, we drew inspiration from a concept pioneered by Jacobsen: hydrogen-bonding organocatalysts can accelerate reactions involving various cationic reactive intermediates by binding their associated counteranions.^{21,22} We hypothesized that the tight pairing between a Lewis basic counteranion and $\text{Ru}^*(\text{btfmb})_3^{2+}$ could be disrupted by addition of an appropriate hydrogen-bonding anion binder, recapitulating the rate increases we observed using weakly coordinating counteranions.

Our investigations focused on the use of sulfonate complex **3d** as a photoredox catalyst for the radical cation Diels–Alder reaction. As

described previously, **3d** is markedly less reactive than the optimal $\text{BAr}^{\text{F}}_4^-$ complex **3a**. Several recent reports have shown that C_{3v} -symmetric thiophosphotriamide **7** is an effective hydrogen-bond donor for binding sulfonate anions,²³ and we imagined that the sequestration of the $\text{Ar}^{\text{F}}\text{SO}_3^-$ counteranion by **7** might attenuate its propensity to participate in tight ion-pairing interactions. To test this hypothesis, we conducted Diels–Alder cycloadditions using 1 mol% of **3d** in the presence and absence of **7** (Table 4). These experiments showed a large rate increase for the Diels–Alder cycloaddition upon addition of just 20 mol% of **7**. Under these conditions, the reaction is complete within 2 h, while only 5% yield of **6** is formed at the same timepoint in the absence of the anion binder. Notably, there is no observable formation of cycloadduct upon irradiation in the presence of **7** without photocatalyst **3d**. This demonstrates that the thiophosphotriamide is not photocatalytically active, and the improvement in photoredox activity thus arises from a synergistic co-catalytic effect.

Table 4. Effect of Ion-binder Co-catalyst **7 on Radical Cation Diels–Alder reaction.**



| entry | [Ru] cat 3d | H-bond cat 7 | time | yield 6 |
|-------|-----------------------|------------------------|------|----------------|
| 1 | 1 mol% | 20 mol% | 2 h | 99% |
| 2 | 1 mol% | none | 2 h | 5% |
| 3 | none | 20 mol% | 2 h | 0% |

We also investigated whether co-catalyst **7** had an influence on the photophysical properties of $\text{Ar}^{\text{F}}\text{SO}_3^-$ photocatalyst **3d** consistent with our proposed model (Figure 3A). The addition of **7** to **3d** induced a large hypsochromic shift in the photoluminescence maximum. This shift was in the direction of the emission maximum of $\text{BAr}^{\text{F}}_4^-$ catalyst **3a**, consistent with the expectation that added **7** would decrease the extent of ion pairing. In contrast, the addition of **7** to the $\text{BAr}^{\text{F}}_4^-$ complex **3a** yielded no change in the emission maximum, even at 50-fold excess of **7** relative to **3a**. This experiment supports the contention that the effect arises from a specific interaction between the thiophosphotriamide and the sulfate counteranion, rather than an interaction with some other component of the reaction mixture or a general medium effect. To further support this contention, we investigated interaction in CD_2Cl_2 using ^1H NMR spectroscopy. Titration of thiophosphotriamide **7** with $n\text{-Bu}_4\text{N}^+ \text{Ar}^{\text{F}}\text{SO}_3^-$ resulted in a significant shift of the aromatic C–H resonances of **7**. A fit of these data to a 1:1 binding model provided an association constant of 1.3×10^6 .




Figure 3. (A) Effect of anion-binding co-catalyst **7** on photoluminescence of **3d**. (B) Stern–Volmer plot in the absence and presence of ion binder **7**.

The influence of added thiophosphotriamide **7** replicated the effect of weakly coordinating anions in other regards as well. First, Stern–Volmer analysis indicates that the rate of quenching of **3d** by anethole is significantly faster upon the addition of the thiophosphotriamide (Figure 3B). This is consistent with the observed increase in triplet excited-state energy with greater concentrations of **7**. Second, we observed a substantial effect on the radical cation chain length (see Supporting Information). The addition of 20 mol% of **7** yielded a 20-fold increase in both the quantum yield and apparent radical chain length of the reaction, suggesting that anion binder **7** can influence the dynamics of the chain process by disrupting ion pairing.

Thus, we have been able to recapitulate the observed effect of non-coordinating anions on the photocatalytic activity of $\text{Ar}^{\text{F}}\text{SO}_3^-$ catalyst **3d** using hydrogen-bonding co-catalyst **7**. The thiophosphotriamide disrupts ion pairing by binding the sulfonate counteranion, which results in significant increases to the photocatalyst's triplet excited-state energy, the rate of photoinduced electron transfer, the length of the radical cation chain process, and the overall efficiency of the photocatalytic Diels–Alder process. These results support the contention that Coulombic effects can be more significant in photoreactions than previously appreciated. Moreover, these results suggest that the use of anion-binding organocatalysts could be a conceptually orthogonal strategy for optimization of the growing class of synthetically useful photoredox transformations.

Conclusion

The studies summarized above suggest several important implications for the design, understanding, and optimization of photocatalytic processes. First, counterion effects can exert a significant impact on the observed rate of radical cation reactions initiated by photoredox catalysis. The degree of ion pairing between the counteranion and both the Ru(II) photoredox catalyst and the oxidized radical cation intermediate can influence the efficiency of multiple steps in the mechanism of these reactions. Thus, these studies indicate that modulating the degree of ion pairing is an important unexplored variable in the optimization of this class of transformations, and we have described two complementary approaches that can successfully increase the overall rate of a radical cation cycloaddition by several orders of magnitude. Second, the Coulombic interactions that are the putative origin of these effects are most significant in relatively non-polar solvents such as those that are often optimal for synthetic applications. This could indicate that counteranion identity is a particularly important variable for organic photoredox reactions compared to better-established applications of Ru(II) photoredox catalysts in solar energy conversion and biology, where the use of water or other high-dielectric solvents might mask the impact of ion pairing. As interest in the use of this class of transition metal photoredox catalyst for synthetic applications continues to grow, a deeper understanding of the impact of ion-pairing effects will be critical to developing a complete, detailed understanding of the mechanisms in this class of synthetically useful transformations.

ASSOCIATED CONTENT

[†] (a) Kalyanasundaram, K. "Photophysics, Photochemistry and Solar Energy Conversion with Tris(bipyridyl)ruthenium(II) and Its Analogues," *Coord. Chem. Rev.* **1982**, 46, 159–244. (b) Roundhill, D. M. "Photochemistry, Photophysics, and Photoredox Reactions of Ru(bpy)₃²⁺ and Related Complexes," in *Modern Inorganic Chemistry: Photochemistry and Photophysics of Metal Complexes*, Springer: New York, 1994, pp. 165–215.

[‡] Lehn, J. M.; Sauvage, J. P. "Chemical Storage of Light Energy. Catalytic Generation of Hydrogen by Visible Light or Sunlight. Irradiation of Neutral Aqueous Solutions," *Nouv. J. Chim.* **1977**, 1, 449–451.

[§] O'Regan, B.; Grätzel, M. "A Low-Cost, High-Efficiency Solar Cell Based on Dye-Sensitized Colloidal TiO₂ Films," *Nature* **1991**, 353, 737–740.

[¶] For reviews, see: a) Narayanam, J. M. R.; Stephenson, C. R. J. "Visible Light Photoredox Catalysis: Applications in Organic Synthesis," *Chem. Soc. Rev.* **2011**, 40, 102–113. b) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. "Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis," *Chem. Rev.* **2013**, 113, 5322–5363. c) Romero, N. A.; Nicewicz, D. A. "Organic Photoredox Catalysis," *Chem. Rev.* **2016**, 116, 10075–10166. d) Twilton, J.; Le, C.; Zhang, P.; Shaw, M. H.; Evans, R. W.; MacMillan, D. W. C. "The Merger of Transition Metal and Photocatalysis," *Nature Rev. Chem.* **2017**, 1, 0052. e) Zou, Y. Q.; Hörmann, F. M.; Bach, T. "Iminium and Enamine Catalysis in Enantioselective Photochemical Reactions," *Chem. Soc. Rev.* **2018**, 47, 278–290. f) Silvi, M.; Melchiorre, P. "Enhancing the Potential of Enantioselective Organocatalysis with Light," *Nature* **2018**, 554, 41–49.

^{||} Juris, A.; Balzani, V.; Barigelli, F.; Campagna, S.; Belser, P.; von Zelewsky, A. "Ru(II) Polypyridine Complexes: Photophysics, Photochemistry, Electrochemistry, and Chemiluminescence," *Coord. Chem. Rev.* **1988**, 84, 85–277.

^{¶¶} (a) Lin, S.; Ischay, M. A.; Fry, C. G.; Yoon, T. P. "Radical Cation Diels-Alder Cycloadditions by Visible Light Photocatalysis," *J. Am. Chem. Soc.* **2011**, 133, 19350–19353. (b) Lin, S.; Padilla, C. E.; Ischay, M. A.; Yoon, T.

(**Supporting Information.** General experimental preparation, synthesis of Ru(II) polypyridyl catalysts, reaction conditions, UV-Vis and photoluminescence spectroscopy, cyclic voltammetry, and NMR characterization. The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

* E-mail: tyoon@chem.wisc.edu

Notes

ORCID:

Steven J. Chapman: 0000-0003-1570-4580

Wesley B. Swords: 0000-0002-2986-326X

Marco D. Torelli: 0000-0002-7691-6491

Robert J. Hamers: 0000-0003-3821-9625

Tehshik P. Yoon: 0000-0002-3934-4973

ACKNOWLEDGMENT

This work was supported by the NIH (GM095666). The NMR and mass spectroscopy facilities at UW–Madison are funded in part by NIH (S10OD020022-1), NSF (CHE-1048642), the University of Wisconsin, and a generous gift from Paul J. and Margaret M. Bender.

REFERENCES

P. "Visible Light Photocatalysis of Intramolecular Radical Cation Diels–Alder Cycloadditions," *Tetrahedron Lett.* **2012**, 53, 3073–3076. (c) Lin, S. S.; Lies, S. D.; Gravatt, C. S.; Yoon, T. P. "Radical Cation Cycloadditions Using Cleavable Redox Auxiliaries," *Org. Lett.* **2017**, 19, 368–371.

[†] (a) Crutchley, R. J.; Lever, A. B. P. "Ruthenium(II) Tris(bipyrazyl) Dication – A New Photocatalyst," *J. Am. Chem. Soc.* **1980**, 102, 7128–7129. (b) Rillema, D. P.; Allen, G.; Meyer, T. J.; Conrad, D. "Redox Properties of Ruthenium(II) Tris Chelate Complexes Containing the Ligands 2,2'-bipyrazine, 2,2'-bipyridine, and 2,2'-bipyrimidine," *Inorg. Chem.* **1983**, 22, 1617–1622.

[‡] Several other photocatalytic radical cation Diels–Alder reactions have been reported using alternate photocatalysts. For leading references, see: (a) Stevenson, S. M.; Shores, M. P.; Ferreira, E. M. "Photooxidizing Chromium Catalysts for Promoting Radical Cation Cycloadditions," *Angew. Chem. Int. Ed.* **2015**, 54, 6506–6510. (b) Pitre, S. P.; Scaiano, J. C.; Yoon, T. P. "Photocatalytic Indole Diels–Alder Cycloadditions Mediated by Heterogenous Platinum-Modified Titanium Dioxide," *ACS Catal.* **2017**, 7, 6440–6444. (c) Gieseler, A.; Steckhan, E.; Wiest, O.; Knoch, F. "Photochemically Induced Radical Cation Diels–Alder Reaction of Indole and Electron-Rich Dienes," *J. Org. Chem.* **1991**, 56, 1405–1411.

[§] Bauld, N. L. "Cation Radical Cycloadditions and Related Sigmatropic Reactions," *Tetrahedron* **1989**, 45, 5307–5363.

^{||} Cismesia, M. A.; Yoon, T. P. "Characterizing Chain Processes in Visible Light Photoredox Catalysis," *Chem. Sci.* **2015**, 6, 5426–5434.

[¶] Furue, M.; Maruyama, K.; Oguni, T.; Naiki, M.; Kamachi, M. "Trifluoromethyl-substituted 2,2'-bipyridine Ligands. Synthetic Control of Excited-State Properties of Ruthenium(II) Tris-Chelate Complexes," *Inorg. Chem.* **1992**, 31, 3792–3795.

[‡] Strauss, S. H. "The Search for Larger and More Weakly Coordinating Anions" *Chem. Rev.* **1993**, 93, 927.

^{††} (a) Ischay, M. A.; Ament, M. S.; Yoon, T. P. "Crossed Intermolecular [2+2] Cycloaddition of Styrenes by Visible Light Photocatalysis," *Chem. Sci.* **2012**, 2046–2050. (b) Douglas, J. J.; Nguyen, J. D.; Cole, K. P.; Stephenson, C. R. J. "Enabling Novel Photoredox Reactivity via Photocatalyst Selection," *Aldrichim. Acta* **2014**, 47, 15–25.

^{†††} (a) Ward, W. M.; Farnum, B. H.; Siegler, M.; Meyer, G. J. "Chloride Ion-Pairing with Ru(II) Polypyridyl Compounds in Dichloromethane," *J. Phys. Chem. A* **2013**, 117, 8883–8894. (b) Wehlin, S. A. M.; Troian-Gautier, L.; Li, G.; Meyer, G. J. "Chloride Oxidation by Ruthenium Excited-States in Solution," *J. Am. Chem. Soc.*, **2017**, 139, 12903–12906. (c) Troian-Gautier, L.; Wehlin, S. A. M.; Meyer, G. J. "Photophysical Properties of Tetracationic Ruthenium Complexes and Their Ter-Ionic Assemblies with Chloride," *Inorg. Chem.*, **2018**, 57, 12232–12244.

^{††††} The chemical shifts associated with the ligands in the ¹H NMR of the photocatalysts vary as a function of counteranion identity and are consistent with observations made by Meyer. These data are summarized in the Supporting Information. See: Li, G.; Swords, W. B.; Meyer, G. J. "Bromide Photo-oxidation Sensitized to Visible light in Consecutive Ion Pairs," *J. Am. Chem. Soc.* **2017**, 139, 14983–14991.

^{†††††} Four recent reports have described the influence of chiral counteranions on the enantioselectivity of photocatalytic radical cation reactions: (a) Morse, P. D.; Nguyen, T. M.; Cruz, C. L.; Nicewicz, D. A. "Enantioselective Counter-Anions in Photoredox Catalysis: The Asymmetric Cation Radical Diels–Alder Reaction," *Tetrahedron* **2018**, 74, 3266–3272. (b) Yang, Z.; Li,

H.; Li, S.; Zhang, M.-T.; Luo, S. "A Chiral Ion-Pair Photoredox Organocatalyst: Enantioselective Anti-Markovnikov Hydroetherification of Alkenols," *Org. Chem. Front.*, **2017**, 4, 1037–1041. (c) Wang, H.; Ren, Y.; Wang, K.; Man, Y.; Xiang, Y.; Li, N.; Tang, B. "Visible Light-Induced Cyclization Reactions for the Synthesis of 1,2,4-Triazolines and 1,2,4-Triazoles," *Chem. Commun.*, **2017**, 53, 9644–9647. (d) Gentry, E. C.; Rono, L. J.; Hale, M. E.; Matsuura, R.; Knowles, R. R. "Enantioselective Synthesis of Pyrrololidines via Noncovalent Stabilization of Indole Radical Cations and Applications to the Synthesis of Alkaloid Natural Products," *J. Am. Chem. Soc.*, **2018**, 140, 3394–3402.

^{‡‡} Adamson, Q. W.; Namnath, J.; Shastry, V. J.; Slawson, V. "Thermodynamic Inefficiency of Conversion of Solar Energy to Work," *J. Chem. Ed.* **1984**, 61, 221.

^{‡‡‡} Flamigni, L.; Barbieri, A.; Sabatini, C.; Ventura, B.; Barigelletti, F. "Photochemistry and Photophysics of Coordination Compounds: Iridium," *Top. Curr. Chem.* **2007**, 281, 143–203.

^{‡‡‡‡} (a) Forster, M.; Hester, R. E. "Resonance Raman Investigation of Electronically Excited Ru(bipyridine)₃²⁺ Using a CW Laser," *Chem. Phys. Lett.* **1981**, 81, 42–47. (b) Bradley, P. G.; Kress, N.; Hornberger, B. A.; Dallinger, R. F.; Woodruff, W. H. "Vibrational Spectroscopy of the Electronically Excited State. 5. Time-Resolved Resonance Raman Study of Tris(bipyridine) Ruthenium(II) and Related Complexes. Definitive Evidence for the Localized MLCT State," *J. Am. Chem. Soc.* **1981**, 103, 7441–7446. (c) Ceulemans, A.; Vanquickenborne, L. G. "On the Charge-Transfer Spectra of Iron(II)- and Ruthenium(II)-Tris(2,2'-bipyridyl) Complexes," *J. Am. Chem. Soc.* **1981**, 103, 2238–2241. (d) Ohsawa, Y.; DeArmond, M. K.; Hanck, K. W.; Morris, D. E.; Whitten, D. G.; Neveux, P. E. "Spatially Isolated Redox Orbitals: Evidence from Low-Temperature Voltammetry," *J. Am. Chem. Soc.* **1983**, 105, 6522–6524.

^{‡‡‡‡‡} Kober, E. M.; Sullivan, B. P.; Meyer, T. J. "Solvent Dependence of Metal-to-Ligand Charge-Transfer Transitions. Evidence for Initial Electron Localization in MLCT Excited States of 2,2'-Bipyridine Complexes of Ruthenium(II) and Osmium(II)," *Inorg. Chem.* **1984**, 23, 2098–2104.

^{‡‡‡‡‡‡} (a) Reisman, S. E.; Doyle, A. G.; Jacobsen, E. N. "Enantioselective Thiourea-Catalyzed Additions to Oxocarbenium Ions," *J. Am. Chem. Soc.* **2008**, 130, 7198–7199. (b) Birrell, J. A.; Desrosiers, J.-N.; Jacobsen, E. N. "Enantioselective Acylation of Silyl Ketene Acetals through Fluoride Anion-Binding Catalysis," *J. Am. Chem. Soc.* **2011**, 133, 13872–13875. (c) Wasa, M.; Liu, R. Y.; Roche, S. P.; Jacobsen, E. N. "Asymmetric Mannich Synthesis of α -Amino Esters by Anion-Binding Catalysis," *J. Am. Chem. Soc.* **2014**, 136, 12872–12875.

^{‡‡‡‡‡‡‡} Gansäuer has recently reported the use of hydrogen bond-donating cocatalysts to influence the rate of titanocene-mediated electron transfer reactions. See: Liedtke, T.; Spanning, P.; Riccardi, L.; Gansäuer, A. "Mechanism-Based Condition Screening for Sustainable Catalysis in Single-Electron Steps by Cyclic Voltammetry," *Angew. Chem. Int. Ed.* **2018**, 57, 5006–5010.

^{‡‡‡‡‡‡‡‡} (a) Cranwell, P. B.; Hiscock, R.; Haynes, C. J. E.; Light, M. E.; Wells, N. J.; Gale, P. A. "Anion Recognition and Transport Properties of Sulfamide-, Phosphoric Triamide- and Thiophosphoric Triamide-Based Receptors," *Chem. Commun.* **2013**, 49, 874–876. (b) Borovika, A.; Tang, P. I.; Klapman, S.; Nagorny, P. "Thiophosphoramidate-Based Cooperative Catalysts for Brønsted Acid Promoted Ionic Diels–Alder Reactions," *Angew. Chem. Int. Ed.* **2013**, 52, 13424–13428.

