

Preliminary communication

Two crystalline toluene-soluble 1,2-organobis(phenylphosphido)dilithium compounds. The monomeric 1,2-C₆H₄(PPh)₂[Li(TMEDA)]₂ and the dimeric [(PhPCH₂CH₂PPh){Li(THF)₂}₂]₂*

David M. Anderson, Peter B. Hitchcock, Michael F. Lappert, Wing-Por Leung
 and Jalal A. Zora

School of Chemistry and Molecular Sciences, University of Sussex, Brighton BN1 9QJ (Great Britain)

(Received June 25th, 1987)

Abstract

Treatment of 1,2-C₆H₄(PPh)₂ with LiBuⁿ in n-C₆H₁₄ and TMEDA (TMEDA = Me₂N[CH₂]₂NMe₂) affords 1,2-C₆H₄(PPh)₂[Li(TMEDA)]₂, which both in the crystal (Li–P 2.58(3) and 2.55(3) Å) and in PhMe solution (*J*(³¹P–⁷Li) 36 Hz) has a monomeric structure with each Li bound to two equivalent P atoms and each P to two equivalent lithium atoms, and yields SnR₂[P(Ph)C₆H₄PPh-1,2] (crystalline *rac* when R = CH(SiMe₃)₂; *rac* + *meso* when R = Me) on treatment with SnR₂Cl₂ or *rac* + *meso*-1,2-C₆H₄[P(Ph)SiMe₃]₂ on treatment with SiClMe₃. By contrast, 1,2-C₂H₄(PPh)₂[Li(THF)₂]₂ (THF = tetrahydrofuran) is an (LiP)₄-skeletal dimer both in the solid and in PhMe (*J*(³¹P–⁷Li) 39 and 47 Hz).

Di(organo)phosphidolithium reagents [Li(PR₂)L_m]_n are of structural interest (X-ray data are available for several compounds, the first being reported in 1983) [1,2] and of value as PR₂-transfer reagents to electrophilic sites. Bifunctional compounds have received scant attention, although other compounds of type (LiX)₂Y are of theoretical concern [3]. Nevertheless, some of them are useful synthons: e.g. Li₂(PhPCH₂CH₂PPh) for phosphorus macrocycles (these ligands are obtained without stereoselectivity at the P-centres) [4a], and Li₂[C₆H₄(PPh)₂-1,2] for benzotriphosphole [4b].

We now report the X-ray structure of the crystalline monomeric 1,2-C₆H₄(PPh)₂[Li(TMEDA)]₂ (TMEDA = *N,N,N',N'*-tetramethylethylenediamine) (1) [11 **], which has each lithium atom bound equivalently to two phosphorus

* No reprints available

** References numbers with asterisks indicate notes in the list of references.




Fig. 1. The molecular structure and atom numbering scheme for 1,2- $\text{C}_6\text{H}_4(\text{PPh})_2[\text{Li}(\text{TMEDA})]_2$ (**1**); for selected data, see Table 1.


atoms. Fig. 1; the two lithium atoms are on opposite sides of the planar 1,2- $\text{C}_6\text{H}_4\text{P}_2$ moiety and the overall configuration of **1** resembles that of 1,2- $\text{C}_6\text{H}_4(\text{CHSiMe}_3)_2[\text{Li}(\text{TMEDA})]_2$ [5]. This is among the first crystalline monomeric $\text{LiP}^<$ compounds; cf. also 1,2- $\text{C}_6\text{H}_4(\text{PPh})_2\text{PLi}(\text{THF})_3$ $\text{LiPPh}_2(\text{PMDETA})$ [**2b**] ($\text{PMDETA} = [\text{Me}_2\text{NCH}_2\text{CH}_2]_2\text{NMe}$), whereas $[\text{Li}(\text{12-crown-4})][\text{PPh}_2]$ has well separated ions [6], and $[\text{Li}(\mu\text{-PPh}_2)(\text{TMEDA})]_2$ is a dimer [2b]. There is a further contrast (Table 1) with the crystalline dimeric Li_4P_4 macrocycle $[(\text{PhPCH}_2\text{CH}_2\text{PPh})\{\text{Li}(\text{THF})_2\}_2]_2$ ($\text{THF} = \text{tetrahydrofuran}$) (**2**) [7].

From variable temperature $^{31}\text{P}\{^1\text{H}\}$ and $^7\text{Li}\{^1\text{H}\}$ NMR data [12 *] we conclude that the structure of **1** in toluene- d_8 solution at low temperature is identical to that


Table 1

Selected bond distances (\AA) and angles ($^\circ$) in compounds **1** and **2**

Li–P		Li–P–Li		P–Li–P	
<i>Compound 1</i>					
Li(1)–P(1)	2.58(3)				
Li(2)–P(1)	2.55(3)	Li(1)–P(1)–Li(2)	103.4(9)	P(1)–Li(1)–P(2)	72(1)
Li(1)–P(2)	2.59(3)				
Li(2)–P(2)	2.59(3)	Li(1)–P(2)–Li(1)	102.0(9)	P(1)–Li(2)–P(2)	72(1)
<i>Compound 2</i> [7]					
Li(1)–P(1)	2.57(1)				
Li(2)–P(1)	2.56(2)	Li(1)–P(1)–Li(2)	138.6(5)	P(1)–Li(1)–P(2)′	113.2(7)
Li(1)–P(2)′	2.57(2)				
Li(2)–P(2)	2.53(2)	Li(1)′–P(2)–Li(2)	128.7(5)	P(1)–Li(2)–P(2)	88.7(5)



in the crystal. Although this situation parallels that found for the di(organo)phosphidolithium compounds $[\text{Li}(\mu\text{-PR}'_2)]_2$ ($\text{R}' = \text{CH}(\text{SiMe}_3)_2$) [8] and $\text{LiPPh}_2(\text{L})$ ($\text{L} = \text{TMEDA}$ or PMDETA) [2b], the solution structures of compounds $[\text{Li}(\text{PR}_2)\text{L}_m]_n$ are sometimes complicated [9]. Likewise, the NMR data for **2** [12 *] were complex, and at first sight at variance with the structure in the crystal, which shows two types of Li atoms but apparently only one type of P atom. However, closer inspection of a molecular model shows that there are two different P sites, labelled P(1) and P(2) in



Scheme 1. The synthesis and some reactions of $1,2\text{-C}_6\text{H}_4(\text{PPh})_2[\text{Li}(\text{TMEDA})]_2$ (**1**, $\text{R} = \text{CH}(\text{SiMe}_3)_2$). Reaction conditions: i: 2LiBu^n , 2TMEDA , $n\text{-C}_6\text{H}_{14}$, 0°C ; ii: 2SiClMe_3 , OEt_2 , 0°C ; iii: SnCl_2R_2 , OEt_2 , 25°C ; iv: SnCl_2Me_2 , OEt_2 , 25°C . Characterisation of compounds **1**–**6**: satisfactory C and H analyses, and NMR and X-ray (for **1** and **2**, Table 1) data.

2, as shown most obviously by the bond angles at these sites: Li(1)–P(1)–Li(2) $138.6(5)^\circ$ and Li(1)'–P(2)–Li(2) $128.7(5)^\circ$. At 305 K, compound **2** in toluene- d_8 showed sharp $^{31}\text{P}\{^1\text{H}\}$ and $^7\text{Li}\{^1\text{H}\}$ singlets; but at 193 K, there were two Li signals each showing coupling to two equivalent phosphorus atoms, whereas there were two P signals each probably coupled to 2 Li's (2 broad signals) [12 *]. A plausible explanation of the low temperature solution spectrum is that it is consistent with structure **2**, in which Li(1) and Li(2) are not, but Li(1) and Li(1)' are, rapidly exchanging [10a *]; this interpretation is also compatible with the ^{31}P and ^7Li solid state NMR data on **2** [10b *].

The compound $[\{\text{Li}(\text{TMEDA})\}_2\{\text{PhPCH}_2\text{CH}_2\text{PPh}\}]_n$ (**3**), obtained from **2** + TMEDA, also showed complicated variable temperature NMR behaviour in THF [12 *]; its sparing solubility in PhMe, its high frequency $^7\text{Li}\{^1\text{H}\}$ signal, and a comparatively sharp $^{31}\text{P}\{^1\text{H}\}$ signal at low temperature are taken as indicative of the presence of a significant proportion of an ionic species.

We suggest that the diphosphidodilithium compounds **1** and **2** have much potential as sources of chelating or bridging 1,2-organobis(phenylphosphido)metal complexes, as illustrated in Scheme 1 for the former by the synthesis of some dialkyltin(IV) (**4** and **5**) or bis(trimethylsilyl) (**6**) derivatives, respectively. Diastereoselectivity is achievable when steric effects are substantial; e.g. for $\text{R}_2\text{Sn}[\text{P}(\text{Ph})\text{C}_6\text{H}_4\text{PPh-}o]$ when $\text{R} = \text{CH}(\text{SiMe}_3)_2$ (**4**) but not when $\text{R} = \text{Me}$ (*rac* (**5a**) (ca. 2/3) + *meso* (**5b**)) (ca. 1/3) (Scheme 1).

Acknowledgements. We thank S.E.R.C. for support, and our colleagues [10 *] for data and discussions.

References

- 1 Cf. E. Hey, P.B. Hitchcock, M.F. Lappert, and A.K. Rai, *J. Organomet. Chem.*, **3** (1987) 1; and refs. therein.
- 2 (a) A. Schmidpeter, G. Burget and W.S. Sheldrick, *Chem. Ber.*, **118** (1985) 3849; (b) R.E. Mulvey, K. Wade, D.R. Armstrong, G.T. Walker, R. Snaith, W. Clegg, and D. Reed, *Polyhedron*, **6** (1987) 987.
- 3 Cf. W.N. Setzler and P. von R. Schleyer, *Adv. Organomet. Chem.*, **24** (1985) 353.
- 4 (a) M. Ciampolini, N. Nardi, P.L. Orioli, S. Mangani and F. Zanobini, *J. Chem. Soc., Dalton Trans.*, (1985) 1179; and refs. therein; (b) F.G. Mann and A.J.H. Mercer, *J. Chem. Soc., Perkin Trans. I*, (1972) 1631.
- 5 M.F. Lappert, C.L. Raston, B.W. Skelton, and A.H. White, *J. Chem. Soc., Chem. Commun.*, (1982) 14.
- 6 H. Hope, M.M. Olmstead, X. Xu, and P.P. Power, *J. Am. Chem. Soc.*, **106** (1984) 819.
- 7 P. Brooks, D.C. Craig, M.J. Gallagher, A.D. Rae, and A. Sarroff, *J. Organomet. Chem.*, **323** (1987) C1; D.M. Anderson, P.B. Hitchcock, M.F. Lappert, and I. Moss, *Inorg. Chim. Acta*, in press.
- 8 P.B. Hitchcock, M.F. Lappert, P.P. Power, and S.J. Smith, *J. Chem. Soc., Chem. Commun.*, (1984) 1669.
- 9 (for $\text{R} = \text{Ph}$) I.J. Colquhoun, H.C.E. McFarlane, and W. McFarlane, *J. Chem. Soc., Chem. Commun.*, (1982) 220; R.A. Bartlett, M.M. Olmstead, and P.P. Power, *Inorg. Chem.*, **25** (1986) 1293; A. Zschunke, M. Riemer, K. Kresch, and K. Issleib, *Phosphorus and Sulfur*, **22** (1985) 349.
- 10 Work in collaboration with A.G. Avent, R.K. Harris, G.A. Lawless, and A. Sabald; (a) 2D NOESY experiments; (b) solid state NMR experiments.
- 11 *Crystal data for 1*: $\text{C}_{30}\text{H}_{46}\text{N}_4\text{Li}_2\text{P}_2$, M 538.6, monoclinic, space group $P2_1/n$, a 17.246(3), b 10.835(3), c 19.399(3) Å; β 109.22(2)°, U 3422.9 Å³, Z = 4, D_c 1.05 g cm⁻³, Mo- K_α radiation, λ 0.71069 Å, μ 1.5 cm⁻¹. The structure of **1** was solved by direct methods and refined by full-matrix least squares to R = 0.13, R' = 0.12, with anisotropic temperature factors for atoms other than

carbon, using 1362 reflections with $|F^2| > \sigma(F^2)$. Atomic coordinates, and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre; thermal parameters and structure factors are available from the authors.

- 12 Selected NMR data (^{31}P at 32.4 MHz (δ in ppm rel. to 85% H_3PO_4), ^7Li at 31.1 MHz (rel. to 5M LiBr), ^{119}Sn at 303 K and 134.3 MHz (rel. to SnMe_4), ^1H at 303 K and 360.1 MHz (rel. to SiMe_4)). 1 (Toluene- d_8): $^{31}\text{P}\{^1\text{H}\}$ at 203 K: δ -45.2 (binomial septet, $^1J(^{31}\text{P}-^7\text{Li})$ 35 Hz); $^7\text{Li}\{^1\text{H}\}$ at 203 K: δ 0.7 (t, $^1J(^{31}\text{P}-^7\text{Li})$ 37 Hz). 2 (Toluene- d_8): $^{31}\text{P}\{^1\text{H}\}$ at 303 K: -73.0 (s, $w_{1/2}$ 60 Hz); $^{31}\text{P}\{^1\text{H}\}$ at 193 K: -65.0 (s, $w_{1/2}$ 200 Hz), -72.0 (s, $w_{1/2}$ 260 Hz); $^7\text{Li}\{^1\text{H}\}$ at 303 K: 1.0 (s, $w_{1/2}$ 20 Hz); $^7\text{Li}\{^1\text{H}\}$ at 193 K: 1.4 (t, $^1J(^{31}\text{P}-^7\text{Li})$ 38 Hz), 0.7 (t, $^1J(^{31}\text{P}-^7\text{Li})$ 47 Hz). 3 (Thf/10% toluene- d_8): $^{31}\text{P}\{^1\text{H}\}$ at 303 K: -60.0 (s, $w_{1/2}$ 30 Hz); $^{31}\text{P}\{^1\text{H}\}$ at 193 K: -56.0 (s, $w_{1/2}$ 30 Hz), -59.0 (s, $w_{1/2}$ 250 Hz), -68.0 (s, $w_{1/2}$ 250 Hz); $^7\text{Li}\{^1\text{H}\}$ at 303 K: 0.5 (s, $w_{1/2}$ 15 Hz); $^7\text{Li}\{^1\text{H}\}$ at 193 K: 2.3 (s, $w_{1/2}$ 60 Hz), 0.2 (s, $w_{1/2}$ 60 Hz). 4 (Benzene- d_6 at 303 K): $^{31}\text{P}\{^1\text{H}\}$: -23.4 (s, $^1J(^{31}\text{P}-^{119}\text{Sn})$ 892, $^1J(^{31}\text{P}-^{117}\text{Sn})$ 856 Hz); $^{119}\text{Sn}\{^1\text{H}\}$: 52.5 (t, $^1J(^{31}\text{P}-^{119}\text{Sn})$ 896 Hz). 5 (Benzene- d_6 at 303 K; peaks marked* refer to 5a): $^{31}\text{P}\{^1\text{H}\}$: -43.4* ($^1J(^{31}\text{P}-^{119}\text{Sn})$ 640 Hz), -50.3 ($^1J(^{31}\text{P}-^{119}\text{Sn})$ 698 Hz); $^{119}\text{Sn}\{^1\text{H}\}$: +64* ($^1J(^{31}\text{P}-^{119}\text{Sn})$ 639 Hz); +59.9 ($^1J(^{31}\text{P}-^{119}\text{Sn})$ 695 Hz); ^1H : 6.95 (m, ArH), 0.03 (s, $\text{Sn}(\text{CH}_3)$), 0.02* (s, $\text{Sn}(\text{CH}_3)$), -0.01 (s, $\text{Sn}(\text{CH}_3)$). 6 (Benzene- d_6 at 303 K): $^{31}\text{P}\{^1\text{H}\}$: -45.3, -45.9 ppm. Variable temperature data on *A* are complicated and will be reported later.