


Carbon Dioxide Chemistry of a Binuclear Iridium(0) Complex. Rapid and Reversible Oxygen Atom Transfer from Carbonate

Mark K. Reinking, Jinfeng Ni, Phillip E. Fanwick,[†] and Clifford P. Kubiak^{*,1}

Department of Chemistry, Purdue University
West Lafayette, Indiana 47907

Received February 13, 1989

We report the activation of CO₂ by a binuclear complex of Ir(0). The activation and reduction of carbon dioxide are areas of potential importance for artificially recycling the products of combustion and respiration.^{2,3} The activation of CO₂ may be accomplished by (i) chemical,²⁻¹⁸ (ii) electrochemical,¹⁹⁻²¹ or (iii) photochemical²²⁻²⁶ means. A serious impediment to chemical or electrochemical reduction of CO₂, however, is the disproportionation of the single-electron-transfer product, CO₂^{•-} (eq 1).



Reactions of low-valent transition-metal complexes with CO₂ often result in the production of 1 equiv of CO₃²⁻ for each equivalent of CO₂ reduced to CO.^{11-14,16} The resulting carbonate ligands are usually unreactive end products that serve to halve the overall efficiency of CO₂ reduction. We describe herein the preparation of the new iridium complexes Ir₂(CO)₄(dmpm)₂ (1) (dmpm = bis(dimethylphosphino)methane) and Ir₂(CO)₃(dmpm)₂ (2) and the reaction of 2 with CO₂(l) to form the μ -carbonate complex Ir₂(μ -CO₃)(CO)₂(dmpm)₂ (3). The carbonate ligand of complex 3 shows unprecedented oxygen atom transfer reactivity and has been found to undergo rapid and reversible oxygen atom transfer to CO.

The reaction of Ir₂(CO)₄(dmpm)₂ (1)²⁷ with Me₃NO produces





Figure 1. ORTEP drawing of Ir₂(CO)₃(CO)₂(dmpm)₂ (3). Selected bond distances and angles: Ir(1)–Ir(2), 2.965 (1) Å; Ir(1)–O(31), 2.08 (1) Å; Ir(2)–O(32), 2.09 (1) Å; O(31)–C(30), 1.29 (3) Å; O(32)–C(30), 1.29 (3) Å; C(30)–O(33), 1.27 (3) Å; Ir–P(av), 2.311 (6) Å; O(31)–C(30)–O(32), 123 (2)°; O(31)–C(30)–O(33), 118 (2)°; O(32)–C(30)–C(33), 119 (2)°; P(11)–Ir(1)–P(12), 176.4 (2)°; P(21)–Ir(2)–P(22), 171.4 (2)°; O(32)–Ir(2)–C(21), 177.8 (9)°; O(31)–Ir(1)–C(11), 176.4 (7)°.

1 equiv of CO₂ and Ir₂(CO)₃(dmpm)₂ (2).²⁸ Complex 2 exhibits $\nu(\text{CO})$ bands at 1902 (s) and 1846 cm⁻¹, indicative of terminal and semibridging CO ligands, respectively. The presence of two $\nu(\text{P}=\text{C})$ dmpm bands at 942 (m) and 929 (m) cm⁻¹ is a signature of a cis,cis or cis,trans (dmpm)₂ conformation.²⁹ The variable-temperature ³¹P{¹H} NMR spectrum of 2 shows a single resonance at -42.9 ppm at 25 °C, coalescence at -10 °C, and an AA'XX' multiplet extending from -22.3 to -61.4 ppm (*J*_{PAP_x} = 74.3 Hz) at -70 °C. The exchange rate at -10 °C is calculated to be extremely rapid, 5000 s⁻¹, with a free energy of coalescence of 11 kcal/mol.^{31,32} The dynamic ³¹P{¹H} data together with IR evidence in the $\nu(\text{P}=\text{C})$ region establish a cis,trans diphosphine conformation for 2. These data suggest that 2 possesses the "non-A-frame" M₂L₇ geometry, analogous to the structurally characterized complex Rh₂(CO)₃(dppm)₂.³⁰



2

[†] Address correspondence pertaining to crystallographic studies to this author.

- (1) Research Fellow of the Alfred P. Sloan Foundation, 1987–1989.
- (2) Eisenberg, R.; Hendriksen, D. E. *Adv. Catal.* **1979**, *28*, 79.
- (3) Palmer, D. A.; van Eldik, R. *Chem. Rev.* **1983**, *83*, 651.
- (4) Darensbourg, D. J.; Kudarski, R. A. *Adv. Organomet. Chem.* **1983**, *22*, 129.
- (5) Inoue, S.; Koinuma, H. *Rev. Inorg. Chem.* **1984**, *6*, 291.
- (6) Lundquist, E. G.; Huffman, J. C.; Caulton, K. G. *J. Am. Chem. Soc.* **1986**, *108*, 8309.
- (7) Lee, G. R.; Cooper, N. J. *Organometallics* **1985**, *4*, 1467.
- (8) DeLaet, D. L.; del Rosario, R.; Fanwick, P. E.; Kubiak, C. P. *J. Am. Chem. Soc.* **1987**, *109*, 754.
- (9) Wu, J.; Fanwick, P. E.; Kubiak, C. P. *Organometallics* **1987**, *6*, 1805.
- (10) Calabrese, J. C.; Herskovitz, T.; Kinney, J. B. *J. Am. Chem. Soc.* **1983**, *105*, 5914.
- (11) Lee, G. R.; Maher, J. M.; Cooper, N. J. *J. Am. Chem. Soc.* **1987**, *109*, 2956.
- (12) Evans, G. O.; Walter, W. F.; Mills, D. R.; Streit, C. A. *J. Organomet. Chem.* **1978**, *144*, C34.
- (13) Maher, J. M.; Cooper, N. J. *J. Am. Chem. Soc.* **1980**, *102*, 7606.
- (14) Chatt, J.; Kubota, M.; Leigh, G. J.; March, T. C.; Mason, R.; Yarrow, D. J. *J. Chem. Soc., Chem. Commun.* **1974**, 1033.
- (15) Bryan, J. C.; Geib, S. J.; Rheingold, A. L.; Meyer, J. M. *J. Am. Chem. Soc.* **1987**, *109*, 2826.
- (16) Fachinetti, G.; Floriani, C.; Chiesi-Villa, A.; Guastini, C. *J. Am. Chem. Soc.* **1979**, *101*, 1767.
- (17) Nicholas, K. M. *J. Organomet. Chem.* **1980**, *188*, C10.
- (18) Maher, J. M.; Lee, G. R.; Cooper, N. J. *J. Am. Chem. Soc.* **1982**, *104*, 6797.
- (19) Lee, G. R.; Cooper, N. J. *Organometallics* **1985**, *4*, 1467.
- (20) Fisher, B. J.; Eisenberg, R. *J. Am. Chem. Soc.* **1980**, *102*, 7361.
- (21) Bruce, M. R. M.; Megehee, E.; Sullivan, B. P.; Thorp, H.; O'Toole, T. R.; Downard, A.; Meyer, T. J. *Organometallics* **1988**, *7*, 238 and references therein.
- (22) Frese, K. W.; Summers, D. P.; Cinibulk, M. J. *Electrochem. Soc.* **1988**, *135*, 264.
- (23) DeLaet, D. L.; Fanwick, P. E.; Kubiak, C. P. *J. Chem. Soc., Chem. Commun.* **1987**, 1412.
- (24) Hawecker, J.; Lehn, J.-M.; Zeissel, R. *J. Chem. Soc., Chem. Commun.* **1983**, 536.
- (25) Lehn, J.-M.; Zeissel, R. *Proc. Natl. Acad. Sci. U.S.A.* **1982**, *79*, 701.
- (26) Kutal, C.; Corbin, A. J.; Ferraudi, G. *Organometallics* **1987**, *6*, 553.
- (27) Belmore, K. A.; Vanderpool, R. A.; Tsai, J.-C.; Khan, M. A.; Nicholas, K. M. *J. Am. Chem. Soc.* **1988**, *110*, 2004.
- (28) Lemke, F. R.; DaLaet, D. L.; Gao, J.; Kubiak, C. P. *J. Am. Chem. Soc.* **1988**, *110*, 6904.

(27) Complex 1 was prepared from [Ir(COD)Cl]₂ (1.41 g, 2.10 mmol) and dmpm (0.57 g, 4.20 mmol) at 0 °C in toluene. The resulting cream-colored slurry was stirred for 5 min at which time CO was added, causing the mixture to turn dark red. The reaction was warmed to room temperature and the solution was stirred 2 h, resulting in a pale yellow solution and a light yellow solid. This product was reduced in situ over an excess of Na/Hg under an atmosphere of CO for 5 days. The yellow solution was filtered and reduced in volume. Precipitation of yellow crystalline material was accomplished by slow addition of hexanes to give 1.34 g (83% yield) of Ir₂(CO)₄(dmpm)₂ (1). Anal. Calcd for C₁₄H₂₈P₄O₄Ir₂: C, 21.87; H, 3.68. Found: C, 21.94; H, 3.64. ¹H NMR (CD₃CN): δ 3.46 (m, 4 H), δ 1.80 (s, 24 H). ³¹P{¹H} NMR (CD₃CN): δ -65.4 (s). IR (CH₂Cl₂): $\nu(\text{CO})$ 1956 (m), 1930 (s), 1890 (s), 1865 (w). Complex 1 was also characterized by a single-crystal X-ray diffraction study, the results of which will be reported separately.


(28) Complex 2 was prepared from 1 (0.025 g, 0.32 mmol) and Me₃NO (0.024 g, 0.32 mmol) at 70 °C in toluene. The solvent was removed to give 0.20 g (83% yield) of red-brown Ir₂(CO)₃(dmpm)₂ (2). Anal. Calcd for C₁₃H₂₈P₄O₃Ir₂: C, 21.08; H, 3.82. Found: C, 21.41; H, 4.06. ¹H NMR (C₆D₆): δ 2.63 (m, 4 H), 1.45 (s, 24 H). ³¹P{¹H} NMR (C₆D₆): δ -42.9 (s). IR (KBr): $\nu(\text{CO})$ 1902 (vs), 1846 (m); $\nu(\text{P}=\text{C})$ 942 (m), 929 (m).

(29) (a) Wu, J.; Fanwick, P. E.; Kubiak, C. P. *J. Am. Chem. Soc.* **1988**, *110*, 1319. (b) Wu, J.; Fanwick, P. E.; Kubiak, C. P. *J. Am. Chem. Soc.*, in press.

(30) Woodcock, C.; Eisenberg, R. *Inorg. Chem.* **1985**, *24*, 1287.


(31) Binsch, G.; Kessler, H. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 411.

(32) Sandstrom, J. *Dynamic NMR Spectroscopy*; Academic Press: New York, 1982.

Scheme 1. Reversible Carbonate Oxygen Atom Transfer from **3** to CO

Complex **2** reacts with both CO_2 and CO. The reaction of **2** with liquid CO_2 affords a mixture of **1** and the new carbonate complex $\text{Ir}_2(\text{CO}_3)(\text{CO})_2(\text{dmpm})_2$ (**3**).³⁶ The reaction of **2** with 1 equiv of CO leads back to **1**. The structure of **3** was determined by single-crystal X-ray diffraction.³⁷ An ORTEP drawing of **3** together with selected bond angles and distances appears in Figure 1. Complex **3** possesses an essentially planar bridging carbonate, torsionally canted at 20° with respect to the Ir–Ir vector. The Ir–Ir separation of 2.965 (1) Å, deep purple color, and overall structure are consistent with a weakly Ir...Ir interacting, d^8 – d^8 face-to-face bi-square-planar system.

The formation of **3** and **1** in the reaction of **2** with $\text{CO}_2(\text{l})$ appears to result from reductive disproportionation of 2CO_2 to CO_3^{2-} and CO (eq 2). Isolated yields of **3** are typically $\sim 40\%$,



indicating that some of the CO produced by CO_2 disproportionation is lost to the vapor phase in equilibrium with CO_2 liquid under reaction conditions.


The reaction of **2** with $^{13}\text{CO}_2$ (99% ^{13}C) leads to a surprising result. Comparison of the IR bands of **3** produced from $^{13}\text{CO}_2$ with those of **3** from CO_2 (natural abundance) reveals new $\nu(^{13}\text{CO})$ at 1896, 1878, 1523, and 1226 cm^{-1} . The disproportionation of $^{13}\text{CO}_2$ to ^{13}CO and $^{13}\text{CO}_3$ was verified by $^{13}\text{C}\{^1\text{H}\}$ NMR. The sample of **3** prepared from $^{13}\text{CO}_2$ showed $^{13}\text{C}\{^1\text{H}\}$ NMR signals at δ 177 (^{13}CO) and δ 164 ($^{13}\text{CO}_3$). However, there are also

Table I. Comparison of IR Data for **3** Prepared from CO_2 vs $^{13}\text{CO}_2$ (cm^{-1})



3 (CO_2)	3 ($^{13}\text{CO}_2$)	assignment ³⁸
1942 (vs)	1942 (vs)	$\nu(^{12}\text{CO})$
	1896 (m)	$\nu(^{13}\text{CO})$
1920 (sh)	1920 (w)	$\nu(^{12}\text{CO})$
	1878 (m)	$\nu(^{13}\text{CO})$
1561 (vs)	1561 (s)	$\nu(^{12}\text{CO}_3)$
	1523 (s)	$\nu(^{13}\text{CO}_3)$
1257 (m)	1257 (m)	$\nu(^{12}\text{CO}_3)$
	1226 (m)	$\nu(^{13}\text{CO}_3)$

intense $\nu(^{12}\text{CO}_3)$ bands evident at 1561 (s) and 1257 (m) cm^{-1} (Table I).³⁸ The key observation from this labeling experiment is that a significant portion of the CO_3^{2-} groups contain ^{12}C ! This suggests that CO from **2** is used to produce some of the CO_3^{2-} . The significance of this result lies in the fact that the products are not consistent with simple disproportionation of two $^{13}\text{CO}_2$ molecules to produce ^{13}CO and $^{13}\text{CO}_3^{2-}$.

Our results suggest that the $\mu\text{-CO}_3$ group of **3** undergoes reversible oxygen atom transfers with CO. This conclusion is supported by additional experimental evidence. A mixture of **1** and **3** reacts at 25°C in acetonitrile to give **2** and CO_2 , establishing the reversibility of eq 2. Complex **3** reacts rapidly with CO to give **1** and CO_2 (eq 3).



The presence of $^{12}\text{CO}_3^{2-}$ in the reaction of **2** with $^{13}\text{CO}_2$ therefore is explained by reverse oxygen atom transfer from co-ordinated $^{13}\text{CO}_3^{2-}$ to ^{12}CO to produce $^{13}\text{CO}_2$ and $^{12}\text{CO}_2$, either of which can disproportionate. Indeed, in the reaction of **3** with ^{13}CO (99% ^{13}C), the products are 50% $^{13}\text{CO}_2$, 50% $^{12}\text{CO}_2$, as determined by GC/MS, and **1** (50% ^{13}CO) (eq 4). The insertion



of CO into the $\mu\text{-CO}_3$ Ir–O bond of **3** to afford a C_2O_4 intermediate is implied by the oxygen atom transfer to CO. The fact that equal $^{13}\text{CO}_2/^{12}\text{CO}_2$ isotope partitioning is observed in the reaction of **3** with ^{13}CO (eq 4), suggests that the C_2O_4 intermediate rapidly collapses to 2 equiv of CO_2 (Scheme 1).³⁹ The reverse

(33) Haymore, B. L.; Ibers, J. A. *J. Am. Chem. Soc.* **1974**, *96*, 3325.
(34) Feltham, R. D.; Kriege, J. C. *J. Am. Chem. Soc.* **1979**, *101*, 5064.
Doughty, D. T.; Gordon, G.; Stewart, R. P., Jr. *J. Am. Chem. Soc.* **1979**, *101*, 2645.

(35) Lawson, H. J.; Atwood, J. D. *J. Am. Chem. Soc.* **1988**, *110*, 3680.

(36) Complex **3** was prepared from **2** and $\text{CO}_2(\text{l})$ in a LECO-MRA-114R pressure reactor at 25°C . Typically, 3 L of CO_2 gas was condensed into a 5-mL reactor containing 0.12 g (0.16 mmol) of **2**, resulting in an average pressure of 1700 psi. The mixture was vented and then opened in an inert-atmosphere box to collect solid products. Solids were washed with ~ 10 mL of Et_2O followed by ~ 1 mL of toluene to remove unreacted **2** and **1**, formed as a coproduct. $\text{Ir}_2(\text{CO}_3)(\text{CO})_2(\text{dmpm})_2$ (**3**) was dried under vacuum to yield 0.051 g (41%) of purple solids. X-ray quality crystals of **3** were obtained by the diffusion of Et_2O into an acetonitrile solution of **3**. Anal. Calcd for $\text{C}_{13}\text{H}_{28}\text{P}_4\text{O}_5\text{Ir}_2$: C, 20.21; H, 3.66. Found: C, 19.53; H, 4.16. ^1H NMR (CD_3CN): δ 3.32 (m, $J_{\text{PH}} = 4.0$ Hz, 4 H), 1.76 (s, 24 H). $^{31}\text{P}\{^1\text{H}\}$ NMR (CD_3CN): δ -6.2 (s). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_3CN): δ 177 (s, ^{13}CO), 164 (s, $^{13}\text{CO}_3$). IR (KBr): $\nu(\text{CO})$ 1942 (s), 1920 (sh), 1602 (sh), 1561 (s), 1257 (m); $\nu(\text{P-C})$ 941 (s).

(37) Crystal data for **3**·MeCN: $\text{Ir}_2\text{P}_4\text{O}_5\text{NC}_5\text{H}_3$, fw = 813.71, monoclinic, space group $P2_1/n$ (No. 14), $a = 11.823$ (2) Å, $b = 14.166$ (3) Å, $c = 14.807$ (4) Å, $\beta = 96.13$ (2)°, $V = 2465$ (2) Å³, $Z = 4$, $d_{\text{calc}} = 2.192$ g cm⁻³. The structure was solved by MULTAN least-squares Fourier methods and was refined to R and R_w values of 0.041 and 0.047 for 214 variables and 1875 unique observations with $I > 3\sigma(I)$ with Mo $K\alpha$ radiation. Data were corrected for absorption empirically.

(38) The $\nu(^{13}\text{CO})$ and $\nu(^{12}\text{CO})$ bands of CO and CO_3^{2-} ligands in complex **3** do not appear to be mixed appreciably. The relative intensities of all $\nu(^{13}\text{CO})$ and $\nu(^{12}\text{CO})$ bands in the IR spectrum of **3** are directly proportional to the total $^{13}\text{C}/^{12}\text{C}$ ratio. Moreover, $^{13}\text{C}\{^1\text{H}\}$ NMR verifies incorporation of ^{13}C into both CO and CO_3^{2-} .

(39) A $\mu\text{-C}(\text{O})\text{OC}(\text{O})$ intermediate formed by CO_2 insertion with **2**¹⁸ is also possible but is not consistent with the formation of carbonate. An Ir_2 -($\mu\text{-O}$) intermediate⁴⁰ has been considered but eliminated on the grounds that the $\mu\text{-CO}_3^{2-}$ carbon of **3** does not equilibrate with $^{13}\text{CO}_2$.

process, head-to-tail dimerization of CO₂, has often been invoked in the disproportionation of CO₂. Oxygen atom transfer to CO from CO₂,¹¹ NO,³³ NO₂,³⁴ and O₂,³⁵ has been reported. The present system is the first to display facile oxygen atom transfer from carbonate. Our studies of oxygen atom transfer from **3** to other substrates are continuing.

Acknowledgment. This work was supported by the NSF (Grant CHE-8707963). We are also grateful to the NSF for support of the Chemical X-ray Diffraction Facility at Purdue. A loan of IrCl₃ from Johnson-Matthey, Inc., is gratefully acknowledged.

Supplementary Material Available: Tables consisting of crystal data and data collection parameters (Table 1), positional parameters (Table 2), temperature factor expressions (Table 3), bond distances (Table 4), and bond angles (Table 5) for **3** (10 pages); a table of observed and calculated structure factors for **3** (14 pages). Ordering information is given on any current masthead page.

(40) Sharp, P. R.; Flynn, J. R. *Inorg. Chem.* **1987**, *26*, 3231.

Total Synthesis and Evaluation of **(±)-N-(tert-Butyloxycarbonyl)-CBI, (±)-CBI-CDPI₁, and (±)-CBI-CDPI₂: CC-1065 Functional Agents** **Incorporating the Equivalent** **1,2,9,9a-Tetrahydrocycloprop[1,2-c]benz[1,2-e]indol-4-one (CBI) Left-Hand Subunit**

Dale L. Boger,^{*,1a} Takayoshi Ishizaki,^{1b}
Ronald J. Wysocki, Jr., and Stephen A. Munk^{1c}


Department of Chemistry, Purdue University
West Lafayette, Indiana 47907

Paul A. Kitos and Oranart Suntornwat

Department of Biochemistry, University of Kansas
Lawrence, Kansas 66045-2500
Received March 30, 1989

(+)-CC-1065 (**1**, NSC-298223), an antitumor antibiotic isolated from cultures of *Streptomyces zelensis*, possesses exceptionally potent in vitro cytotoxic activity, broad spectrum antimicrobial activity, and confirmed in vivo antitumor activity.²⁻³ In a series of extensive investigations the site and mechanism of the (+)-CC-1065 antitumor activity have been related to its irreversible covalent alkylation of sequence-selective B-DNA minor groove sites [5'-d(A/GNTTA)-3' and 5'-d(AAAAA)-3'] that has been demonstrated to proceed by 3'-adenine N-3 alkylation of the electrophilic cyclopropane present in the left-hand (CPI) subunit of (+)-CC-1065.^{4,5} The demonstration that simplified agents including CDPI₃ methyl ester exhibit a substantial preference for A-T rich noncovalent minor groove binding⁷ attributable to preferential stabilization of a noncovalent complex within the

Scheme I



Scheme II^a

