

Generation and reaction of carbenes from α -oxacyclo-*N*-aziridinyl imines—a new method for ring expansion of cyclic ethers

Sunggak Kim,* Joo-Yong Yoon and Chang Mook Cho


Department of Chemistry, Korea Advanced Institute of Science and Technology, Taejeon 305-701, Korea

α -Oxetanyl-*N*-aziridinyl imines undergo ring expansion on heating to afford dihydrofurans *via* alkylidenecarbenes, whereas heating α -tetrahydrofuranyl and α -tetrahydropyranyl *N*-aziridinyl imines affords the ring expanded cyclic enol ethers *via* alkyl carbenes.

Since Eschenmoser reported that *N*-aziridinyl imines of α,β -epoxy ketones underwent thermal fragmentation,¹ *N*-aziridinyl imines have been utilized as precursors of both diazoalkanes² and carbenes.³ In connection with our interest in the synthetic utility of *N*-aziridinyl imines,⁴ we recently reported that the thermal reaction of α,β -epoxy-*N*-aziridinyl imines in refluxing toluene generates (β -hydroxyalkylidene)carbenes, which underwent 1,5-C–H insertion and 1,5-O–Si insertion to afford cyclopentenols⁵ and dihydrofurans.⁶ As an extension to this work, we were interested in the generation and reaction of carbenes from α -oxacyclo-*N*-aziridinyl imines.

The reaction of oxetanes with alkoxycarbonylcarbenes has been previously studied and the reaction was known to proceed *via* competitive C–H insertion and oxonium ylide formation.⁷ However, with tetrahydrofuran, carbenes reacted to give only C–H insertion products.⁸ We turned our attention to the question of whether (i) α -oxacyclo-*N*-aziridinyl imines would generate alkylcarbenes **2** or alkylidenecarbenes **4** (path a) and (ii) alkylcarbene intermediates would involve C–C insertion to afford **5** (path b) or oxonium ylide formation to afford **6** (path c), Scheme 1.

When the thermal reaction of **1b** was carried out in refluxing toluene for 5 h, we were somewhat surprised to find that the expected **6b** was not obtained. Compound **5b** was instead isolated in 68% yield. Apparently, the reaction did not proceed *via* the oxonium ylide **3b** or alkylidenecarbene **4b** from the structure of the product **5b**. Thus, the most reasonable explanation for the formation of **5b** would involve the direct insertion of alkylcarbene **2b** into C–C bond.⁹ The same result was also obtained with α -tetrahydropyranyl-*N*-aziridinyl imine **1c**. For the thermal reaction of bicyclic ether **7**, in the carbene intermediate **8**, migration of bond *a* would be expected to afford **9**. Alternatively, migration of bond *b* is unlikely due to the formation of the bridged product **10**, eqn. (1). Thus, the thermal reaction of **7** in refluxing toluene for 7 h afforded **9** in 72% yield. Further examples are shown in Table 1.



We next examined the thermal reaction of α -oxetanyl-*N*-aziridinyl imines. When the thermal reaction of **1a** was carried out in refluxing toluene for 8 h, **6a** was isolated in 59% yield, indicating that the reaction did not undergo C–C insertion *via*






Table 1 Ring expansion of cyclic ethers

Entry	Substrate ^a	t/h	Product (%) ^b
1		5	(60)
2		8	(62)
3		5	(85)
4		5	(91)
5		5	(59)
6		8	(58)
7		5	(69)

^a All substrates are a mixture of *syn*- and *anti*- isomers. ^b The yield refers to the isolated yield. ^c A = 2-phenylaziridinyl.

alkylcarbene **2a**. Since the thermal reaction of **1b** and **1c** did not involve the formation of oxonium ylides **3b** and **3c**, the reaction seems to proceed *via* alkylidenecarbene **4a**, indicating the dependence of the mode of the reaction on the ring size of cyclic ethers. Our previous results on the thermal reaction of α,β -epoxy-*N*-aziridinyl imines also support the intermediacy of **4a**.⁵ For most of the cases observed, the reaction was complete within 8 h in refluxing toluene affording dihydrofurans as shown in Table 1. The reaction was also successful with bicyclic oxetanes.

To study the relative ease of the two processes, 1,2-H migration and direct ring expansion into the C–C bond in alkylcarbenes, we examined the thermal reaction of **11**, eqn. (2). The thermal reaction of **11** in refluxing toluene for 8 h gave only pyran **14** in 84% yield as a result of selective 1,2-H migration in



the alkyl carbene **12** and the following isomerization of the double bond. However, it is noteworthy that the 1,5-OH insertion in alkylidenecarbenes proceeded exclusively without 1,2-H migration. Thus, the thermal reaction of **15** in refluxing toluene for 8 h afforded **17** in 55% yield, eqn. (3).

We are grateful to the Korea Science and Engineering Foundation and the Organic Chemistry Research Center for financial support of our research programme.

References

- 1 D. Felix, R. K. Mueller, U. Horn, R. Joos, J. Schreiber and A. Eschenmoser, *Helv. Chim. Acta*, 1972, **55**, 1276.
- 2 A. G. Schultz, J. P. Dittami and K. K. Eng, *Tetrahedron Lett.*, 1984, **25**, 1255; G. B. Jones, C. J. Moody, A. Padwa and J. M. Kassir, *J. Chem. Soc., Perkin Trans. I*, 1991, 1721.
- 3 A. Padwa, Y. Gareau and S. L. Xu, *Tetrahedron Lett.*, 1991, **32**, 983; N. Chen, M. Jones, W. R. White and M. S. Platz, *J. Am. Chem. Soc.*, 1991, **113**, 4981; A. Padwa, D. J. Austin, Y. Gareau, J. M. Kassir and S. L. Xu, *J. Am. Chem. Soc.*, 1993, **115**, 2637.
- 4 For use of *N*-aziridinyl imines in radical reactions, S. Kim, I. S. Kee and S. Lee, *J. Am. Chem. Soc.*, 1991, **113**, 9882; S. Kim and I. S. Kee, *Tetrahedron Lett.*, 1993, **34**, 4213.
- 5 S. Kim and C. M. Cho, *Tetrahedron Lett.*, 1994, **35**, 8405.
- 6 S. Kim and C. M. Cho, *Tetrahedron Lett.*, 1995, **36**, 4845.
- 7 H. Nozaki, H. Takaya and R. Noyori, *Tetrahedron*, 1966, **22**, 3393; W. Kirmse, R. Lelgemann and K. Friedrich, *Chem. Ber.*, 1991, **124**, 1853.
- 8 W. E. Doering, L. H. Knox and M. Jones, *J. Org. Chem.*, 1959, **24**, 136.
- 9 P. Yates and S. Danishefsky, *J. Am. Chem. Soc.*, 1962, **84**, 879; J. W. Wilt, J. M. Kosturik and R. C. Orlowski, *J. Org. Chem.*, 1965, **30**, 1052; J. W. Wilt, J. F. Zawadzki and D. G. Schultenover, *J. Org. Chem.*, 1966, **31**, 876.

Received, 23rd January 1996; Com. 6/00543H