


equilibrating bicyclobutonium ions



equilibrating "bisected" cyclopropylcarbinyl cations

methylcyclopropylcarbinyl (7) < cyclopropylcarbinyl (6). All of the cyclopropylcarbinyl cations show delocalization of charge by the cyclopropyl ring, but only with the primary system do we observe the truly unique behavior which indicates the limiting nonclassical nature of the ion. Present work reinforces our contention that the  $^{13}\text{C}$  chemical shifts are an excellent, direct experimental way of determining the "non-classical" character of carbonium ions.

**Acknowledgment.** Support of this work by a grant of the National Science Foundation is gratefully acknowledged.

G. A. Olah, D. P. Kelly, C. L. Jeuell, R. D. Porter  
Department of Chemistry, Case Western Reserve University  
Cleveland, Ohio 44106  
Received October 2, 1969


### Stable Carbonium Ions. IC. The Benzenonium Ion ( $\text{C}_6\text{H}_7^+$ ) and Its Degenerate Rearrangement<sup>1</sup>

Sir:


Mass spectroscopic studies<sup>2</sup> have shown the formation of  $\text{C}_6\text{H}_7^+$  as a fragmentation ion produced upon electronic bombardment of benzyl alcohol. Substantial research has been reported on proton-addition complexes of aromatic hydrocarbons.<sup>3-12</sup> Olah and

Kuhn<sup>6,9</sup> achieved the synthesis and isolation of stable alkylbenzenonium fluoroborate and fluoroantimonate salts. Although Perkampus and Baumgarten<sup>13</sup> reported solid-state infrared and ultraviolet spectra for the ternary system benzene-HCl-GaCl<sub>3</sub>, a well-characterized description of  $\text{C}_6\text{H}_7^+$  in solution has not been presented until now.<sup>14</sup>


We wish to report the observation of the simple benzenonium ion ( $\text{C}_6\text{H}_7^+$ ) as a stable long-lived species. When a saturated solution of benzene in sulfuryl chloride fluoride at  $-50^\circ$  is added dropwise with vigorous stirring to a 1:2 (v/v) solution of 1:1 (mol/mol) hydrogen fluoride-antimony pentafluoride in  $\text{SO}_2\text{ClF}$  at  $-78^\circ$ , a suspension is formed. The clear supernatant solution has a pmr spectrum (Figure 1) consisting of a single sharp peak at  $\delta$  8.09 (apart from the acid and the  $\text{H}_3\text{O}^+$  peaks), which we assign to the degenerate  $\text{C}_6\text{H}_7^+$  ion 1a-c, undergoing rapid hydrogen migration.



1a



1b



1c

etc.

When the mixed solvent system  $\text{SO}_2\text{ClF}-\text{SO}_2\text{F}_2$  was employed,<sup>15</sup> the pmr spectrum could be observed at as low a temperature as  $-134^\circ$ . At this temperature the separate absorptions for the four different types of protons of the "frozen-out" benzenonium ion are apparent (cf. 1a): the methylene protons ( $\text{H}_1, \text{H}_2$ ) giving a broad signal at  $\delta$  5.84,  $\text{H}_4$  and  $\text{H}_6$  at 8.40,  $\text{H}_5$  at 9.20, and  $\text{H}_3$  and  $\text{H}_7$  at 9.42 ppm (Figure 2). Warming the solution to  $-70^\circ$  regenerates the singlet absorption at  $\delta$  8.09 ppm.

The temperature dependence of the spectrum was used to calculate the rate constants and activation parameters of the hydrogen migration. The line shapes for various rate constants were calculated and recorded by means of a multiple-site exchange program.<sup>16</sup> The axial ( $\text{C}_{2v}$ ) symmetry of the ion allowed us to treat this case as a four-site problem. Fitting the data to the Arrhenius equation gave an activation

(8) E. L. Mackor, A. Hofstra, and J. H. van der Walls, *Trans. Faraday Soc.*, **54**, 186 (1958).

(9) G. A. Olah, *J. Amer. Chem. Soc.*, **87**, 1103 (1965).

(10) C. MacLean and E. L. Mackor, *Discuss. Faraday Soc.*, **34**, 165 (1962).

(11) T. Birchall and R. J. Gillespie, *Can. J. Chem.*, **42**, 502 (1964).

(12) E. M. Arnett and J. W. Larsen, *J. Amer. Chem. Soc.*, **90**, 792 (1968).

(13) H. H. Perkampus and E. Baumgarten, *Angew. Chem. Int. Ed. Engl.*, **3**, 776 (1964).

(14) The species thought to be  $\text{C}_6\text{H}_7^+$  in  $\text{SO}_3$  solution of benzene-HF-SbF<sub>5</sub> was obviously the sulfinated product (*vide infra*).

(15) The solution (0.5 ml) is added to  $\text{SO}_2\text{F}_2$  (ca. 0.2 ml) in an nmr tube at  $-78^\circ$ .

(16) The Fortran IV program was written by C. S. Johnson, Jr., *Advan. Magnetic Resonance*, **1**, 33 (1965), and adapted for use on a Univac 1108 computer by Mr. W. Heyd of this department.

(1) Part XCVIII: G. A. Olah, D. P. Kelly, C. L. Jeuell, and R. D. Porter, *J. Amer. Chem. Soc.*, **92**, 2544 (1970).

(2) H. Budzikiewicz, C. Djerassi, and D. H. Williams, "Mass Spectroscopy of Organic Compounds," Holden-Day, San Francisco, Calif., 1967, p 119.

(3) V. Gold and F. L. Tye, *J. Chem. Soc.*, 2172 (1952).

(4) C. Reid, *J. Amer. Chem. Soc.*, **76**, 3264 (1954).

(5) D. A. McCauley and A. P. Lien, *ibid.*, **73**, 2013 (1951).

(6) G. A. Olah and S. J. Kuhn, *Nature*, **178**, 693 (1956); *J. Amer. Chem. Soc.*, **80**, 6535, 6541 (1958).

(7) W. von E. Doering, M. Saunders, H. G. Boyton, H. W. Earhart, E. F. Wadley, W. R. Edwards, and G. Taber, *Tetrahedron*, **4**, 178 (1958).





Figure 1. Spectrum (60 MHz) of  $C_6H_7^+$  in  $HF-SbF_5-SO_2ClF$  at  $-80^\circ$ .

energy,  $E_a$ , of  $10 \pm$  kcal mol $^{-1}$  and a preexponential factor,  $A$ , of  $10^{15.9 \pm 1.6}$ .

The broadness of the methylene absorption at  $\delta$  5.84 ppm can be explained by proton exchange with the acid solvent system, which is not completely stopped at the low temperature attained. The deuterated analog of ion **1** formed from benzene- $d_6$  in  $HF-SbF_5-SO_2ClF$  indeed undergoes slow deuterium-hydrogen exchange at  $-134^\circ$  which is much accelerated at higher temperatures. Broadening due to the nonequivalence of the aliphatic methylene protons in a 4- $\pi$ -electron Möbius-type<sup>17</sup> analog (**2**) of the homotropylium cation (a homo-Möbius cyclopentadienyl cation) seems un-



likely due to steric prohibition of a Möbius strip closure in a five-membered ring system.<sup>18</sup>

The singlet peak at  $\delta$  8.09 ppm is flanked by large  $^{13}C$  satellites (Figure 1), as would be expected for the degenerate ion (theoretically, each 3.3% of the  $^{12}C$  peak), giving a value of  $J(^{13}C-H)_{av} = 26$  Hz. This average value of a C-H coupling constant is a result of a rapid exchange involving all seven protons and six carbon atoms, similarly as observed by us in the case of the cyclopentyl cation.<sup>19</sup> Using the indor technique previously described<sup>19</sup> (monitoring the downfield  $^{13}CH$  satellite or the main  $^{12}CH$  peak) the  $^{13}C$  chemical shift of **1** at  $-78^\circ$  was found to be  $\delta(^{13}C) + 48$  ppm (from

(17) (a) E. Heilbronner, *Tetrahedron Lett.*, 1923 (1964); (b) H. E. Zimmerman, *J. Amer. Chem. Soc.*, **88**, 1564 (1966).

(18) Larger cyclic  $4n$   $\pi$ -electron cations like the cyclononatetraenyl cation could be stabilized by a Möbius-type interaction; our studies in this regard will be reported.

(19) A. M. White and G. A. Olah, *J. Amer. Chem. Soc.*, **91**, 2943 (1969).





Figure 2. Computed and observed spectra of degenerate (top) and "frozen-out" (bottom) benzenonium ion. Dashed line indicates computed methylene proton absorption in the absence of exchange with acid (see text).

$CS_2$ ). Both the average coupling constant and chemical shift values are in good agreement with those calculated for the rapidly equilibrating set of benzenonium ions ( $\sigma$  complexes) **1a-c** using either a static model, **3a-c**, or reported  $^{13}C$  data from the mesitylium ion (2,4,6-trimethylbenzenonium ion).<sup>20</sup>




From the static model,  $J(^{13}C-H)_{av}$  is calculated from four cyclohexadiene-type olefinic (167 Hz), two  $sp^3$ -type cyclohexane methylene (125 Hz), and one isopropyl cation-type (169 Hz)<sup>21</sup> couplings, equilibrated over six carbons as:  $J(^{13}C-H)_{av} = [(4 \times 167) + (2 \times 125) + (1 \times 169)]/(6 \times 7) = 26$  Hz.

Similarly the  $^{13}C$  chemical shift can be calculated from the values for olefinic carbons in cyclohexadiene, the methylene carbon in the 2,4,6-trimethylbenzenonium ion,<sup>20</sup> and the isopropyl cation carbon:<sup>19</sup>  $\delta(^{13}C)_{av} = [(4 \times 66) + (1 \times 139) + (1 \times -125)]/6 = +46$  ppm ( $CS_2$ ). The  $^{13}C$  chemical shifts for the 2,4,6-trimethylbenzenonium ion (**4**) have been reported by Koptug<sup>20</sup> and serve as an excellent model for **1**. Since the sub-


(20) V. Koptug, A. Rezvukhin, E. Lippmaa, and T. Pehk, *Tetrahedron Lett.*, 4009 (1968).

(21) G. A. Olah and A. M. White, *J. Amer. Chem. Soc.*, **91**, 5801 (1969).



stitution of a hydrogen atom for a methyl group in carbonium ions generally causes only a slight change (shielding of about 10 ppm) in the  $^{13}\text{C}$  shift,<sup>22</sup> the data from the 2,4,6-trimethylbenzenonium ion can be used directly to calculate the average  $^{13}\text{C}$  shift in 1:  $\delta(^{13}\text{C})_{\text{av}} = [139 + (3 \times -9.5) + (2 \times 58)]/6 = 47 \text{ ppm}$  ( $\text{CS}_2$ ).

When sulfur dioxide is added to the  $\text{HF-SbF}_5\text{-SO}_2\text{ClF}$  solution of 1 at  $-78^\circ$ , an immediate reaction occurs to form protonated benzenesulfonic acid,  $\text{C}_6\text{H}_5\text{-S(OH}_2^+)$ , as evidenced by a two-proton pmr singlet at  $\delta$  9.54 ppm (S-OH) and the five-proton aromatic absorption at 7.96 ppm.



Winstein's group<sup>23</sup> has reported that monoalkylbenzenes in  $\text{FSO}_3\text{H-SbF}_5\text{-SO}_2$  solution form sulfinated alkylbenzenes and suggested that the  $\text{SO}_2\text{-SbF}_5$  complex is responsible for the electrophilic sulfinations. This suggestion could explain the results of previous investigations of monoalkylbenzenonium ions in sulfur dioxide solutions,<sup>9</sup> i.e., formation of sulfinated products. Similar observations were made concerning ions produced from monoalkyl- and halobenzenes, which were obtained and studied in detail in super acid systems free of  $\text{SO}_2$  and diluted where necessary with  $\text{SO}_2\text{ClF}$ .<sup>24</sup> While the  $\text{SO}_2\text{-SbF}_5$  complex is an effective electrophilic sulfinating agent, we felt that in super acid media the reaction of the benzenonium ions themselves with  $\text{SO}_2$  can also be responsible for the observed sulfinations. In our previous studies the reaction of benzenonium ions as isolated salts with nucleophiles was studied. Ion 1 is not sufficiently stable for isolation, but its solution behavior supports its reactivity with  $\text{SO}_2$ . Alternatively there could be an equilibrium, even at  $-78^\circ$ , of 1 and benzene, with added  $\text{SO}_2$  promoting deprotonation. Benzene then could undergo sulfination by  $\text{SO}_2\text{-SbF}_5$  or  $\text{SO}_2\text{H}^+$ . At this low temperature sulfination would be expected to be slow and would hardly account for the experimental observation of fast quantitative conversion of 1 into benzenesulfonic acid. It should be mentioned that dialkylbenzenes and higher polyalkylbenzenes are sufficiently basic to form alkylbenzenonium ions which, at low temperatures, generally do not react with  $\text{SO}_2$ . Observation of these ions in  $\text{SO}_2$  solution was therefore not affected by the solvent.

(22) Substitution of a methyl by a hydrogen brings about a shielding of 3–10 ppm: (a) B. V. Cheney and D. M. Grant, *J. Amer. Chem. Soc.*, **89**, 5319 (1967); (b) D. M. Grant and E. G. Paul, *ibid.*, **86**, 2984 (1964); (c) L. M. Jackman and D. P. Kelly, *J. Chem. Soc., B*, in press.

(23) M. Brookhart, F. A. L. Anet, and S. Winstein, *J. Amer. Chem. Soc.*, **88**, 5657 (1966); M. Brookhart, Ph.D. Thesis, University of California, Los Angeles, Calif., 1968.

(24) G. A. Olah, C. U. Pittman, Jr., E. Namanworth, and M. B. Comisarow, *J. Amer. Chem. Soc.*, **88**, 5571 (1966); G. A. Olah and T. E. Kiovsky, *ibid.*, **89**, 5692 (1967); **90**, 2583 (1968).

Our studies indicate that we have observed the heretofore elusive benzenonium ion (1) in solution as an equilibrating set of degenerate ions which can be frozen out at  $-134^\circ$  to the static benzenonium ion. Our investigations relating to the benzenonium ion and monoalkylbenzenonium ions are published in full in a forthcoming paper.

**Acknowledgment.** Support of this work by the National Science Foundation and the C. F. Mabery Fund of Case Western Reserve University is gratefully acknowledged.

G. A. Olah, R. H. Schlosberg,  
D. P. Kelly, Gh. D. Mateescu

Department of Chemistry, Case Western Reserve University  
Cleveland, Ohio 44106

Received November 17, 1969

### Structure of the Methylcyclobutyl Cation

Sir:

Methylcyclopentyl cation was recently reported to undergo ready interchange of the  $\alpha$  and  $\beta$  ring protons and to exchange ring and methyl protons and carbons less readily.<sup>1</sup> The rates were similar to those found for analogous processes in the classical *t*-amyl cation.<sup>1,2</sup>

With the hope of studying similar processes in methylcyclobutyl cation, we treated the corresponding tertiary chloride I (prepared from methylenecyclobutane and  $\text{HCl}$  gas at  $0^\circ$ ) with  $\text{SbF}_5$  and  $\text{SO}_2\text{ClF}$  in the usual manner at low temperature. A solution giving an nmr ( $-80^\circ$ ) spectrum which consisted of two singlets at  $\tau$  6.11 and 7.13 (measured from external  $(\text{CH}_3)_4\text{Si}$ ) with an area ratio of 2:1 was obtained. Cooling a sample prepared with a  $\text{SO}_2\text{ClF-SO}_2\text{F}_2\text{-SbF}_5$  solvent system<sup>4</sup> to  $-125^\circ$  produced only broadening of both peaks, probably due to increased viscosity. Splitting was observed ( $-25^\circ$ ) appropriate for coupling between groups of three and six equivalent protons (Figure 1).

1-Chlorotrideuteriomethylcyclobutane (II) was prepared from trideuteriomethyl iodide (approximately 90% deuterium) via a Grignard reaction on cyclobutanone and subsequent treatment of the alcohol formed with concentrated  $\text{HCl}$  and pentane at room temperature. The ion from II was prepared and its nmr spectrum at  $-60^\circ$  was only a singlet (half-height width 1.7 Hz) at  $\tau$  6.07 and a small peak at 7.1 due to residual hydrogen from the methyl group of the chloride. Apparently the ring hydrogens are identical and 1 ppm downfield from the methyl position. The coupling constant (0.9 Hz) is unusually low compared to that for the coupling of methyl and  $\alpha$  hydrogens of the methylcyclopentyl cation (4.0 Hz).<sup>5</sup>

Several possible explanations might fit these results: (1) a series of rapid hydride and possibly methide shifts could exchange ring hydrogens; (2) the ion could have a static highly symmetrical structure with six equivalent methylene protons; (3) it could be a rapidly equilibrat-

(1) M. Saunders and J. Rosenfeld, *J. Amer. Chem. Soc.*, **91**, 7756 (1969).

(2) M. Saunders and E. L. Hagen, *ibid.*, **90**, 2436 (1968).

(3) H. C. Brown and M. Borkowski, *ibid.*, **74**, 1894 (1952); we are grateful to Gary Taylor for supplying the methylenecyclobutane.

(4) G. A. Olah and A. M. White, *ibid.*, **91**, 3956 (1969).

(5) G. A. Olah, J. M. Bollinger, C. A. Cupas, and J. Lukas, *ibid.*, **89**, 2692 (1967).