

Cite this: *Org. Biomol. Chem.*, 2014, **12**, 3238

Synthesis and evaluation of monophosphoryl lipid A derivatives as fully synthetic self-adjuvanting glycoconjugate cancer vaccine carriers†

Zhifang Zhou,‡ Mohabul Mondal,‡ Guochao Liao‡ and Zhongwu Guo*

A fully synthetic carbohydrate-based cancer vaccine is an attractive concept, but an important topic in the area is to develop proper vaccine carriers that can improve the immunogenicity and other immunological properties of tumor-associated carbohydrate antigens (TACAs). In this context, four monophosphoryl derivatives of *Neisseria meningitidis* lipid A were synthesized via a highly convergent and effective strategy and evaluated as vaccine carriers and adjuvants. The conjugates of these monophosphoryl lipid A (MPLA) derivatives with a modified form of the sTn antigen were found to elicit high titers of antigen-specific IgG antibodies, indicating a T cell-dependent immune response, in the absence of an external adjuvant. It was concluded that MPLAs could be utilized as potent vaccine carriers and built-in adjuvants to create fully synthetic self-adjuvanting carbohydrate-based cancer vaccines. The lipid composition and structure of MPLA were shown to have a significant influence on its immunological activity, and among the MPLAs examined, natural *N. meningitidis* MPLA exhibited the most promising properties. Moreover, Titermax Gold, a conventional vaccine adjuvant, was shown to inhibit, rather than promote, the immunological activity of MPLA conjugates, maybe via interacting with MPLA.

Received 20th February 2014,
Accepted 11th March 2014

DOI: 10.1039/c4ob00390j

www.rsc.org/obc

Introduction

Fully synthetic glycoconjugate cancer vaccines are currently a hot topic, since they have well-defined structures, reproducible physical, chemical and biological properties, and promising immunological activities.^{1–7} To develop functional conjugate cancer vaccines, a vital issue is the carrier molecule. An ideal vaccine carrier should be rather small to be synthetically manageable and highly immunoactive to be able to improve the immunogenicity and promote T cell-dependent immunity of tumor-associated carbohydrate antigens (TACAs) that are often poorly immunogenic and T cell-independent. While several vaccine carriers have been explored for this purpose,^{1–7} this paper presents a new type of vaccine carrier derived from lipid A for fully synthetic self-adjuvanting carbohydrate-based cancer vaccines.

Lipid A is the conserved hydrophobic core of lipopolysaccharides (LPSs) – the main component and virulent factor

on the Gram negative bacterial cell surface.⁸ Lipid A is of great importance in that it not only serves as an anchor to attach bacterial O-polysaccharides to the bacterial cell membrane⁸ but also is a potent toxin that can stimulate strong immune responses in eukaryotic hosts.⁹ It binds specifically to the toll or interleukin-1 receptor domain-containing adaptor-inducing interferon- β (TRIF) of toll-like receptors (TLRs) in association with MD88 to induce a downstream signaling cascade, stimulate the release of cytokines and chemokines, such as tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), IL-6 and interferon- β (IFN- β), and upgrade immune cell expression.¹⁰ Even though the immunostimulatory activity of lipid A can be extremely useful, its pro-inflammatory and septic properties are a serious problem.¹¹

Regardless of the source, all lipid As contain the highly conserved construct **1** (Fig. 1) with several lipids, varying considerably in chain length, saturation, number and distribution, linked to the N-2-, N-2'-, O-3- and O-3'-positions and two phosphate groups linked to the O-1- and O-4'-positions of the β -1,6-linked disaccharide of D-glucosamine (GlcNH₂). Biological assays demonstrated that the number, structure and position of lipids and the degree of phosphorylation are important factors determining the bioactivity of lipid A.^{12–14} A typical lipid A that exhibits a full spectrum of immunological and endotoxic activities consists of six lipids that can be distributed symmetrically (3 + 3), such as in the lipid A of *Neisseria*

Department of Chemistry, Wayne State University, 5101 Cass Avenue, Detroit, MI 48202, USA. E-mail: zwguo@chem.wayne.edu; Fax: +1-313-577-5060; Tel: +1-313-577-2557

†Electronic supplementary information (ESI) available: Experimental, ¹H, ¹³C, and 2D NMR spectra and MS spectra of all new compounds. See DOI: 10.1039/c4ob00390j

‡These authors contributed equally to the present paper.





Fig. 1 The conserved construct of lipid A (1) and the structures of *N. meningitidis* lipid A (2) and designed MPLA derivatives and their sTn conjugates (3–6).

meningitidis (2, Fig. 1),¹⁵ or asymmetrically (4 + 2), such as in the lipid A of *Escherichia coli*, on the two GlcNH₂ units. The two phosphate groups are essential for the endotoxic activity of lipid A. However, modification of the O-1-position, for example, removal of the phosphate group at this position to generate monophosphoryl lipid A (MPLA), was shown to greatly reduce its endotoxicity with little impact on its immunostimulatory activity.¹⁶ Thus, MPLA has emerged as a potential adjuvant used in human vaccine formulation.^{17,18} However, the optimal lipid A structure required for binding to the cell surface receptors is different from the structure required for the activation of the host immune system, supported by the finding that most biosynthetic precursors of lipid A are nontoxic and that the toxicity of lipid A has multiple structural requirements.¹⁹ Consequently, a systematic study on the structure–activity relationship of MPLA is necessary to discover proper vaccine carriers and adjuvants.

In this regard, many lipid A and MPLA derivatives have been prepared and evaluated in the literature.^{13,14,17,20–31} In association with our efforts to develop fully synthetic carbohydrate-based cancer vaccines, we synthesized a monophosphoryl analog³² of *N. meningitidis* lipid A and coupled it to *N*-phenylacetyl GM3, a modified form of TACA.³³ The resulting conjugate alone induced promising immune responses in animals in the absence of any external adjuvant, suggesting that, as an immunostimulant, MPLA was functional not only as a vaccine carrier but also as a built-in adjuvant.³⁴ Inspired by this discovery, here we have designed and synthesized three other monophosphoryl analogs of the lipid A of *N. meningitidis*, containing lipids of different chain lengths and linkages as well as different patterns of lipidation (Fig. 1). To investigate the immunological properties of these MPLA derivatives as vaccine carriers and adjuvants, we conjugated them with *N*-phenylacetyl sTn (sTnNPhAc), a modified TACA that is currently studied for anticancer vaccine development,³⁵ and examined in mice the specific immune responses induced by the resulting MPLA-sTnNPhAc conjugates 3–6 (Fig. 1). Through these studies, we anticipated to identify novel vaccine carriers and adjuvants with improved immunological properties useful for the development of fully synthetic glycoconjugate cancer vaccines.




Scheme 1 The retrosynthesis for 3–6.

Results and discussion

In our synthetic design for target molecules 3–6 (Scheme 1), the key intermediates were fully protected MPLA derivatives 7–10, which had a 2-azidoethyl group at the reducing end of the disaccharide backbone. Global deprotection of 7–10 would afford free MPLA derivatives that can be explored as independent vaccine adjuvants. On the other hand, the azido group in 7–10 could be selectively reduced to form a primary amine that would facilitate conjugation with sTnNPhAc or other TACAs to generate conjugate vaccines. In return, the MPLA derivatives would be constructed from a common disaccharide intermediate 11 having the 2,2'-*N*- and 3,3'-*O*-positions differently protected from other positions to enable regioselective deprotection and acylation later on. Moreover, the 4',6'-*O*-positions in 11 were protected with a benzylidene group that can be regioselectively opened to expose the 4'-*O*-position for the introduction of the conserved phosphoryl group.

The synthesis started with the preparation of properly protected D-glucosamine derivatives 12 and 13 (Scheme 2), employed as a glycosyl donor and acceptor, respectively, according to reported procedures.³² Glycosylation of 12 with 13 was realized with *N*-iodosuccinimide (NIS) and silver triflate (AgOTf) as promoters, and was unexpectedly slow and took two days to complete. The doublet signal of H-1' at δ 5.88 in the ¹H NMR spectrum of the resulting disaccharide 11 gave a coupling constant ($J_{1',2'}$) of 8.8 Hz, which confirmed the β -conformation of the newly generated glycosidic linkage. Treatment of 11 with hydrazine hydrate in refluxing ethanol removed both *N*-phthalyl and *O*-acetyl groups simultaneously to form



Scheme 2 Synthesis of the key intermediates 7–10.


14, which was ready for the introduction of lipids. In the meantime, the chiral lipids **15** and **18–20** (Scheme 2) required for these syntheses were also prepared according to reported procedures³² with commercial (*R*)-epichlorohydrin as the starting material.

For lipid installation (Scheme 2), the free amino groups in **14** were acylated first by reacting with **15** under the influence of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI). Despite the presence of two free hydroxyl groups in **14**, the reaction was fast and selective for the amino groups, probably owing to their higher reactivity, to produce **16** in an excellent yield (94%). The two NH signals as doublets at δ 6.19 (*J* 4.8 Hz) and 5.99 (*J* 6.8 Hz) as well as the down-field shift of the H-2 and H-2' signals in the ¹H NMR spectrum of **16** confirmed the desired N-acylation. Compound **16** was employed as a common intermediate for the synthesis of all MPLA derivatives designed. Acylation of the 3-O- and 3'-O-positions in **16** with lauric acid (**17**) and EDCI required DMAP as a catalyst to afford **21**. It was observed that the reaction to introduce the last lipid chain into either the 3-O- or the 3'-O-position was very slow and could be complete only after prolonged reaction time, probably due to the further increased steric hindrance after introducing the third lipid into one of these two positions. Similarly, the coupling reactions between **16** and fatty acids **18–20** in the presence of EDCI and DMAP generated **22–24**, respectively. Regioselective opening of the benzylidene ring in **21–24** was accomplished using NaBH₃CN under acidic conditions to result in **25–28**, which had a free hydroxyl group at the 4-O'-position. The regiochemistry of these intermediates was confirmed by the observation of a down-field shift of 4'-H in the ¹H NMR spectra of the acetylation products as well as by a similar down-field shift of the 4'-H observed in the O-phosphoryl products of the next step. Phosphorylation of **25–28** was achieved by the phosphoramidite approach in a two-step one-pot manner to afford the key intermediates **7–10**. As mentioned above, significant down-field shift of the H-4' signals was observed with the ¹H NMR spectra of **7–10** (δ 4.3–4.5), as compared to that of **25–28** (δ 3.5–3.6), which further substantiated the attachment

of the phosphate group to the 4'-O-position. All of the intermediates involved in the syntheses were fully characterized by ¹H, ¹³C and correlation NMR spectroscopy and by MALDI-TOF MS.

To be readied for conjugation with sTnNPhAc, MPLA derivatives **7–10** were treated with Zn and acetic acid to selectively reduce the azido group (Scheme 3). The resulting amines **30–33** were then acylated with succinic acid anhydride to produce **34–37**, which had a free carboxylic acid group attached to the disaccharide reducing end. For coupling with sTnNPhAc derivative **42** that carried a free amino group, acids **34–37** were converted to active esters **38–41** by reacting with *p*-nitrophenol and EDCI. These reactions were fast, and the active esters formed were stable enough to enable silica gel column purification. Subsequently, **38–41** were coupled with **42** in DMF. These reactions were smooth and clean to afford **43–46** that were readily purified by silica gel column chromatography and by preparative TLC. Finally, catalytic hydrogenolysis of the benzyl ether linkages in **43–46** under a H₂ atmosphere in the presence of 10% Pd/C afforded the desired glycoconjugates **3–6**, which were confirmed by both NMR and MS. Thereafter, **3–6** were subjected to biological evaluation in mice.

Immunological investigation of the MPLA-sTnNPhAc conjugates **3–6** was carried out with female C57BL/6 mouse, a well-characterized inbred strain for vaccine evaluation and comparable to the human in immune reactions.³³ To improve the solubility of these conjugates in buffer, they were incorporated in liposomes formed by using 1,2-distearoyl-*sn*-glycero-3-phosphocholine and cholesterol. Delivering glycoconjugate vaccines in liposomal forms can further improve their immunogenicity.^{36,37} For immunization, the liposomal preparations of **3–6** were each injected subcutaneously (s.c.) to a group of six mice. In the meantime, groups of mice were simultaneously inoculated with emulsions of **3–6** liposomes and Titermax Gold to evaluate the potential influence of an external adjuvant on the activities of these glycoconjugate vaccines. The adopted vaccination schedule was to inject into each



Scheme 3 Assembly of the target conjugates 3–6.

mouse 0.1 mL of a vaccine preparation containing *ca.* 3 μg of sTnNPhAc on days 1, 14, 21 and 28, respectively. Each mouse was subjected to bleeding on day 0 before the first injection (used as blank controls) and on days 27 and 38 after immunization. The blood samples were treated according to standard protocols to prepare antisera for the analysis of sTnNPhAc-specific antibodies by enzyme-linked immunosorbent assay (ELISA) with the human serum albumin (HSA) conjugate of sTnNPhAc (sTnNPhAc-HSA)³⁸ as a capture antigen. In addition to total antibodies, antibody isotypes such as IgG1, IgG2a, IgG3 and IgM were also individually assessed. Antibody titers were calculated from linear regression analysis of the curves of the optical density (OD) value against serum dilution number, and were defined as the dilution number yielding an OD value of 0.2.³⁴

The ELISA results of antisera obtained with 3 without the use of an external adjuvant (Fig. 2A) revealed that the conjugate itself provoked a strong sTnNPhAc-specific immune response in mice. More importantly, in addition to IgM antibody, a high titer of IgG3 and some IgG1 antibodies were also observed, suggesting a T cell-dependent immune response.^{39,40} Interestingly, 3 plus Titermax Gold failed to stimulate a significant antibody response (Fig. 2B), although IgM antibodies were observed. These results suggest that the external adjuvant had an inhibitory effect on the T cell-mediated immune response against 3 but a relatively small influence on the B cell response. The results agreed well with our previous report about other MPLA-TACA conjugates.³⁴

The ELISA results of 4 (Fig. 2C and 2D) were similar to that of 3 in terms of the type of immune responses provoked and the influence of Titermax Gold. However, the IgM and IgG




Fig. 2 ELISA results of day 38 antisera of mice immunized with 3 alone (A), 3 plus Titermax Gold (B), 4 alone (C), 4 plus Titermax Gold (D), 5 alone (E), and 6 alone (F), respectively. The titers of various sTnNPhAc-specific antibodies are displayed. Each dot represents the antibody titer of an individual mouse, and the black bar shows the average antibody titer of a group of six mice.




Fig. 3 Competitive ELISA results of the pooled mouse antisera obtained with conjugates 4, 5, 6, and 4 plus adjuvant. The OD values at 405 nm reflected the levels of total antibodies bound to the sTnNPhAc-HSA-coated ELISA plates in the presence of specified concentrations of 42.

antibody titers for 4 were much higher than that for 3. Moreover, 4 stimulated a strong immune response in all of the mice. Another interesting finding was that 4 also induced a significant level of IgG1 antibody which is usually observed with neoglycoprotein vaccines. The ELISA results of 5 (Fig. 2E) and 6 (Fig. 2F) were similar to that of 3 and 4, but their antibody titers were lower than that of 4 and higher than that of 3. In all of these cases, Titermax Gold showed inhibition of the immune responses against the conjugate vaccines.

To further verify that the detected antibodies were indeed sTnNPhAc-specific, a competitive ELISA experiment for the pooled mouse antisera obtained with 4, 5, 6, and 4 plus adjuvant was performed. In this study, various concentrations (0, 0.00152, 0.0152, 0.152, 1.52, 15.2 and 152 μM) of free sTnNPhAc (42) were added to compete with sTnNPhAc-HSA attached to the ELISA plate for antibody binding. Again, alkaline phosphatase-linked goat anti-mouse kappa antibody was employed as the secondary antibody. The OD values at 405 nm reflected the levels of total antibodies bound to the sTnNPhAc-HSA-coated plates. These studies (Fig. 3) have revealed that 42 had concentration-dependent inhibition of antibody binding to the plates and that, at concentrations higher than 152 μM, 42 could essentially completely inhibit the binding, proving that the elicited antibodies were indeed specific to the sTnNPhAc antigen.

The above ELISA results disclosed that the MPLA conjugates 3–6 alone, that is, in the absence of any external adjuvant, provoked a robust antigen-specific immune response in mice. More importantly, they stimulated high titers of IgG3 and IgG1 antibodies, indicating a T cell-mediated immune response desirable for cancer immunotherapy.^{39,40} Therefore, all of the synthetic MPLA derivatives could act both as a vaccine carrier and as an adjuvant to effectively enhance the immunogenicity of sTnNPhAc and to efficiently promote a T cell-dependent immune response toward TACAs.^{18,34,41} These properties of MPLA should be particularly useful for the design and development of new fully synthetic carbohydrate-based conjugate vaccines.

Titermax Gold is a commercial adjuvant that is commonly used with vaccines such as carbohydrate-protein conjugates⁴²




Fig. 4 Comparison of the average antibody titers of sTnNPhAc-specific kappa (total), IgG3, and IgM antibodies in day 38 antisera of mice immunized with 3, 4, 5, and 6.

to improve their immunogenicity. However, we showed in this work that it had an inhibitory impact on the immunological activities of 3–6. As Titermax Gold mainly inhibited the elicitation of IgG antibodies, it seemed to mainly affect the T cell-mediated immune response. We proposed that Titermax Gold might interact with MPLA in the conjugate vaccine to prevent MPLA from binding to cell surface TLRs and affect its engagement in T cell-mediated immunological pathways. Another hypothesis is that Titermax Gold might interact with MPLA to affect vaccine delivery to the lymph system or antigen presenting cells required for effective T cell-mediated immunity.^{43,44} This discovery and further detailed studies may help understand vaccine adjuvants, such as their binding sites, functional mechanisms and so on.

Despite the fact that 3–6 stimulated similar patterns of immune responses in mice, they did show differences in immunological activity. For example, conjugate 4 elicited much higher and more consistent titers of both total antibodies and IgG3 antibody than 3, 5 and 6 (Fig. 4). Moreover, all of the mice showed strong immune responses to 4 but not to other conjugates. Clearly, 4 was the best vaccine among the MPLA conjugates investigated.

Structurally, 3–6 were different only in their MPLA moiety, and their antisera were obtained by means of the same immunization protocol and schedule. Consequently, any difference in their immunological activity should reflect the impact of the MPLA structure. Conjugate 4 contained the monophosphoryl form of natural lipid A of *N. meningitidis* (H44/76 strain).¹⁵ In 3, the two free hydroxyl groups on the lipid chains of the MPLA moiety were removed; 5 and 6 were different in that their MPLA contained different lengths and different numbers of lipid chains than that of 4. As all these conjugates had essentially the same immunological profile, it seems that

the free hydroxyl groups on the lipid chains and the length and number of lipid chains of MPLA had a quantitative, rather than qualitative, impact on its biological activities. From what was said, it is clear that the hydroxyl groups on the lipid chains play an important role in the MPLA interaction with its receptors, as **3** was significantly less potent than both **4** and **5**. Elongating the length of lipids at the 3'-O- and 3'-O-positions and additional lipid chains linked to these lipids seem to have a relatively small impact, although such changes did result in reduced immunological activity and some inconsistency of immune responses in individual mice.

Conclusion

In summary, a convergent and efficient strategy was developed for the synthesis of MPLA derivatives suitable for conjugation with carbohydrate antigens, and was employed to synthesize four monophosphoryl derivatives of *N. meningitidis* lipid A. These MPLA derivatives were coupled with sTnNPhAc to form fully synthetic glycoconjugate cancer vaccines. The strategy should be generally applicable in preparing other MPLA derivatives and MPLA-carbohydrate conjugates. Studies on the resulting MPLA-sTnNPhAc conjugates revealed that they elicited strong and T cell-dependent immune responses without the use of any external adjuvant. Our previous work revealed that antisera derived from mice immunized with MPLA conjugates could effectively bind to and kill cancer cells metabolically engineered to express the corresponding antigen.³⁴ MPLA has thus been demonstrated to be a useful platform for the development of new vaccine carriers and adjuvants and for the development of novel types of fully synthetic carbohydrate-based cancer vaccines with self-adjuvanting properties. Our results have also revealed that MPLA derivatives containing six lipid chains exhibited more potent immunostimulatory activities than those with eight lipid chains (conjugate **6**) and that the lipid structure and length had a significant impact on the immunology of MPLA. The monophosphoryl form of natural *N. meningitidis* lipid A was found to have the most promising immunological properties and its sTnNPhAc conjugate elicited the most potent and the most consistent T cell-dependent anti-sTnNPhAc immunity. As a result, the MPLA moiety in **4** was identified as the first generation of optimized vaccine carriers and adjuvants that is under further optimization and additional investigation.

On the other hand, Titermax Gold was found to inhibit the immunological activity of MPLA-sTnNPhAc conjugates, whereas it has the opposite influence on the activity of protein-sTnNPhAc conjugates.^{35,38} It is proposed that Titermax Gold may interact with MPLA to affect its binding to cell surface receptors and/or its delivery to the lymph system or antigen presenting cells. It is anticipated that these issues may be clarified by studies utilizing labeled MPLA derivatives and conjugates, the results of which should be useful for understanding the immunostimulatory and adjuvant activities and the functional mechanisms of MPLA and Titermax

Gold, which is a very important topic in cancer vaccine immunology.

Experimental section

Synthesis of the target glycoconjugates 3–6

Detailed experimental procedures for these syntheses and analytical data of compounds are described in the ESI† document.

Preparation of liposomes of glycoconjugates 3–6

The protocol was similar to that reported in the literature.³⁴ In brief, a mixture of a specific conjugate **3–6** (0.474 μmol), 1,2-distearoyl-*sn*-glycero-3-phosphocholine (2.44 mg, 3.08 μmol), and cholesterol (0.92 mg, 2.37 μmol) (in a 10:65:50 molar ratio) was dissolved in CH_2Cl_2 and MeOH (1:1, v/v, 2 mL) in a round-bottom flask. The solvents were then removed under vacuum to form a thin lipid film on the flask wall, which was hydrated by adding 2.0 mL of HEPES buffer (20 mM, pH 7.5) containing NaCl (150 mM) and shaking the mixture under an argon atmosphere at 40 °C for 1 h. The suspension was finally sonicated for 1 min to obtain the desired liposomes.

Immunization of mice

Each group of six female C57BL/6 mice was inoculated on day 1 by subcutaneous (s.c.) injection of 0.1 mL of the liposomal solution of a specific glycoconjugate containing 10 μg of the carbohydrate antigen or by injection of an emulsion of the liposomal solution (0.1 mL) of a specific glycoconjugate with Titermax Gold adjuvant (0.05 mL) prepared according to the manufacturer's protocol. Following the initial immunization, the mice were boosted 3 times on day 14, day 21 and day 28 by s.c. injection of the same conjugate and by means of the same immunization protocol. Mouse blood samples were collected prior to the initial immunization on day 0 and after immunization on day 27 and day 38, and were clotted to obtain antisera that were stored at $-80\text{ }^\circ\text{C}$ before use. The animal protocol (#A 03-10-11) for this investigation was approved by the Institutional Animal Care and Use Committee (IACUC) of Wayne State University, and all animal experiments were performed in compliance with the relevant laws and institutional guidelines.

ELISA

ELISA plates were treated with 100 μL of a solution of the sTnNPhAc-HSA conjugate (2 $\mu\text{g mL}^{-1}$) dissolved in coating buffer (0.1 M bicarbonate, pH 9.6) at 37 °C for 1 h, which was followed by treatment with a blocking buffer and washing 3 times with phosphate-buffered saline (PBS) containing 0.05% Tween-20 (PBST). Then, the pooled or an individual mouse antiserum with serial half-log dilutions from 1:300 to 1:656 100 in PBS was added to the coated plates (100 μL per well) and incubated at 37 °C for 2 h. The plates were washed with PBS and incubated at rt for another hour with a 1:1000 diluted solution of alkaline phosphatase linked goat anti-

mouse kappa, IgM or IgG2a antibody or a 1 : 2000 diluted solution of alkaline phosphatase linked goat anti-mouse IgG1 or IgG3 antibody (100 μ L per well), respectively. Finally, the plates were washed with PBS and developed with 100 μ L of a *p*-nitrophenylphosphate (PNPP) solution (1.67 mg mL⁻¹ in buffer) for 30 min at rt for colorimetric readout using a microplate reader at 405 nm wavelength. For titer analysis, OD values were plotted against antiserum dilution values, and a best-fit line was obtained. The equation of this line was used to calculate the dilution value at which an OD of 0.2 was achieved, and the antibody titer was calculated at the inverse of the dilution value. For the competitive ELISA experiments, before the addition of antiserum (80 μ L) to each well of the plate, a solution of free sTnNPhAc derivative **42** (20 μ L) was added to reach the final concentration of 0, 0.00152, 0.0152, 0.152, 1.52, 15.2 and 152 μ M, respectively. The remaining operations were the same as those described above.

Acknowledgements

This research work was supported by an NIH/NCI grant (R01 CA95142).

Notes and references

- Q. Wang and Z. Guo, *Curr. Opin. Chem. Biol.*, 2009, **13**, 608–617.
- F. Peri, *Chem. Soc. Rev.*, 2013, **42**, 4543–4556.
- A. Kaiser, N. Gaidzik, T. Becker, C. Menge, K. Groh, H. Cai, Y. Li, B. Gerlitzki, E. Schmitt and H. Kunz, *Angew. Chem., Int. Ed.*, 2010, **49**, 3688–3692.
- Z. Huang, L. Shi, J. Ma, Z. Sun, H. Cai, Y. Chen, Y. Zhao and Y. Li, *J. Am. Chem. Soc.*, 2012, **134**, 8730–8733.
- V. Lakshminarayanan, P. Thompson, M. A. Wolfert, T. Buskas, J. M. Bradley, L. B. Pathange, C. S. Madsen, P. A. Cohenc, S. J. Gendler and B.-J. Boons, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 261–266.
- A. M. Abdel-Aal, D. El-Naggar, M. Zaman, M. Batzloff and I. Toth, *J. Med. Chem.*, 2012, **55**, 6968–6974.
- H. Cai, Z. Sun, Z. Huang, L. Shi, Y. Zhao, H. Kunz and Y. Li, *Chem.-Eur. J.*, 2013, **19**, 1962–1970.
- C. Erridge, E. Bennett-Guerrero and I. R. Poxton, *Microbes Infect.*, 2002, **4**, 837–851.
- C. A. Alving and M. Rao, *Vaccine*, 2008, **26**, 3036–3045.
- M. A. Dobrovolskaia and S. N. Vogel, *Microbes Infect.*, 2002, **4**, 903–914.
- E. S. Van Amersfoort, T. J. C. Van Berkel and J. Kuiper, *Clin. Microbiol. Rev.*, 2003, **16**, 379–414.
- E. T. Rietschel, T. Kirikae, F. U. Schade, U. Mamat, G. Schmidt, H. Loppnow, A. J. Ulmer, U. Zahring, U. Seydel, F. Di Padova, M. Schreier and H. Brade, *FASEB J.*, 1994, **8**, 217–225.
- A. G. Stover, J. D. S. Correia, J. T. Evans, C. W. Cluff, M. W. Elliott, E. W. Jeffery, D. A. Johnson, M. J. Lacy, J. R. Baldridge, P. Probst, R. J. Ulevitch, D. H. Persing and R. M. Hershsberg, *J. Biol. Chem.*, 2004, **279**, 4440–4449.
- S. Kanegasaki, K. Tanamoto, T. Yasuda, J. Y. Homma, M. Matsuura, M. Nakatsujka, Y. Kumazawa, A. Yamamoto, T. Shiba, S. Kusumoto, M. Imoto, H. Yoshimura and T. Shimamoto, *J. Biochem.*, 1986, **99**, 1203–1210.
- V. A. Kulshin, U. Zahring, B. Lindner, C. E. Frasch, C. M. Tsai, B. A. Dmitriev and E. T. Rietschel, *J. Bacteriol.*, 1992, **174**, 1793–1800.
- E. Ribi, J. Cantrell, T. Feldner, K. Myers and J. Peterson, *Microbiol.*, 1986, 9–13.
- J. Baldridge, K. Myers, D. Johnson, D. Persing, C. Cluff and R. Hershsberg, in *Vaccine Adjuvants*, ed. C. J. Hackett and D. A. J. Harn, Humana Press Inc., Totowa, NJ, 2006, pp. 235–255.
- C. R. Casella and T. C. Mitchell, *Cell. Mol. Life Sci.*, 2008, **65**, 3231–3240.
- K. Takayama, N. Qureshi, E. Ribi and J. Cantrell, *Rev. Infect. Dis.*, 1984, **6**, 439–443.
- Y. Fukase, Y. Fujimoto, Y. Adachi, Y. Suda, S. Kusumoto and K. Fukase, *Bull. Chem. Soc. Jpn.*, 2008, **81**, 796–819.
- Z. Jiang, W. A. Budzynski, D. Qiu, D. Yalamati and R. R. Koganty, *Carbohydr. Res.*, 2007, **342**, 784–796.
- A. Zamyatina, H. Sekljic, H. Bradeb and P. Kosmaa, *Tetrahedron*, 2004, **60**, 12113–12137.
- B. Santhanam, M. A. Wolfert, J. N. Moore and G.-J. Boons, *Chem. – Eur. J.*, 2004, **10**, 4798–4807.
- Y. Watanabe, K. Miura, M. Shiozaki, S. Kanai, S. Kurakata and M. Nishijima, *Carbohydr. Res.*, 2003, **338**, 47–54.
- A. V. Demchenko, M. A. Wolfert, B. Santhanam, J. N. Moore and G.-J. Boons, *J. Am. Chem. Soc.*, 2003, **125**, 6103–6112.
- T. Mochizuki, Y. Iwano, M. Shiozaki, S. Kurakata, S. Kanae and M. Nishijima, *Tetrahedron*, 2000, **56**, 7691–7703.
- D. A. Johnson, D. S. Keegan, C. G. Sowell, M. T. Livesay, C. J. Johnson, L. M. Taubner, A. Harris, K. R. Myers, J. D. Thompson, G. L. Gustafson, M. J. Rhodes, J. T. Ulrich, J. R. Ward, Y. M. Yorgensen, J. L. Cantrell and V. G. Brookshire, *J. Med. Chem.*, 1999, **42**, 4640–4649.
- K. Fukase, Y. Fukase, M. Oikawa, W. Liu, Y. Suda and S. Kusumoto, *Tetrahedron*, 1998, 4033–4050, (and references therein).
- W. J. Christ, P. D. McGuinness, O. Asano, Y. Wang, M. A. Mullarkey, M. Perez, L. D. Hawkins, T. A. Blythe, G. R. Dubuc and A. L. Robidoux, *J. Am. Chem. Soc.*, 1994, **116**, 3637–3638.
- A. J. Ulmer, H. Heine, W. Feist, S. Kusumoto, T. Kusama, H. Brade, F. U. Schade, E. T. Rietschel and H. D. Flad, *Infect. Immun.*, 1992, **60**, 3309–3314.
- C. Galanos, O. Luderitz, E. T. Rietschel, O. Westphal, H. Brade, L. Brade, M. Freudenberg, U. Schade, M. Imoto and H. Yoshimura, *FEBS*, 1985, **148**, 1–5.
- Q. Wang and Z. Guo, *Chem. Commun.*, 2009, 5536–5537.
- Y. Pan, P. Chefalo, N. Nagy, C. V. Harding and Z. Guo, *J. Med. Chem.*, 2005, **48**, 875–883.
- Q. Wang, Z. Zhou, S. Tang and Z. Guo, *ACS Chem. Biol.*, 2012, **7**, 235–240.

- 35 J. Wu and Z. Guo, *Bioconjugate Chem.*, 2006, **17**, 1537–1544.
- 36 T. Buskas, S. Ingale and G. J. Boons, *Angew. Chem., Int. Ed.*, 2005, **44**, 5985–5988.
- 37 F. Estevez, A. Carr, L. Solorzano, O. Valiente, C. Mesa, O. Barroso, G. V. Sierra and L. E. Fernandez, *Vaccine*, 1999, **18**, 190–197.
- 38 Q. Wang, S. A. Ekanayaka, J. Wu, J. Zhang and Z. Guo, *Bioconjugate Chem.*, 2008, **19**, 2060–2068.
- 39 R. B. Markham, G. B. Pier and J. R. Schreiber, *J. Immunol.*, 1991, **146**, 316–320.
- 40 A. L. Gavin, N. Barnes, H. M. Dijkstra and P. M. Hogarth, *J. Immunol.*, 1998, **160**, 20–23.
- 41 D. H. Persing, R. N. Coler, M. J. Lacy, D. A. Johnson, J. R. Baldridge, R. M. Hershberg and S. G. Reed, *Trends Microbiol.*, 2002, **10**(Suppl.), S32–S37.
- 42 Q. Wang and Z. Guo, *ACS Med. Chem. Lett.*, 2011, **2**, 373–378.
- 43 R. M. Zinkernagel, S. Ehl, P. Aichele, S. Oehen, T. Kundig and H. Hengartner, *Immunol. Rev.*, 1997, **156**, 199–209.
- 44 D. T. O'Hagan, M. L. MacKichan and M. Singh, *Biomol. Eng.*, 2011, **18**, 69–85.