

Figure 5. Field desorption spectrum of potassium dihydrogen phosphate, 306 °C.

mental conditions result in the formation of Al^+ and Fe^+ ions from the cathode surface, in addition to Na^+ and Mg^+ ions present as impurities. This spectrum, obtained at a higher field strength, is therefore not a pure field desorption spectrum but a composite of field desorption and ion bombardment. We may be observing here behavior similar to that observed under secondary ion mass spectrometry (SIMS).^{12,13}

An FD spectrum of potassium phosphate, a material containing a polyvalent anion, is shown in Figure 5. Here we can identify, in addition to K^+ ions, KH_3PO_4^+ , $\text{K}_2\text{H}_2\text{PO}_4^+$, K_3HPO_4^+ , and K_4PO_4^+ . Analogously, the analysis of K_2CO_3 showed only K_3CO_3^+ ions in addition to K^+ . In contrast, under electron impact one might observe a minute yield of K_2CO_3^+ ¹⁴ but no K_3CO_3^+ .

Again it should be emphasized that these are merely rudimentary illustrations of the type of chemistry that can be studied by field desorption mass spectrometry. Unlike classical electron impact mass spectrometry, which is highly useful in the study of the concentration and the structure of inorganic species in the gas phase, and in contrast to SIMS, which is highly useful in determining the elemental composition of thin solid layers, field desorption can teach us about the structure of inorganic solids. This technique could demonstrate nearest neighbor relations in solutions of inorganic materials and provide information on the relative stability of cluster ions. It could also indicate charge-transfer processes in solution, in addition to allowing the assessment of the isotopic composition of oxy anions, PO_4^{3-} in particular, using ^{18}O as a tracer. The great number of possible applications of this technique and its potential as a novel research tool are the main reasons for bringing it this early to the attention of the inorganic chemistry research community.

Registry No. Potassium iodide, 7681-11-0; potassium fluoride, 7789-23-3; ammonium chloride, 12125-02-9; potassium chloride, 7447-40-7; lithium iodide, 10377-51-2; potassium bromide, 7758-02-3; sodium iodide, 7681-82-5; ammonium nitrate, 6484-52-2; calcium fluoride, 7789-75-5; calcium chloride, 10043-52-4; magnesium chloride, 7786-30-3; uranium fluoride, 10049-14-6; potassium dihydrogen phosphate, 7778-77-0.

References and Notes

- H. D. Beckey and H. R. Schulten, *Angew. Chem., Int. Ed. Engl.*, **14**, 403 (1975).
- M. Anbar and G. A. St. John, *Anal. Chem.*, **48**, 198 (1976).
- C. A. Spindt, *J. Appl. Phys.*, **39**, 3504 (1968).
- H. D. Beckey, "Field Ionization Mass Spectrometry", Pergamon Press, Elmsford, N.Y., 1971.
- A. J. Robertson, *J. Phys. E.*, **7**, 321 (1974).
- M. Anbar and W. H. Aberth, *Anal. Chem.*, **46**, 59A (1974).
- M. Anbar and G. A. St. John, *J. Am. Chem. Soc.*, **97**, 7195 (1975).
- R. W. Kiser, J. G. Dillard, and D. L. Dugger, *Adv. Chem. Ser.*, No. **72**, 153 (1968).
- P. Kearnle, *Adv. Chem. Ser.*, No. **72**, 24 (1968).
- I. Lauder, C. Towler, and T. Wuth, *Adv. Mass Spectrom.*, **5**, 379 (1971).
- R. C. Schoonmaker and R. F. Porter, *J. Chem. Phys.*, **28**, 168 (1958).
- R. F. K. Herzog and F. P. Viehblock, *Phys. Rev.*, **76**, 855 (1949).
- C. A. Andersen, *Int. J. Mass Spectrom. Ion Phys.*, **2**, 61 (1969).
- L. N. Gorokhov and G. A. Semenov, *Adv. Mass Spectrom.*, **5**, 349 (1971).

Contribution No. 2733 from the Department of Chemistry, Indiana University, Bloomington, Indiana 47401

Zerovalent Triarylphosphine Complexes of Nickel, Palladium, and Platinum

Philip E. Garrou*¹ and George E. Hartwell*¹

Received July 16, 1975

AICS0501N

Although $\text{Pd}(\text{PPh}_3)_4$,² $\text{Pt}(\text{PPh}_3)_4$,³ and $\text{Ni}(\text{PPh}_3)_4$ have been known for some time, the nature of the species present in solution remained obscure until recently.⁵ Tolman examined solutions of the ML_4 complexes by ^{31}P and ^1H NMR and uv-visible spectroscopy. All observations are consistent with complete dissociation of the ML_4 complexes to ML_3 and L [$\text{L} = \text{PPh}_3$, $\text{P}(p\text{-CH}_3\text{C}_6\text{H}_4)_3$]. Recent studies have reported a relationship between the steric requirements of the phosphine ligands and the degree of dissociation to $\text{M}(\text{PR}_3)_3$ ($\text{M} = \text{Ni}$,^{6,7} Pd ⁸). The $\text{Ni}(0)$ complexes were categorized according to the degree of ligand dissociation and the rate of exchange, on the NMR time scale, between free and complexed ligands.

Unlike the triarylphosphine complexes, the zerovalent triarylphosphine complexes of Ni, Pd, and Pt have not been extensively studied. $\text{Ni}(\text{SbPh}_3)_4$ was reported by Wilke.⁹ $\text{Pd}(\text{SbPh}_3)_4$, the only reported palladium complex, was prepared by the reaction of (acetylacetonato)[8-(acetylacetonyl)cyclooct-4-enyl]palladium with Ph_3Sb .¹⁰ There are no literature references to $\text{Pt}(\text{SbR}_3)_4$. The mixed-ligand complexes $\text{Pd}(\text{PPh}_3)_2(\text{SbPh}_3)_2$ and $\text{Pt}(\text{PPh}_3)_3(\text{SbPh}_3)$ were formed from the reaction of excess SbPh_3 with $\text{Pd}(\text{PPh}_3)_4$ and $\text{Pt}(\text{PPh}_3)_4$.¹¹ All of the reported triarylphosphine complexes are insoluble and thus do not lend themselves to solution studies. We therefore undertook the study of soluble zerovalent triarylphosphine complexes in order to compare preparations, stabilities, and solution properties with those of the well-known phosphine analogues. This work is a continuation of our studies involving triarylphosphine-transition metal complexes.^{12,13}

Experimental Section

All reactions were carried out in Schlenk apparatus. Solvents were dried prior to use. Manipulations of the compounds were carried out in an inert-atmosphere box or in Schlenk apparatus under argon. The complexes $\text{Ni}(\text{COD})_2$,¹⁴ ($\text{COD} = 1,5\text{-cyclooctadiene}$), $\text{Pd}(\text{PPh}_3)_4$,¹⁵ $\text{Pt}(\text{PPh}_3)_3$,¹⁶ and $\text{Pt}(\text{PPh}_3)_2(\text{C}_2\text{H}_4)$ ¹⁷ and the ligands $\text{Sb}(p\text{-CH}_3\text{C}_6\text{H}_4)_3$,¹⁸ $\text{Sb}(p\text{-CH}_3\text{OC}_6\text{H}_4)_3$,¹⁸ and $\text{Sb}(\text{SPh})_3$ ¹⁹ were prepared by literature methods. Analytical data are given in Table I. Preparations are shown in Figure 1.

(a) **Preparation of $\text{Ni}(\text{SbR}_3)_4$ ($\text{R} = \text{Ph}$, $p\text{-CH}_3\text{C}_6\text{H}_4$, $p\text{-CH}_3\text{OC}_6\text{H}_4$), Method I.** $(\text{CH}_3\text{CH}_2)_3\text{Al}$ (0.50 ml, 0.46 mmol) was added slowly in a Schlenk filtration apparatus to a dried and deaerated diethyl ether solution (30 ml) of $\text{Ni}(\text{acac})_2$ (0.5 g, 0.19 mmol) and SbR_3 (0.80 mmol) at 0°C. A yellow crystalline material precipitated ($\text{R} = \text{Ph}$). For the more soluble complexes ($\text{R} = p\text{-CH}_3\text{C}_6\text{H}_4$, $p\text{-CH}_3\text{OC}_6\text{H}_4$), the solution was reduced in volume to ca. 15 ml and 10 ml of hexane was added, inducing the precipitation of an off-white solid. The complexes were collected by filtration, washed with hexane, and dried under vacuum; yield ca. 75%.


(b) **Preparation of $\text{Ni}(\text{SbR}_3)_4$ ($\text{R} = \text{Ph}$, $p\text{-CH}_3\text{C}_6\text{H}_4$, $p\text{-CH}_3\text{OC}_6\text{H}_4$), Method II.** A solution of SbR_3 (0.15 mmol) was added slowly to a benzene solution (20 ml) of $\text{Ni}(\text{COD})_2$ (0.10 g, 0.036 mmol) in a Schlenk filtration apparatus. Further work-up was identical with that described in section a.

(c) **Preparation of $\text{Pd}(\text{SbR}_3)_4$ ($\text{R} = \text{Ph}$, $p\text{-CH}_3\text{C}_6\text{H}_4$, $p\text{-CH}_3\text{OC}_6\text{H}_4$).** $(\text{CH}_3\text{CH}_2)_3\text{Al}$ (0.21 ml, 0.20 mmol) was added slowly in a Schlenk filtration apparatus to a dried and deaerated diethyl ether solution (30 ml) of $\text{Pd}(\text{acac})_2$ ($\text{acac} = \text{acetylacetonate}$) (0.25 g, 0.08 mmol) and SbR_3 (0.34 mmol) at 0°C. The yellow solution was stirred at 0°C for 1 hr. Further work-up was identical with that described in section a.

(d) **Preparation of $\text{Pd}(\text{PPh}_3)_2(\text{SbR}_3)_2$ ($\text{R} = \text{Ph}$, $p\text{-CH}_3\text{C}_6\text{H}_4$), Method I.** A benzene solution (10 ml) of SbR_3 (0.086 mmol) was added dropwise to a suspension of $\text{Pd}(\text{PPh}_3)_4$ (0.25 g, 0.022 mmol)

Table I. Analytical Data for Zerovalent Ni, Pd, and Pt Complexes of Triarylstibines^a

Compd	C	H	M	S
Ni[Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₄	61.56 (60.75)	5.13 (4.93)	3.69 (3.27)	
Pd[SbPh ₃] ₄	56.94 (56.75)	4.97 (4.83)	7.01 (6.71)	
Pd[Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₄	59.91 (59.61)	5.03 (4.95)	6.12 (5.80)	
Pd[Sb(<i>p</i> -CH ₃ OC ₆ H ₄) ₃] ₄	57.71 (58.02)	4.85 (4.90)		
Pd(PPh ₃) ₂ [Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₂	64.52 (64.06)	4.54 (4.55)	8.01 (7.67)	
Pd(PPh ₃) ₂ (SbPh ₃) ₂	66.08 (65.78)	5.08 (4.76)	7.27 (7.01)	
Pt(PPh ₃) ₂ (SbPh ₃) ₂	60.88 (59.94)	4.23 (4.01)		
Pt(PPh ₃) ₂ [Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₂	62.05 (61.61)	4.75 (4.95)	12.87 (12.83)	
Pt(PPh ₃) ₂ [Sb(SPh) ₃] ₂	53.35 (52.84)	3.70 (3.76)		11.85 (11.42)

^a Calculated (found).**Figure 1.** Preparation of zerovalent nickel, palladium, and platinum complexes of triarylstibines.

in benzene (20 ml) at room temperature. The resulting white powder (R = Ph) was collected by filtration, washed with hexane, and dried under vacuum. For the more soluble complex (R = *p*-CH₃C₆H₄) the solvent was removed under vacuum to give a yellow oil. Washing the oil with acetone-hexane gave a yellow powder which was collected by filtration, washed with hexane, and vacuum dried; yield ca. 70%.

(e) **Preparation of Pd(PPh₃)₂[Sb(*p*-CH₃C₆H₄)₃]₂, Method II.** A suspension of Pd(CS₂)(PPh₃)₂ (0.10 g, 0.015 mmol) and Sb(*p*-CH₃C₆H₄)₃ (0.12 g, 0.30 mmol) in benzene (25 ml) was stirred overnight under argon. Unreacted Pd(CS₂)(PPh₃)₂ was removed by filtration and the solvent was removed under vacuum to give a yellow oil. The complex was further worked up as described in section d; yield ca. 50–60%.

(f) **Preparation of Pt(PPh₃)₂(SbR₃)₂ (R = Ph, *p*-CH₃C₆H₄), Method I.** A benzene solution (10 ml) of SbR₃ (0.114 mmol) was added dropwise to a suspension of Pt(PPh₃)₃ (0.25 g, 0.025 mmol) in benzene (20 ml) at room temperature. The resulting yellow powder (R = Ph) was collected by filtration, washed with hexane, and dried under vacuum.

For R = *p*-CH₃C₆H₄, after 3.5 h the solvent was removed under vacuum to give a yellow oil. Further work-up was as described in section d. The yield of Pt(PPh₃)₂[Sb(*p*-CH₃C₆H₄)₃]₂ was 60–70% and that of Pt(PPh₃)₂(SbPh₃)₂ was 90–95%.

(g) **Preparation of Pt(PPh₃)₂(SbR₃)₂ (R = Ph, *p*-CH₃C₆H₄), Method II.** A benzene solution (10 ml) of SbR₃ (0.04 mmol) was added dropwise to a benzene solution (10 ml) of Pt(PPh₃)₂(C₂H₄) (0.15 g, 0.02 mmol). After 1 h of stirring at room temperature, the solvent was removed under vacuum to give an oil. Further work-up was as in section d; yield ca. 80–85%.

(h) **Preparation of Pt(PPh₃)₂[Sb(*p*-CH₃C₆H₄)₃]₂, Method III.** A suspension of Pt(CS₂)(PPh₃)₂ (0.3 g, 0.025 mmol) and Sb(*p*-CH₃C₆H₄)₃ (0.2 g, 0.05 mmol) in benzene (25 ml) was stirred overnight under argon. Further work-up was as in section d; yield 60–65%.

(i) **Preparation of Pt(PPh₃)₂[Sb(SPh)₃]₂.** A benzene solution (10 ml) of Sb(SPh)₃ (0.18 g, 0.04 mmol) was added dropwise to a benzene solution (10 ml) of Pt(PPh₃)₂(C₂H₄) (0.15 g, 0.02 mmol). After 1.5 h of stirring at room temperature, the lime green solution was reduced in volume to ca. 5 ml and 10 ml of hexane was added to induce precipitation. The yellow complex was collected by filtration, washed with Et₂O, and dried under vacuum; yield ca. 75%.

Discussion

Ni(PPh₃)₄ is easily prepared by the zinc dust reduction of NiCl₂ in the presence of PPh₃.⁵ When the reaction was attempted with SbR₃, a black metallic precipitate was observed and no triarylstibinenickel complex was obtained. Ni(SbR₃)₄,

Table II. 220-MHz ¹H NMR^a Data for Zerovalent Ni, Pd and Pt Complexes of Triarylstibines

Compd	Aromatic	CH ₃ or OCH ₃
Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃	1.80, 2.00	7.69
Sb(<i>p</i> -CH ₃ OC ₆ H ₄) ₃	2.75, 3.20	6.27
Pd[Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₄	3.00, 3.32	7.74
Pd[Sb(<i>p</i> -CH ₃ OC ₆ H ₄) ₃] ₄	3.05, 3.64	6.39
Ni[Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₄	3.10, 3.36	7.77
Ni[Sb(<i>p</i> -CH ₃ OC ₆ H ₄) ₃] ₄	3.12, 3.72	6.42
Pd(PPh ₃) ₂ [Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₂	2.27–3.40 ^b	7.75
Pt(PPh ₃) ₂ [Sb(<i>p</i> -CH ₃ C ₆ H ₄) ₃] ₂	2.27–3.41 ^b	7.73, 7.77

^a τ scale, CDCl₃. ^b Multiplets due to several species present in solution.

was, however, prepared by the (CH₃CH₂)₃Al reduction of Ni(acac)₂ in the presence of SbR₃ (R = Ph, *p*-CH₃C₆H₄, *p*-CH₃OC₆H₄) or by the replacement of cyclooctadiene from Ni(COD)₂.

Pd(PPh₃)₄ is easily prepared by the hydrazine hydrate reduction of PdCl₂ in the presence of PPh₃ or by NaBH₄ reduction of K₂PdCl₄ in the presence of PPh₃.²⁰ When either of these standard methods was attempted with SbR₃, a black metallic precipitate was obtained. The palladium complexes Pd(SbR₃)₄ (R = Ph, *p*-CH₃C₆H₄, *p*-CH₃OC₆H₄) were obtained by the (CH₃CH₂)₃Al reduction of Pd(acac)₂ in the presence of SbR₃.

Attempted preparation of Pt(SbR₃)₄ by standard procedures for preparing Pt(PPh₃)₄²⁰ also failed. When the preparation was attempted by the (CH₃CH₂)Al reduction of Pt(acac)₂, a very air-sensitive red oil was obtained in addition to a large amount of black metallic material. No triarylstibineplatinum complex could be isolated.

Reaction of Pd(PPh₃)₄ with excess SbR₃ (R = Ph, *p*-CH₃C₆H₄) resulted in a white powder which analyzed as Pd(PPh₃)₂(SbR₃)₂, in agreement with the formulation proposed in a previous study for R = Ph.¹¹ Pd(PPh₃)₂(SbR₃)₂ was also isolated from the reaction of Pd(PPh₃)₂(CS₂) with SbR₃.

Reaction of Pt(PPh₃)₃ with excess SbR₃ (R = Ph, *p*-CH₃C₆H₄) resulted in a yellow powder which analyzed as Pt(PPh₃)₂(SbR₃)₂—which is not in agreement with the previous formulation, Pt(PPh₃)₃(SbPh₃), obtained from Pt(PPh₃)₄.¹¹ Pt(PPh₃)₂(SbR₃)₂ was also isolated from the reaction of Pt(PPh₃)₂(C₂H₄) or Pt(CS₂)(PPh₃)₂ with SbR₃.

The reaction of Pt(PPh₃)₂(C₂H₄) with Sb(SPh)₃ resulted in the isolation of Pt(PPh₃)₂[Sb(SPh)₃]₂ the first reported thioantimonite-transition metal complex. The complex is insoluble in all common organic solvents and thus its ¹H NMR spectrum could not be obtained.

The zerovalent triarylstibine complexes of nickel, platinum, and palladium were found to have air stability similar to that of the platinum- and palladium-phosphine analogues whereas the nickel-triarylphosphine complexes are more reactive.


¹H NMR Data. ¹H NMR data for solutions of the free ligands and isolated complexes are given in Table II. Examination of the chemical shifts of the M(SbR₃)₄ complexes shows an upfield shift of the aromatic protons and a corre-

sponding though smaller upfield shift in the methyl or methoxy resonances vs. the free ligands.

Addition of free ligand to a solution of any of the $M(\text{SbR}_3)_4$ ($M = \text{Pd}, \text{Ni}$) complexes simply added the resonances of the ligands to the spectra, indicating little dissociation and slow exchange or type I behavior as discussed by Tolman.⁷ Analogous $P(p\text{-CH}_3\text{C}_6\text{H}_4)_3$ complexes exhibit extensive dissociation and fast exchange,⁵ i.e., type IV behavior.⁷

The steric bulk of the triarylphosphine and stibine ligands can be compared by methods previously outlined.^{6,7} The covalent radius of the coordinated antimony is about 0.3 Å larger than for phosphorus³¹ and results in a lengthening of the $M\text{--Sb}$ and Sb--C bonds such that the cone angle for the SbPh_3 is $<132^\circ$ vs. that for PPh_3 , 145° . The steric bulk is more similar to that of $P(p\text{-CH}_3\text{C}_6\text{H}_4\text{O})_3$, 128° , which also exhibits type I behavior.²²

The ^1H NMR spectrum of $\text{Pd}(\text{PPh}_3)_2(\text{SbR}_3)_2$, $R = p\text{-CH}_3\text{C}_6\text{H}_4$, exhibited a single methyl resonance (peak width at half-height 7 Hz) and a very complex phenyl region. Examination of the phenyl region indicated the presence of several species, consistent with the redistribution of $\text{Pd}(\text{PPh}_3)(\text{SbR}_3)_2$ in solution



The observation of a single broadened methyl resonance is consistent with nearly identical methyl chemical shifts for the species present. The material obtained from solution analyzing as $\text{Pd}(\text{PPh}_3)_2(\text{SbR}_3)_2$ could be either a single product or a complex mixture, $\text{Pd}(\text{PPh}_3)_n(\text{SbR}_3)_{4-n}$ ($n = 4-0$), in appropriate ratios.

The ^1H NMR spectrum of $\text{Pt}(\text{PPh}_3)_2(\text{SbR}_3)_2$ showed methyl resonances at τ 7.73 and 7.77. The aromatic region was again more complex than expected for a single species. ^1H NMR spectra of the complexes obtained from $\text{Pt}(\text{PPh}_3)_3$, $\text{Pt}(\text{CS}_2)(\text{PPh}_3)_2$, or $\text{Pt}(\text{PPh}_3)_2(\text{C}_2\text{H}_4)$ were identical. An equilibrium similar to that for $\text{Pd}(\text{PPh}_3)_2(\text{SbR}_3)_2$ must again be proposed.

Triarylstibinerhodium(I) complexes favor five-coordination in the solid state whereas analogous phosphine complexes are four-coordinate.¹² We now find that $M(\text{SbR}_3)_4$ ($M = \text{Pd}, \text{Ni}$) are not extensively dissociated in solution, unlike the analogous phosphine complexes. These observations can be rationalized on the basis of steric considerations. Measurements of cone angles indicate the steric nature of SbR_3 is closer to that of P(OR)_3 than of PR_3 . It appears that, compared with triarylphosphine ligands, the lower steric requirements of the triarylstibine ligands allow higher coordination and do not force dissociation in such systems.

Registry No. $\text{Ni}[\text{Sb}(p\text{-CH}_3\text{C}_6\text{H}_4)_3]_4$, 57527-33-0; $\text{Ni}[\text{Sb}(p\text{-CH}_3\text{OC}_6\text{H}_4)_3]_4$, 57527-34-1; $\text{Pd}[\text{SbPh}_3]_4$, 23854-68-4; $\text{Pd}[\text{Sb}(p\text{-CH}_3\text{C}_6\text{H}_4)_3]_4$, 57527-39-6; $\text{Pd}[\text{Sb}(p\text{-CH}_3\text{OC}_6\text{H}_4)_3]_4$, 57527-40-9; $\text{Pd}(\text{PPh}_3)_2[\text{Sb}(p\text{-CH}_3\text{C}_6\text{H}_4)_3]_2$, 57527-35-2; $\text{Pd}(\text{PPh}_3)_2(\text{SbPh}_3)_2$, 14263-83-3; $\text{Pt}(\text{PPh}_3)_2(\text{SbPh}_3)_2$, 57527-36-3; $\text{Pt}(\text{PPh}_3)_2[\text{Sb}(p\text{-CH}_3\text{C}_6\text{H}_4)_3]_2$, 57527-37-4; $\text{Pt}(\text{PPh}_3)_2[\text{Sb}(\text{SPh})_3]_2$, 57527-38-5; $\text{Ni}(\text{acac})_2$, 3264-82-2; $\text{Ni}(\text{COD})_2$, 1295-35-8; $\text{Pd}(\text{acac})_2$, 14024-61-4; $\text{Pd}(\text{PPh}_3)_4$, 14221-01-3; $\text{Pd}(\text{PPh}_3)_2\text{CS}_2$, 57527-52-3; $\text{Pt}(\text{PPh}_3)_3$, 13517-35-6; $\text{Pt}(\text{PPh}_3)_2\text{CS}_2$, 50588-64-2; $\text{Pt}(\text{PPh}_3)_2(\text{C}_2\text{H}_4)$, 12120-15-9; AlEt_3 , 97-93-8.

References and Notes

- (1) To whom correspondence should be addressed at Dow Chemical U.S.A., New England Laboratory, Wayland, Mass. 01778.
- (2) L. Malatesta and M. Angoletta, *J. Chem. Soc.*, 1186 (1957).
- (3) L. Malatesta and C. Cariello, *J. Chem. Soc.*, 2323 (1958).
- (4) G. Wilke, E. W. Miller, and M. Kroner, *Angew. Chem.*, 73, 33 (1961).
- (5) C. A. Tolman, W. C. Seidel, and D. H. Gerlach, *J. Am. Chem. Soc.*, 94, 2669 (1972).
- (6) C. A. Tolman, *J. Am. Chem. Soc.*, 92, 2956 (1970).
- (7) C. A. Tolman, W. C. Seidel, and L. W. Gosser, *J. Am. Chem. Soc.*, 96, 53 (1974).
- (8) W. Kuran and A. Musco, *Inorg. Chim. Acta*, 12, 187 (1975).
- (9) G. Wilke, E. W. Mueller, M. Kroner, P. Heimbach, and M. Breil, French Patent 1320729 (Mar 15, 1963).

- (10) B. F. G. Johnson, T. Keating, J. Lewis, M. S. Subramanian, and D. A. White, *J. Chem. Soc. A*, 1793 (1969).
- (11) S. Takahashi, K. Sonogashira, and N. Hagihara, *Nippon Kagaku Zasshi*, 87, 610 (1966).
- (12) P. E. Garrou and G. E. Hartwell, *J. Organomet. Chem.*, 69, 445 (1974).
- (13) P. E. Garrou and G. E. Hartwell, *Inorg. Chem.*, 14, 194 (1975).
- (14) B. Bogdanovic, M. Kroner, and G. Wilke, *Justus Liebig's Ann. Chem.*, 699, 1 (1966).
- (15) D. R. Coulson, *Inorg. Synth.*, 13, 121 (1972).
- (16) R. Ugo, F. Cariati, and G. LaMonica, *Inorg. Synth.*, 11, 105 (1968).
- (17) C. D. Cook and G. S. Jauhal, *J. Am. Chem. Soc.*, 90, 1464 (1968).
- (18) T. B. Brill and G. G. Long, *Inorg. Chem.*, 11, 225 (1972).
- (19) T. B. Brill and N. C. Campbell, *Inorg. Chem.*, 12, 1884 (1973).
- (20) F. R. Hartley, *Organomet. Chem. Rev.*, 6, 119 (1970).
- (21) J. T. Mague, *Inorg. Chem.*, 9, 1610 (1970).
- (22) Recent work²³ assumes cone angles for AsPh_3 and SbPh_3 are similar to that of PPh_3 which is obviously invalid.
- (23) L. E. Manzer and C. A. Tolman, *J. Am. Chem. Soc.*, 97, 1955 (1975).

Contribution from the Department of Chemistry, University of Nevada, Reno, Nevada 89507, and the Life Sciences Division, Stanford Research Institute, Menlo Park, California 94025

Investigation of the Utility of "Virtual Coupling" in the $^{13}\text{C}\{^1\text{H}\}$ Nuclear Magnetic Resonance Spectra of Bis-Phosphite Complexes of Palladium and Platinum. Algebraic Cancellation of Spin-Spin Coupling

A. W. Verstuyft,^{1a} J. H. Nelson,^{*1a} and L. W. Cary^{1b}

Received July 23, 1975

AIC50532I

Following the early observation of "virtual coupling" in the ^1H NMR spectra of bis-organophosphorus compounds by King² and by Jenkins and Shaw,³ this phenomenon has become a fundamental tool for geometry assignment. Several hundred articles have appeared in which it has been utilized to assign the geometry of transition metal-phosphine complexes. Unfortunately, there are a large number of compounds for which ^1H NMR virtual coupling is not applicable. The method generally cannot be applied unless the phosphorus ligand bears an α -methyl, α -methoxy, α -methylene, or ortho methylene group which is not coupled to other protons. This excludes such commonly utilized ligands⁴ as triphenylphosphine, triethylphosphine, and triethyl phosphite for which the method is applicable only in fortuitous circumstances.⁵ Because of this limitation as well as the increasing availability of Fourier transform spectrometers $^{13}\text{C}\{^1\text{H}\}$ NMR has been briefly investigated for similar use.⁶ In several instances it has been shown that ^{13}C NMR can provide structural information where ^1H NMR cannot.⁷ Yet few systematic investigations have appeared. In the course of our studies, we noted that the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of *cis*-[(MeO)₃P]₂PdCl₂ was a singlet instead of the anticipated multiplet.⁶ We had also observed other instances of the apparent lack of phosphorus-carbon coupling, e.g., in the aryl groups of arylphosphine complexes.⁸ In order to seek an explanation for this phenomenon and to investigate the general utility of $^{13}\text{C}\{^1\text{H}\}$ NMR for geometry assignments we undertook an investigation of the $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of complexes of the type L_2MCl_2 , where L is a phosphite or phosphine and M is Pd(II) or Pt(II). The chosen ligands are ones for which ^1H virtual coupling for the most part is not applicable. Furthermore, no previous $^{13}\text{C}\{^1\text{H}\}$ studies of phosphite complexes have been reported. For comparison, a few compounds of the type *cis*-LL'₂PdCl₂ were also investigated.⁹ While this work was being completed, a similar study of phosphine complexes appeared.^{7b} Our results on the