# Total Synthesis of Spatol and Other Spatane Diterpenes

## Robert G. Salomon,\* Basudeb Basu, Subhas Roy, and Navzer D. Sachinvala

Contribution from the Department of Chemistry, Case Western Reserve University, Cleveland, Ohio 44106-2699. Received September 6, 1990

Abstract: Total syntheses of three spatane diterpenes stoechospermol (1), 5(R), 15(R), 18-trihydroxyspata-13, 16(E)-diene (2), and (+)-spatol (3) were accomplished from a common intermediate, diol 7. The total synthesis established as R the absolute configuration at the 15-position in 2. Novel stereospecific transformations of 2,3-epoxy-1,4-diols into vicinal diepoxides were demonstrated and exploited for assembling the sensitive allylic diepoxide in the side chain of spatol. This new synthetic method allows the conversion of both 1,2-threo-2,3-trans- and 1,2-erythro-2,3-trans-2,3-epoxy-1,4-diols into vicinal 1,2-cis-2,3erythro-1,3-diepoxides. Unexpected stability toward hydroxide anion was found for the allylic diepoxide functional array. This observation provides presumptive evidence that acid catalysis is operative in the epoxide cleaving substitution reaction of spatol by the weakly nucleophilic chloride anion that gives chlorohydrin 4. A proclivity for erythro-selective epoxidation of allylic silyl ethers was found. The utility of C-silyl substituents for reversing this stereoselectivity, i.e., favoring threo-selective epoxidation of allylic silyl ethers, was established.

### Introduction

The isolation<sup>1</sup> of stoechospermol (1) was soon followed by the discovery of a family of marine diterpenes that possesses a structurally novel carbon skeleton.<sup>2</sup> Besides simple dienes as in



1, the side chains of most spatane diterpenes incorporate one or more hydroxyl groups as in 2. A unique allylic vicinal diepoxide occurs in the side chain of spatol (3). The high electrophilicity of this functional array is evidenced by the cleavage of an epoxide ring by chloride. Thus, esterification of 3 by treatment with p-bromobenzoyl chloride and pyridine is accompanied by an intriguing attack by the weakly nucleophilic chloride anion at the 17-position of the allylic diepoxide producing 4. The reactive diepoxide in spatol is reminiscent of the electrophilic vicinal diepoxide in the cytotoxin crotepoxide<sup>3</sup> and the vicinal triepoxide in the antileukemic diterpenes triptolide and tripdiolide.<sup>4</sup> However, while the epoxy groups in these natural products are confined to rigid cyclohexane rings, those in spatol are located in a flexible acyclic side chain. The reactivity of the allylic diepoxide side chain is almost certainly at the heart of spatol's biological activities. Spatol inhibits mitosis of the fertilized sea urchin egg and cell division in human T242 melanoma and 224C astrocytoma neoplastic cell lines.<sup>2</sup> However, investigations of its chemotherapeutic

- (2) (a) Gerwick, W. H.; Fenical, W.; Van Engen, D.; Clardy, J. J. Am.
  Chem. Soc. 1980, 102, 7991. (b) Gerwick, W. H.; Fenical, W.; Sultanbawa, M. U. S. J. Org. Chem. 1981, 46, 2233. (c) Gerwick, W. H.; Fenical, W. J.
  Org. Chem. 1983, 48, 3325. (d) Ravi, B. N.; Wells, R. J. Aust. J. Chem. 1982, 35, 129.
- 101, 5679.

utility are hampered by the difficulty of obtaining sufficient quantities of spatol from natural sources. This need and spatol's structural and functional complexity make the development of a practical total synthesis a worthy challenge.

We now report the total syntheses of three spatane diterpenes stoechospermol (1), 5(R), 15(R), 18-trihydroxyspata-13, 16(E)diene (2), and (+)-spatol (3) from a common intermediate, diol 7. This diol is available in 16% overall yield in 21 steps from cyclopent-2-en-1-one (5) and 6-methylbicyclo[2.2.1]hept-5-en-2-one (6). Our stereocontrolled synthesis<sup>5</sup> features  $2\pi + 2\pi$ 





photocycloaddition of 5, as an A-ring precursor, with a carbonyl-masked derivative of 6<sup>6</sup> as a temporarily bridged C-ring precursor that incorporates the hydroxyl substituent at position 5 in latent form. Efficient, virtually quantitative resolution of an early intermediate provides homochiral 7 in 8% yield from racemic 6.

### **Results and Discussion**

Stoechospermol. Completion of the spatane skeleton requires the addition of five carbons to the side chain of diol 7. The first total synthesis<sup>7</sup> of a spatane diterpene was accomplished by selective bromodehydroxylation of the primary allylic hydroxyl group in  $(\pm)$ -7 followed by a copper(I) iodide catalyzed coupling of the resulting allylic bromide 8 with an excess of prenylmagnesium chloride.<sup>8</sup> The <sup>1</sup>H NMR spectrum of the product, (±)-stoechospermol (1), is identical with that of (+)-1 isolated from Stoechospermum marginatum.<sup>1,2</sup>



5(R),15(R),18-Trihydroxyspata-13,16(E)-diene. For the first total synthesis of trihydroxylated spatane diterpene 2, the vinyl

(5) Salomon, R. G.; Sachinvala, N. D.; Roy, S.; Basu, B.; Raychaudhuri, S. R.; Miller, D. B.; Sharma, R. B. J. Am. Chem. Soc. 1991, 113, preceding article in this issue.

- (6) Lal, K.; Salomon, R. G. J. Org. Chem. 1989, 54, 2628.
- (7) For a preliminary report of this work see: Salomon, R. G.; Sachinvala, N. D.; Raychaudhuri, S. R.; Miller, D. B. J. Am. Chem. Soc. 1984, 106, 2211.
- (8) Kwart, H.; Miller, R. K. J. Am. Chem. Soc. 1954, 76, 5403.

<sup>(1)</sup> Fernandes, S. L.; Kamat, S. Y.; Paknikar, S. K. Tetrahedron Lett. 1980, 21, 2249.

Scheme I



trans nucleophile **10** was generated stereoselectively from 2methyl-3-butyn-2-ol through a vinyl tin intermediate **9**. The



 $\alpha,\beta$ -unsaturated aldehyde (+)-11, available from diol (+)-7 in 96% yield by MnO<sub>2</sub> oxidation, underwent exclusive 1,2-addition of 10 to give two epimeric alcohols 12 and 13 in 91% yield.



Desilylation of the minor adduct 12 delivered the natural product 2 with  $[\alpha]^{22}_D - 32.2^\circ$  (reported<sup>2</sup>  $[\alpha]^{22}_D - 33.4^\circ$ ). Desilylation of the major adduct 13 afforded 14, which exhibits  $[\alpha]^{22}_D - 1.01^\circ$  and is an epimer of the natural product at position 15. The configuration of 2 at position 15 had not been established previously. Our assignment of the *R* configuration to this center in the natural product is based on a conversion of the epimer 14 into spatol (vide infra).

Vicinal Diepoxides from 2,3-Epoxy-1,4-diols. The unique allylic vicinal diepoxide in spatol (3) constitutes by far the most challenging aspect of any synthetic approach to this target.<sup>9</sup> Our strategy for assembling the spatol side chain was based on the premise that an appropriately activated derivative, e.g., *threo*,-*trans*-15 of a 15,16-threo-16,17-trans (spatane numbering) 2,3-



epoxy-1,4-diol (functional array numbering), could stereospecifically interconvert with the corresponding derivative of a 3,4-epoxy-1,2-diol, e.g., *threo,erythro*-16 by a reversible intra-

Scheme II



molecular  $S_N^2$  displacement, a Payne rearrangement.<sup>10</sup> Subsequent irreversible intramolecular  $S_N^2$  displacement of a leaving group from the 15-position would deliver the 15,16-cis-16,17-erythro diepoxide array, as in *cis,erythro*-17, required for the spatol side chain. A similar synthesis of *cis,threo*-17 from *threo,cis*-15 might be derailed by a competing cyclization to an epoxy furan 18. However, furan formation from *threo,trans*-15 would not



interfere with the generation of cis, erythro-17 owing to the ring strain that would be created upon formation of a trans fused epoxy furan. A similar double  $S_N 2$  displacement process should deliver *trans,erythro-17* from *erythro,trans-15* via *erythro,erythro-16*. To test this hypothesis, model threo and erythro *trans-2,3-ep*oxy-1,4-diols were assembled as outlined in Scheme I. To assure epoxidation of the appropriate C=C bond, the tertiary hydroxyl in 21 was depyranylated and the secondary hydroxyl in 22 was selectively silylated to provide 23. Vanadium-catalyzed epoxidation produced a 4:1 mixture of erythro and threo epoxides 24e and 24t, respectively. Desilylation delivered the corresponding epoxy diols.

In an early attempt at generating a vicinal diepoxide, the monomesylate derivative 26e of the erythro diol 25e was boiled under reflux in 2-propanol in the presence of solid  $K_2CO_3$ . Payne rearrangement followed by heterocyclization did not occur. Rather, intermolecular  $S_N 2$  displacement produced the isopropyl ether 27 of three epoxy diel 25t (Scheme II). To explore the utility of such intermolecular S<sub>N</sub>2 displacement for interconversion of threo and erythro epoxy diols 25t and 25e, the monomesylate 26e was heated with aqueous bicarbonate. A low yield of the three epoxy diol 25t was obtained. A better route to 25t was devised, exploiting intermolecular  $S_N 2$  displacement of the mesylate in 26e by levulinate (Scheme II). Whereas reaction in 2-methyl-2propanol solvent favored net S<sub>N</sub>2' substitution delivering 28 virtually quantitatively, in most solvents  $S_N 2$  substitution produced the threo levulinate ester 29 (Table I). Reaction in DME provided 29 exclusively in 91% yield. Removal of the levulinate ester to provide three epoxy diel 25t in 85% yield was readily accomplished

<sup>(9)</sup> For an alternative strategy for construction of the spatol allylic diepoxide side chain, see: Tanaka, M.; Tomioka, K.; Koga, K. Tetrahedron Lett. **1985**, 26, 6109.

<sup>(10)</sup> Payne, G. B. J. Org. Chem. 1962, 27, 3819.

 
 Table I. Solvent Effects on Reaction of 26e with Tetramethylammonium Levulinate

Solvent	temp (°C)	time (h)	29:28	yield (%)
CHCl <sub>3</sub>	61	30	68:32	98
t-BuOH	80	4	0:100	96
CH <sub>3</sub> CN	80	4	65:35	94
DMF	80	3.5	100:0	77
DME	80	4.5	100:0	91

under mild conditions upon treatment of **29** with sodium borohydride.<sup>11</sup>

Since the tertiary hydroxyl group in 26e is apparently not sufficiently nucleophilic to displace the epoxy leaving group, conditions were sought that would generate an alkoxide from the tertiary hydroxyl. Treatment of the erythro monomesylate 26e with t-BuOK in t-BuOH promoted a clean, stereospecific rearrangement and heterocyclization to deliver the diepoxide trans, erythro-31. Similar treatment of the threo monomesylate



26t provided *cis,erythro-31*, which contains an allylic diepoxide resembling the side chain of spatol. Considering the substitution reaction of monomesylate 26e with aqueous bicarbonate to give 25t (Scheme II), we expected a similar substitution with tetrabutylammonium hydroxide. Instead, a high yield of diepoxide trans.ervthro-31 was obtained. The unexpected stability of this allylic diepoxide product toward hydroxide is especially interesting in view of the epoxide cleaving substitution reaction of spatol with the weakly nucleophilic chloride anion to give chlorohydrin 4 (vide supra). Apparently, the latter reaction is an acid-catalyzed epoxide opening that is induced by pyridinium hydrochloride, a byproduct of the acylation with p-bromobenzoyl chloride in the presence of pyridine. It remains an interesting question why chloride attacks the 17-position in spatol to generate 4 rather than inducing cleavage of the presumably weaker allylic C-O bond at position 15. That the reactivity of the side chain functionality is important for spatol's biological activity is supported by the fact that both of the isomeric allylic diepoxides 31 are biologically active.<sup>12</sup>

Recall that epoxidation of 23 favored the erythro over the threo epoxy TBDMS ether by 4:1 (Scheme I). Furthermore, the threo mesyloxy epoxide 26t is the precursor of the cis,erythro diepoxide required for spatol. Therefore, we sought a route from the *erythro* epoxy alcohol 25e to an activated derivative of the *threo* epoxy alcohol, i.e., activation with concomitant inversion of configuration.



This can be accomplished by the Still modification of the Mitsunobu reaction.<sup>13</sup> Treatment of 25e with  $Zn(OTs)_2$ , diethyl

Scheme III



Scheme IV



azodicarboxylate, and triphenylphosphine gave tosylate **32**, which upon treatment with *t*-BuOK in *t*-BuOH, afforded allylic diepoxide *cis,erythro*-**31** in only 66% yield together with an unidentified byproduct (17%). Solvent strongly influences the ratio of these products (see Experimental Section) with the desired diepoxide being the only product using DMF as reaction solvent.

Threo Epoxy Alcohols. Since the epoxidation of 23 favored generation of the erythro over the threo epoxy TBDMS ether by 4:1 (Scheme I) and since the threo mesyloxy epoxide 26t is the precursor of the cis,erythro diepoxide required for spatol, we sought an epoxidation route favoring threo epoxy TBDMS ethers. The possible utility of a C-silyl substituent as a stereocontrol element was explored (Scheme III). Indeed, molybdenum-catalyzed epoxidation of 36 favors the threo epoxide 37t in contrast with the epoxidation of 23, which favored the erythro epoxide 24e. This inversion of stereoselectivity by a C-silyl substituent is analogous to that observed previously for similar epoxidations of secondary allylic  $alcohols^{14}$  as opposed to the TBDMS ethers.

<sup>(11)</sup> Hassner, A.; Strand, G.; Rubinstein, M.; Patchornik, A. J. Am. Chem. Soc. 1975, 97, 1614.

<sup>(12)</sup> The ability of these allylic diepoxides to inhibit microtubule assembly and the mitosis of sea urchin eggs was examined. *cis,erythro-31* showed 36% inhibition at 16  $\mu$ g/mL in the sea urchin egg assay and 74% inhibition of microtubule assembly at 70  $\mu$ g/mL, and *trans,erythro-31* showed 28% inhibition at 16  $\mu$ g/mL in the sea urchin egg assay and 64% inhibition of microtubule assembly at 70  $\mu$ g/mL. For details of these techniques, see: Murthi, K. K.; Salomon, R. G.; Sternlicht, H. *Prostaglandins* **1990**, 39, 611.

<sup>(13)</sup> Galynker, I.; Still, W. C. Tetrahedron Lett. 1982, 23, 4461.

<sup>(14)</sup> For erythro stereoselectivity in metal-catalyzed epoxidations of allylic alcohols, see: (a) Mihelich, E. D. Tetrahedron Lett. 1979, 4729. (b) Rossiter, B. E.; Verhoeven, T. R.; Sharpless, K. B. Tetrahedron Lett. 1979, 4733. (c) Takai, K.; Oshim, K.; Nozaki, H. Tetrahedron Lett. 1980, 21, 1657. (d) Martin, V. S.; Woodard, S. S.; Katsuki, T.; Yamada, Y.; Ikeda, M.; Sharpless, K. B. J. Am. Chem. Soc. 1981, 103, 6237. For threo stereoselectivity in metal catalyzed epoxidations of C-silyl allylic alcohols, see: (e) Tomioka, H.; Suzuki, T.; Oshima, K.; Nozaki, H. Tetrahedron Lett. 1982, 23, 3387.

Scheme V



The utility of C-silyl groups as stereocontrol elements in metalcatalyzed epoxidations depends on their stereospecific removal. Treatment of the silyl epoxides 37 with tetrabutylammonium fluoride delivered the corresponding epoxides 38 with retention of configuration as expected.<sup>15</sup>

(+)-Spatol. An early attempt at generating the allylic diepoxide side chain in the spatane ring system is outlined in Scheme IV. A selectively disilylated derivative 39 was readily available from triol 14. Vanadium-catalyzed epoxidation of 39 produced a 1:6 mixture of disilylated epoxides 40t and 40e in good yield. The corresponding triols 41 were obtained upon treatment with  $Bu_4NF$ . However, generation of an inverted tosylate from the major epimer 41e followed by treatment with base produced an allylic diepoxide 42 with a rearranged tricyclic nucleus. The *cis,anti,cis*-tricyclo[5.3.0.0<sup>2.5</sup>]decane nucleus of 42 is formed by a Wagner-Meerwein rearrangement of the *cis,anti,cis*-tricyclo[5.3.0.0<sup>2.6</sup>]decane nucleus of 41e, apparently owing to unintended activation of the 5-hydroxyl that accompanied the desired activation of the hydroxyl at position 15.

These results suggested that derivatives of epoxy triols 41e and 41t with the hydroxyl at position 5 masked would be desirable for generation of the spatol side chain without accompanying rearrangement of the tricyclic nucleus. The lability of the allylic diepoxide array in spatol (3) under acidic conditions limited the choice of derivatives to those with masking groups that would be removable under neutral or basic reaction conditions. The further requirement for stability toward a vinyllithium reactant and the presence of unsaturation in the synthetic target agreed with the choice of *p*-methoxybenzyl (MBn) ether derivatives. The MBn masking group is removable under mild conditions by oxidative cleavage with DDQ.<sup>16</sup> The MBn derivatives erythro-50b and threo-50b of 41e and 41t were prepared from diol (+)-7 as outlined in Scheme V. Another isomer, threo-50a was prepared from erythro-50b by inversion of configuration at position 15 through levulinate ester 51.

An allylic cis diepoxide (-)-52 was generated upon treatment of *threo*-50a with methanesulfonic anhydride and *p*-(dimethylamino)pyridine followed by reaction of the intermediate secondary monomethanesulfonate with t-BuOK. The same diepoxide was



also generated more directly from *erythro*-50b in better overall yield upon treatment with Zn(OTs)<sub>2</sub>, diethyl azodicarboxylate, and triphenylphosphine followed by reaction of the intermediate secondary tosylate with *t*-BuOK. Removal of the MBn masking group by treatment with DDQ converted (-)-52 into an allylic cis diepoxide (-)-53. That (-)-53 was not spatol (3) was evident from its optical activity,  $[\alpha]_D = -10.0^\circ$  in contrast with  $[\alpha]_D = +45.6^\circ$  reported for the natural product. Small chemical shift differences, e.g., vinyl <sup>1</sup>H NMR resonances at  $\delta$  5.14 and 5.09, confirmed that (-)-53 is epimeric at positions 15, 16, and 17 with (+)-spatol (3), which exhibits vinyl resonances at  $\delta$  5.13 and 5.02.



Similar stereospecific transformations of *threo*-**50b** produced an allylic cis diepoxide that was identical with the natural product. Each resonance in the <sup>1</sup>H NMR spectrum of synthetic (+)-spatol (3) coincided within 0.01 ppm with a spectrum of an authentic sample of natural spatol.<sup>17</sup>

**Prospects for a Practical Total Synthesis of (+)-Spatol.** Two approaches for achieving a practical synthesis of spatol are suggested by our studies. One route exploits the stereoselectivity discovered for nucleophilic additions to the aldehyde (+)-46. The



13:1 diastereoselectivity favoring generation of the S configuration at position 15 during the (+)-46 to 47 conversion (Scheme V) did not result in an efficient, practical synthesis of spatol since epoxidation of the major epimer, (-)-49b, is erythro-selective. Only the minor epoxidation product from (-)-49b, threo-50b, is a precursor for spatol. We did demonstrate the feasibility of using a C-silyl substituent to promote threo-selective epoxidation of an allylic TBDMS ether (Scheme III). However, the route established for assembling the C-silylated allylic alcohol 36 (Scheme III) is too long to provide a workable solution to this problem. If a vinyllithium 55 were available, a more convergent route to the requisite C-silyl derivative 56 would be possible. Diastereoselective generation of the S configuration at the 15-position in 56 is anticipated for the addition of 55 to aldehyde (+)-46. Threo-selective epoxidation, as for 36 (Scheme III), would then

 <sup>(15)</sup> Chan, T. J.; Lau, P. W. K.; Li, M. P. Tetrahedron Lett. 1976, 2667.
 (16) Oikawa, W.; Yoshioka, T.; Yonemitsu, O. Tetrahedron Lett. 1982, 23, 885.

<sup>(17)</sup> All <sup>1</sup>H and <sup>13</sup>C NMR spectra of (-)-53 and (+)-spatol (3), including those of an authentic sample of natural spatol, were compared in CDCl<sub>3</sub> solutions on the same Brucker 9.4 Tesla MSL-400 FT-NMR spectrometer.

provide 57 from which the intermediate *threo*-50b in our spatol synthesis would be available by a stereospecific, one-pot, double desilylation as demonstrated earlier for the 37t to 38t conversion.

A second approach for achieving a practical synthesis of spatol that is suggested by our studies depends on the availability of a synthetic equivalent of the hypothetical  $\alpha$ -lithioepoxide **58**.<sup>18</sup>



Efficient diastereoselective generation of *threo*-50b can be anticipated from the addition of homochiral 58 to aldehyde (+)-46. Stereospecific conversion of *threo*-50b into spatol was demonstrated earlier. Furthermore, application of the novel transformations of 2,3-epoxy-1,4-diols reported herein to the conversion of the expected minor epimeric adduct, *erythro*-50a, into an allylic diepoxide should also produce (+)-spatol (3) stereospecifically. This exceptionally convergent strategy for construction of the allylic diepoxide side chain of spatol is especially attractive since it does not depend on stereocontrol at position 15. These plans for optimizing our synthesis are the subject of ongoing research.

## Experimental Section<sup>19</sup>

Allylic Bromide 8. A solution of the 5-hydroxy allylic alcohol 7<sup>5</sup> (9.6 mg, 0.040 mmol) in dry acetonitrile (225  $\mu$ L) was mixed with triphenylphosphine (12.2 mg, 0.047 mmol, 1.15 equiv). To this mixture was added carbon tetrabromide (44.1 mg, 0.133 mmol, 3.3 equiv), and the mixture was stirred at room temperature for 12 h. Solvents were then evaporated from the reaction mixture under a stream of dry nitrogen. Flash chromatography on a 10 × 150 mm column of silica gel 230-400 mesh with 1% ethyl acetate in methylene chloride afforded the allylic bromide 8 (7.6 mg, 0.025 mmol, 63%): mp 72-73 °C; <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  0.85 (3 H, d, J = 6.1 Hz), 0.99 (3 H, s), 1.19-2.05 (10 H), 2.23 (1 H, dd, J = 5.4, 14.5 Hz), 3.35 (1 H, m), 3.75 (1 H, d, J = 4.7 Hz), 3.84 (2 H, dd, J = 10.2, 15.6 Hz), 4.97 (1 H, s), 5.26 (1 H, s).

(+)-Stoechospermol (1). To copper(I) iodide (1.4 mg, 0.007 mmol) at 0 °C was added prenylmagnesium chloride<sup>8</sup> in tetrahydrofuran (0.7 M, 100  $\mu$ L, 0.070 mmol). To this reagent at -50 °C<sup>20</sup> was added the allylic bromide 8 (3.0 mg, 0.01 mmol) in tetrahydrofuran (50  $\mu$ L) and the resulting mixture allowed to warm to room temperature and stirred 14 h. The reaction mixture was quenched with aqueous ammonium chloride (50 mg in 300  $\mu$ L water) then concentrated aqueous ammonium hydroxide (50  $\mu$ L) and extracted with ether (5 mL). The aqueous layer was reextracted with ether (5 mL), and the ether extracts were then washed serially with water. The ether layers were combined and concentrated in vacuo to give a product mixture that was flashed chromatographed on a  $10 \times 150$  mm column of 230-400 mesh silica gel with 1% ethyl acetate in methylene chloride. The product, racemic 5hydroxyspata-13,17(E)-diene (1) (1.63 mg, 5.6  $\mu$ mol, 56% yield) exhibited a <sup>1</sup>H NMR spectrum identical with that found for the natural product (+)-1 isolated from S. marginatum.<sup>1</sup>

2-Methyl-2-{(triethylsilyl)oxy}-3-butyne. Triethylsilyl chloride (3.0 g, 20 mmol) was added dropwise to a stirring mixture of 2-methyl-3-butyn-2-ol (1.0 g, 12 mmol), triethylamine (2.53 g, 25 mmol), and 4-(dimethylamino)pyridine (145 mg, 1.2 mmol) in methylene chloride (25 mL) at 0 °C under nitrogen. After being stirred an additional 1 h at 0 °C, the mixture was washed with water and dried (MgSO<sub>4</sub>). The volatiles were removed under water aspirator reduced pressure and the resulting residue was flash chromatographed (hexane) to give the title compound as a colorless oil (2.35 g, 100%): <sup>1</sup>H NMR  $\delta$  0.46–0.74 (6 H), 0.87–0.98 (6 H), 1.43–1.47 (6 H), 2.36 (H, s). Anal. Calcd for

C<sub>11</sub>H<sub>22</sub>OSi: C, 66.59; H, 11.17. Found: C, 66.43; H, 11.12.

3-Methyl-1-(tri-n-butylstannyl)-3-[(triethylsilyl)oxylbut-1(E)-ene (9). A mixture of 2-methyl-2-[(triethylsilyl)oxy]-3-butyne (1.0 g, 5.1 mmol), tri-n-butyltin hydride (1.6 g, 5.6 mmol), and azo(bisisobutyro)nitrile (20 mg) was irradiated with a sunlamp (110 V, 250 W) for 10 min and then stirred at 90 °C for 2 h. The crude mixture was flash chromatographed (hexane) to yield 9 as a colorless oil (1.76 g, 92%): <sup>1</sup>H NMR  $\delta$  0.48–0.61 (6 H), 0.82-0.96 (24 H), 1.23-1.60 (12 H), 5.86-6.26 (H). Anal. Calcd for C23H50OSiSn: C, 56.44; H, 10.29. Found: C, 56.35; H, 10.32. (+)-5(R),15(R)-Dihydroxy-18-[(triethylsilyl)oxy]spata-13,16(E)diene (12) and (+)-S(R),15(S)-Dihydroxy-18-[(triethylsilyl)oxy]spata-13,16(E)-diene (13). n-Butyllithium (546 µL, 1.37 mmol, 2.5 M in hexanes) was added dropwise to a stirred solution of the vinylstannane 9 (668 mg, 1.36 mmol) in tetrahydrofuran (3 mL) at -78 °C under nitrogen. After being stirred 2 h at -78 °C and 1 h at -10 °C, a solution of the aldehyde (+)-11, (60 mg, 0.256 mmol) in tetrahydrofuran (1 mL) was added dropwise at -78 °C. The reaction mixture was allowed to warm to room temperature over 3 h and then quenched with methanol (500  $\mu$ L). After the volatiles were removed by rotary evaporation, the residue was flash chromatographed to afford a minor epimer 12 (14.2 mg, 13%) as a colorless oil (<sup>1</sup>H NMR δ 0.44-0.60 (6 H), 0.88-0.95 (15 H), 1.28-2.05 (17 H), 2.26 (H, dt, J = 12 and 4 Hz), 2.78-3.0 (H, m), 3.71 (H, d, J = 4 Hz), 4.38 (H, d, J = 8 Hz), 4.90 (H, s), 5.29 (H, s), 5.48 (H, dd, J = 16 and 8 Hz), 5.79 (H, d, J = 16 Hz)) and a major epimer 13 (80 mg, 78%) as a semisolid mass: <sup>1</sup>H NMR δ 0.47-0.60 (6 H), 0.82-0.95 (15 H), 1.27-2.04 (17 H), 2.22 (H, dt, J = 12 and 4 Hz),

3.14-3.22 (H, m), 3.70 (H, d, J = 4 Hz), 4.41 (H, bs), 4.88 (H, s), 5.05 (H, s), 5.60 (H, dd, J = 16 and 4 Hz), 5.75 (H, d, J = 16 Hz). **5(R),15(R),18-Trihydroxyspata-13,16(E)-diene (2).** A mixture of the triethylsilyl ether **12** (12 mg, 0.028 mmol) in THF (1 mL) and tetrabutylammonium fluoride (140  $\mu$ L, 0.14 mmol, 1 M in THF) was stirred at room temperature for 40 h. Volatiles were removed by rotary evaporation, and the residue was flash chromatographed on silica gel with 80% ethyl acetate in hexanes as eluant to give the known triol **2**<sup>i</sup> (8.5 mg, 95%):  $[\alpha]^{22}_{D} -33.2^{\circ}$  (c 1.3, CDCl<sub>3</sub>), reported  $[\alpha]_{D} -33.4$  (c 0.73,

CHCl<sub>3</sub>). (+)-5(R),15(S),18-Trihydroxyspata-13,16(E)-diene (14). A mixture of the triethylsilyl ether 13 (75 mg, 0.173 mmol) in THF (2 mL) and tetrabutylammonium fluoride (1 mL, 1.0 mmol, 1 M in THF) was stirred overnight at room temperature. The volatiles were removed by rotary evaporation, and the residue was flash chromatographed with 80% ethyl acetate in hexanes as eluant to yield the triol 14 (55.1 mg, 99%) as colorless crystals: mp 124-125 °C;  $[\alpha]^{22}_{D}$  +1.01° (c 2.0, CHCl<sub>3</sub>); <sup>1</sup>H NMR  $\delta$  0.85 (3 H, d, J = 6.4 Hz), 0.96 (3 H, s), 1.18-2.10 (18 H), 2.23 (H, dt, J = 16 and 4 Hz), 3.16-3.32 (H, m), 3.72 (H, d, J = 4 Hz), 4.44 (H, d, J = 5.5 Hz), 4.90 (H, s), 5.06 (H, s), 5.68 (H, dd, J = 16 and 4 Hz), 5.84 (H, d, J = 16 Hz). Anal. Calcd for C<sub>20</sub>H<sub>32</sub>O<sub>3</sub>: C, 74.95; H, 10.06. Found: C, 74.91; H, 10.12.

Ethyl 2-Cylopentylacrylate. To a solution of the diethyl cyclopentylmalonate<sup>21</sup> (22.8 g, 0.1 mol) in ethanol (64 mL) at 0 °C was added potassium hydroxide (6.72 g, 0.12 mol) in water (28 mL) very slowly over 1 h. The temperature of the reaction was maintained at 0 °C during the course of the addition and for approximately 2 h after the addition was complete. The reaction mixture was then allowed to attain room temperature and then stirred overnight (10 h). The ethanol was then removed by vacuum transfer to a receiver cooled to -78 °C and the residue partitioned between ether and water. The aqueous layer was washed once more with ether and the ether layers were combined, washed with brine, dried over sodium sulfate, and concentrated in vacuo to recover unreacted starting diethyl cyclopentylmalonate (1.94 g, 8.5 mmol). The aqueous layer was then cooled to -10 °C and acidified to pH 1 at that temperature by slow addition of concentrated hydrochloric acid. Extraction of the aqueous layer with ether ( $6 \times 50$  mL) and drying over anhydrous magnesium sulfate followed by rotary evaporation of the solvent yielded the half-acid (17.8 g, 0.089 mol, 89% yield).

A stock solution containing sodium acetate (8.2 g), acetic acid (296.4 mL), 30 aqueous formalin (216 mL), and diethylamine (freshly distilled, 74.1 mL) was prepared at 10 °C. The previous half-acid (63.1 g, 0.32 mol) was added the stock solution (400 mL) at room temperature. The mixture was immersed in an oil bath preheated to 90 °C. After carbon dioxide evolution had ceased, the reaction mixture was cooled and transferred to a separatory funnel with water and ether. The aqueous phase was then extracted with ether  $(5 \times 50 mL)$  and the organic extracts combined. Removal of acetic acid was accomplished by washing with saturated aqueous sodium bicarbonate solution. The organic phase was then with a 10% solution of hydrochloric acid, then with water, and then with brine. The organic phase was dried over anhydrous magnesium sulfate and concentrated in vacuo. Ethyl 2-cyclopentyl-

<sup>(18)</sup> Silyl-stabilized  $\alpha$ -lithio epoxides might serve as synthetic equivalents of **58** since they are readily available by metallation of the corresponding epoxy silanes (Eisch, J. J.; Galle, J. E. J. Am. Chem. Soc. 1976, 98, 4646. J. Organomet. Chem. 1976, 121, C10; 1976, 41, 2615; 1988, 341, 293) and since desilylation can be accomplished with retention of configuration as in the 37t to 38t conversion.

<sup>(19)</sup> For information on "General" experimental details and "Materials" see the Experimental Section of ref 5.

<sup>(20)</sup> Derguini-Boumechal, F.; Lorne, R.; Linstrumelle, G. Tetrahedron Lett. 1977, 1181.

acrylate thus obtained weighed 28.1 g (0.17 mol, 53% yield): <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  1.3–2.1 (10 H), 2.89 (1 H, quin, J = 8.3 Hz), 4.25 (2 H, q, J = 6.9 Hz), 5.56 (1 H, s), 6.14 (1 H, s). On smaller scale, i.e., 8–10 g, the yields are generally 10% higher. Without further purification, this product was used for preparing 2-cyclopentylallyl alcohol.

2-Cyclopentylallyl Alcohol. To a solution of diisobutylaluminum hydride (12.1 mL, 9.65 g, 68 mmol, 3.0 equiv) in diethyl ether (360 mL) was added ethyl 2-cyclopentylacrylate (3.80 g, 22.6 mmol) dropwise over 10 min at -78 °C. The reaction mixture was then allowed to gradually warm to room temperature over 3 h and stirred for another 2 h. The course of the reaction was monitored by TLC with 10% ethyl acetate in methylene chloride. The  $R_f$  valves of the ester and allyl alcohol were 0.69 and 0.33, respectively. The acrylic ester is UV and iodine active but doesn't stain with vanillin; however, the allyl alcohol shows no activity in UV or iodine but stains a blue-green color with the vanillin indicator. Upon loss of UV activity at  $R_{f}$  0.69, the reaction mixture is cooled to -10 °C and treated very cautiously with an excess of freshly distilled methanol (20 mL) added dropwise over approximately 20 min. The mixture containing a suspension of aluminates is then allowed to attain room temperature, the aluminates are dissolved in 10% hydrochloric acid (20 mL) and extracted immediately into ether, and the phases are separated. The aqueous layer is then reextracted with ether  $(4 \times 20 \text{ mL})$ . The combined organic extracts were washed once with dilute bicarbonate water and brine. Drying over anhydrous magnesium sulfate and concentration in vacuo afforded 2-cyclopentylallyl alcohol (2.84 g, 88% yield). Without further purification, this product was used for preparing 2-cyclopentylacrylaldehyde. A purified sample was recovered from that oxidation by chromatography on silica gel (vida infra): <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  1.0–1.9 (8 H), 2.41 (1 H, quin, J = 8.3 Hz), 4.10 (2 H, s), 4.88 (1 H, s), 4.98 (1 H, s). Anal. Calcd for C<sub>8</sub>H<sub>14</sub>O: C, 76.14; H, 11.18. Found: C, 75.98; H, 11.12.

2-Cyclopentylacrylaidehyde (19). To a solution of the allyl alcohol (5.1 g, 40 mmol) in methylene chloride (160 mL) was added manganese dioxide (24-g portions added every 24 h for 3 days). A total of 72 g (0.83 mol, 20.6 equiv) of manganese dioxide was consumed and the reaction mixture stirred for 4 days. The reaction was monitored by TLC with methylene chloride as the mobile phase.  $R_f$  values for the allyl alcohol and the acryl aldehyde were 0.27 and 0.55. The allyl alcohol shows no UV activity but stains a deep blue-green color with vanillin. The acryl aldehyde is UV active, shows an orange color with dinitrophenylhydrazine, but stains only a very faint navy blue to grey with vanillin indicator. After 4 days, the reaction was still incomplete and a strongly vanillin staining (blue) impurity was developing at  $R_f 0.69$ . The reaction mixture was filtered through a 3-cm thick layer of Celite that was washed with copious amounts of methylene chloride, and the solvent was then removed at 10 °C by rotatory evaporation. Flash chromatography of the oil (5.3 g) thus obtained on a silica gel column  $80 \times 180$  mm with 10%ether in hexanes (acrylaldehyde  $R_f$  0.19, impurity  $R_f$  0.36, and allyl alcohol stays at the origin), yielded the acrylaldehyde 19 (3.97 g, 32 mmol, 80% yield). The column was then eluted with methylene chloride to recover unreacted allyl alcohol (810 mg, 6.4 mmol, 16% recovery). The acrylaldehyde 19 is very sensitive and should be stored under argon at -78 °C. A 70% solution of 19 in hexanes or pentanes (dry, olefin-free) under argon at -78 °C is stable for months: <sup>1</sup>H NMR (200 MHz,  $CDCl_3$ )  $\delta$  1.2-2.1 (8 H), 2.84 (1 H, quin, J = 8.2), 5.95 (1 H, s), 6.25 (1 H, s), 9.56 (1 H, s).

2-Methyl-2-[(tetrahydropyranyl)oxy]-3-butyne. The procedure of Corey, Ulrich, and Fitzpatrick<sup>22</sup> was modified. A mixture of 2methyl-3-butyn-2-ol (8.4 g, 0.1 mol, 9.67 mL), dihydropyran, freshly distilled over sodium (11 mL, 10.1 g, 0.12 mmol, 1.2 equiv), and methylene chloride (84 mL) was cooled to 0 °C and treated with a catalytic amount of p-toluenesulfonic acid (7 mg). The reaction was exothermic, and without the ice-water bath, it was violent. The mixture was stirred and allowed to attain room temperature. The progress of the reaction was monitored by TLC with neat methylene chloride as the eluant;  $R_{f}$ values for the alkynol, dihydropyran, and the tetrahydropyranyl ether were 0.18, 0.38, and 0.53 respectively. The spots were visualized with vanillin indicator. Within 2-2.5 h, all the starting alkynol was consumed. The reaction mixture was then washed with a solution of saturated sodium bicarbonate and then brine and dried over anhydrous potassium carbonate. Methylene chloride and some dihydropyran were removed by rotory evaporation, and the resulting liquid was fractionally distilled to remove excess dihydropyran (85–87 °C (760 mmHg)). The tetrahydropyranyl ether boiled between 120-124 °C (760 mmHg) (15.9 g, 94.5% yield).22

3-Methyl-3-[(tetrahydropyranyl)oxy]-1-(tri-*n*-butylstannyl)-1(*E*)butene. In a 300-mL round-bottom flask, a mixture containing tri-*n*- butyltin hydride (47.5 g, 0.163 mol), 2-[(tetrahydropyranyl)oxy]-2methyl-3-butyne (24.9 g, 0.148 mol) and AIBN (900 mg) was irradiated with a GE 475-W, 110-V sunlamp and then, after ~10 min, immersed in an oil bath preheated to 80 °C. Although the initial reaction is very (often violently) exothermic, heating is necessary to ensure that all of the alkyne is consumed. The use of a reaction flash 4× the total volume of the liquid facilitates control of the reaction. The progress of the reaction was monitored by TLC with 10% ethyl acetate in hexanes as the mobile phase.  $R_f$  values of the hydride, the alkyne, and the vinylstannane were 0.91, 0.48, and 0.65, respectively. The vinyl stannane was less polar than the alkyne and stained a deep purple with vanillin indicator. The reaction went to completion in 3 h. The crude mixture was chromatographed in 20-g batches on a  $120 \times 180$  mm silica gel flash column and eluted with 5% ethyl acetate in hexanes ( $R_f$  trans vinyl stannane 0.40). Removal of solvents in vacuo gave vinylstannane (57 g, 85%).<sup>22</sup> Note that the vinyl stannane can trap a considerable amount of solvents; therefore, it was stirred vigorously at room temperature under high vacuum (0.02 mmHg) to remove all traces of solvents.

6-Cyclopentyl-5-hydroxy-2-methyl-2-[(tetrahydropyranyl)oxy]-3-(E),6-heptadiene (21). To a solution of 3-methyl-3-[(tetrahydropyranyl)oxy]-1-(tributylstannyl)-1(E)-butene (4.59 g, 10 mmol) in THF (15 mL) at -78 °C was added n-BuLi in hexane (2.6 N, 10.5 mmol, 4.43 mL). The mixture was stirred 1 h at -78 °C and 1 h at -5 °C. It was again cooled to -78 °C, and a solution of acrylaldehyde 19 (595 mg, 4.80 mmol) in THF (1 mL) was added dropwise. Aftering the solution was stirred 2 h at -78 °C, excess vinyl anion was destroyed by addition of methanol (5 mL) and the mixture was allowed to attain room temperature. Water (50 mL) was added, and the resulting mixture was extracted with ether. The ether layer was washed with H<sub>2</sub>O and dried (MgSO<sub>4</sub>). Removal of the solvent followed by flash chromatography over silica gel with 10% EtOAc in hexane gave the (tetrahydropyranyl)oxy dienol 21 (994 mg, 75%) as a viscous oil: <sup>1</sup>H NMR δ 1.17-1.82 (21 H), 2.37 (H, quin, J = 8 Hz), 3.32–3.43 (H, m), 3.87–3.96 (H, m), 4.58 (2 H, bs, J = 4.7 Hz), 4.93 (H, s), 5.08 (H, s), 5.56 (H, ddd, J = 3.6, 16 Hz), 5.79 (H, dd, J = 6 and 16 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>18</sub>H<sub>30</sub>O<sub>3</sub> calcd 294.2195, found 294.2203.

6-Cyclopentyl-2,5-dihydroxy-2-methyl-3(*E*),6-heptadiene (22). A solution of the tetrahydropyranyl ether 21 (220 mg, 0.79 mmol) in EtOH (5 mL) was stirred with pyridinium *p*-toluenesulfonate (20 mg, 0.079 mol) at 55 °C in a preheated oil bath for 20 min. The solution was then immediately cooled to 0 °C and ethanol was removed at 0 °C by rotary evaporation. The residue was subjected to flash chromatography over silica gel with 30% EtOAc in hexane to afford the diol 22 (149 mg, 90%) as a semisolid: <sup>1</sup>H NMR  $\delta$  1.18–1.86 (16 H, including a 6 H singlet at  $\delta$  1.32 owing to 2 methyls), 2.37 (H, quin, J = 8 Hz), 4.59 (H, d, J = 6 Hz), 4.93 (H, s), 5.07 (H, s), 5.65 (H, dd, J = 6, 16 Hz); mass Spectrum m/z (M<sup>+</sup>) for C<sub>13</sub>H<sub>22</sub>O<sub>2</sub> calcd 210.1620; found 210.1609.

**5-**[(*tert*-**Butyldimethylsily**])oxy]-6-cyclopentyl-2-hydroxy-2-methyl-3-(*E*),6-heptadiene (23). To a solution of the diol 22 (100 mg, 0.475 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was added Et<sub>3</sub>N (50.8 mg, 0.502 mmol, 70  $\mu$ L) and 4-(dimethylamino)pyridine (10 mg, 0.082 mmol) at room temperature. The mixture was stirred at that temperature for 10 min. Then, *tert*-butyldimethylsilyl chloride (93 mg, 0.62 mmol) was added, and the solution was stirred at room temperature overnight. The solvent was removed, and the residue was flash chromatographed over silica gel with 10% EtOAc in hexane as eluant to give the silyl ether 23 (145 mg, 94%): <sup>1</sup>H NMR  $\delta$  0.03-0.10 (6 H), 0.86-0.91 (9 H), 1.25-1.82 (15 H, including one singlet at  $\delta$  1.32), 2.38 (H, quin, J = 8 Hz), 4.56 (H, d, J = 6 Hz), 4.85 (H, d, J = 0.92 Hz), 5.05 (H, t, J = 1.36 Hz), 5.54 (H, dd, J = 6 and 16 Hz), 5.80 (H, dd, J = 1.1 and 16 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>19</sub>H<sub>36</sub>O<sub>2</sub>Si caled 324.2485, found 324.2488.

5-[(tert-Butyldimethylsilyl)oxy]-6-cyclopentyl-3,4-epoxy-2-hydroxy-2-methyl-6-heptenes 24e and 24t. To a solution of the hydroxydiene 23 (130 mg, 0.477 mmol) in benzene (4 mL) was added vanadyl acetylacetonate (12 mg, 0.045 mmol) at room temperature. The mixture was stirred at 40 °C for 4 min while the solution became blueish green. Then, tert-butyl hydroperoxide (60.4 mg, 0.67 mmol) was added at room temperature, and the resulting mixture was stirred at 40 °C for 30 min. After the solvent was removed by rotary evaporation, the residue was chromatographed over silica gel with 10% EtOAc in hexane as eluent to afford a mixture of epoxides 24 (117 mg, 86%) in a ratio of 4:1 according to <sup>1</sup>H NMR analysis. The <sup>1</sup>H NMR spectrum of the mixture showed different signals for the SiOCH proton. The major isomer showed a peak at  $\delta$  4.10 (d, J = 4 Hz) and the minor isomer at  $\delta$  3.90 (d, J = 8 Hz): Mass spectrum m/z (M<sup>+</sup>) for C<sub>19</sub>H<sub>36</sub>O<sub>3</sub>Si calcd 340.2432, found 340.2398.

6-Cyclopentyl-2,5-dihydroxy-3,4-epoxy-2-methyl-6-heptenes 25e and 25t. A solution of the epoxy silyl ethers 24 (123 mg, 0.367 mmol) in THF (1.2 mL) was stirred with tetrabutylammonium fluoride (2.32 mL,

<sup>(22)</sup> Corey, E. J.; Ulrich, R.; Fitzpatrick, M. J. J. Am. Chem. Soc. 1976, 98, 222.

2.32 mmol, 1 M in THF) at room temperature overnight. Solvent was removed, and the residue was purified by flash chromatography over silica gel with 50% EtOAc in hexane to furnish the epoxy diols **25e** and **25t** (80 mg, 98%). The two isomers were separated by HPLC on a  $\mu$ -Partisil column with 5% 2-propanol-10% EtOAc-85% isooctane as eluting solvent to provide **25e** and **25t** in a 4:1 ratio, respectively (75 mg, 92% total yield). Major product **25e**: <sup>1</sup>H NMR  $\delta$  1.20 (3 H, s), 1.31 (3 H, s), 1.38-1.98 (10 H), 2.48 (H, quin, J = 8 Hz), 3.09 (H, d, J = 2.4 Hz), 3.21 (H, t, J = 2.7 Hz), 4.35 (H, brs), 5.02 (H, s), 5.12 (H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>13</sub>H<sub>22</sub>O<sub>3</sub> calcd 226.1569, found 226.1532.

Minor product 25t: <sup>1</sup>H NMR  $\delta$  1.21 (3 H, s), 1.30 (3 H, s), 1.40–2.16 (10 H), 2.50 (H, quin, J = 8 Hz), 2.98 (H, d, J = 2.34 Hz), 3.18 (H, dd, J = 2.38 and 4.8 Hz), 3.99 (H, bs), 5.01 (H, s), 5.14 (H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>13</sub>H<sub>22</sub>O<sub>3</sub> calcd 226.1569, found 226.1546.

Mesylate 26e. To a solution of epoxy diol 25e (15 mg, 0.066 mmol) in anhydrous methylene chloride (2.0 mL) was added 4-(N,N-dimethylamino)pyridine (48 mg, 0.392 mmol), and the resulting mixture was stirred 10-15 min at 0 °C under a blanket of nitrogen. Methanesulfonic anhydride (recrystallized from dry ether) (34 mg, 0.194 mmol) was then added, and the reaction mixture was stirred 4 h at 0 °C. The solvent was removed, and the residue was triturated with carbon tetrachloride and passed through a pipette containing a cotton plug. Evaporation of the solvent furnished a residue that was subjected to HPLC with 50% EtOAc in hexane as a solvent. Some diol 25e was recovered (1.5 mg). The mesylate 26e was obtained as an oil (12 mg, 66%): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.27 (1 H, s), 5.22 (1 H, d, J = 1.2 Hz), 5.01 (1 H, d, J = 4.4 Hz), 3.28 (1 H, dd, J = 4.4 and 2.2 Hz), 3.12 (1 H, d, J = 2.2 Hz), 3.02 (3 H, s), 2.53 (1 H, m), 2.08-1.46 (9 H), 1.33 (3 H, s), 1.27 (3 H, s).

Isopropyl Ether 27. Methanesulfonyl chloride (228 mg, 2.00 mmol) was added dropwise to a stirring mixture of erythro epoxy diol 25e (50 mg, 0.221 mmol) and pyridine (536 µL, 6.63 mmol) in methylene chloride (3 mL) at 0 °C. The mixture was stirred 6 h at 0 °C. Solvent was then removed by rotary evaporation, and the residue was triturated with CCl4 and the solution filtered. Solvent was removed by rotary evaporation, the residue was dissolved in 2-propanol (1 mL), and the resulting solution was boiled 6 h under reflux over  $K_2CO_3$  (50 mg). After being cooled to room temperature, the solution was filtered and the 2-propanol was removed by rotary evaporation. The residue was purified by HPLC with 20% ethyl acetate in hexanes to give the isopropyl ether 27 as an oil (40 mg, 67%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 5.08 (1 H, s), 5.02 (1 H, s), 3.65 (1 H, hept, J = 6.1 Hz), 3.57 (1 H, d, J = 6.7 Hz), 3.14 (1 H, dd, J = 6.1 Hz)2.3, 6.8 Hz), 2.84 (1 H, d, J = 2.4 Hz), 2.43 (1 H, quin, J = 7.2 Hz), 1.3-2.1 (9 H), 1.28 (3 H, s), 1.18 (3 H, s), 1.14 (6 H, d, J = 6.2 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>16</sub>H<sub>28</sub>O<sub>3</sub> calcd 268.2038, found 268.2040.

Reaction of Mesylate 26e with Tetramethylammonium Levulinate. To a solution of mesylate 26e (9 mg, 0.03 mmol) and tetramethylammonium levulinate (28 mg, 0.15 mmol) in freshly distilled chloroform (4 mL) was added sodium bicarbonate (2 mg), and the reaction mixture was heated 30 h under gentle reflux under nitrogen. TLC showed the disappearance of starting material. Solvent was removed in vacuo, and the residue was flash chromatographed over silica gel, eluting with 35% EtOAc in hexance to afford a mixture of levulinate esters 28 and 29. The mixture was separated by HPLC with 35% EtOAc in hexane as solvent. S<sub>N</sub>2' product 28 (2.9 mg, 31.5%): 'H NMR (CDCl<sub>3</sub>)  $\delta$  5.20 (1 H, d, J = 9 H), 4.56 (2 H, s), 3.74 (1 H, dd, J = 9 and 2.4 Hz), 3.07 (1 H, m), 2.88 (1 H, d, J = 2.4 Hz), 2.76 (2 H, t, J = 6.2 Hz), 2.60 (2 H, t, J = 6.2 Hz), 2.19 (3 H, s), 1.88–1.46 (9 H), 1.34 (3 H, s), 1.27 (3 H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>18</sub>H<sub>28</sub>O<sub>5</sub> calcd 324.1937, found 324.1934.

S<sub>N</sub>2 product, levulinate ester **29** (6.3 mg, 67%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 5.2-5.0 (3 H), 3.20 (1 H, dd, J = 6.4 and 2.4 Hz), 2.90 (1 H, d, J =2.4 Hz), 2.78-2.68 (2 H), 2.64-2.50 (2 H), 2.42 (1 H, quin, J = 8 Hz), 2.16 (3 H, s), 1.92 (9 H), 1.29 (3 H, s), 1.20 (3 H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>18</sub>H<sub>28</sub>O<sub>5</sub> calcd 324.1937, found 324.1925. The ratio of products **28:29** and overall yield varied with reaction solvent (See Results and Discussion; Table 1).

**Deprotection of Levulinate 29.** Sodium borohydride (5 mg, 0.132 mmol) was added to a stirring solution of **29** (9 mg, 0.028 mmol) in methylene chloride (1 mL) and water (120  $\mu$ L) at room temperature, and the mixture was stirred for an additional 30 min. Volatiles were removed in vacuo, and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and dried (MgSO<sub>4</sub>). Removal of solvent and flash chromatography of the resulting residue (50% ethyl acetate in hexanes) gave **25t** (5.4 mg, 85%).

Allylic Diepoxide trans, erythro-31 from the Erythro Epoxy Diol 25e. To a solution of the erythro epoxy diol 25e (20 mg, 0.11 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) at -78 °C was added pyridine (600  $\mu$ L, 587 mmol), and the solution was stirred at that temperature for 15 min. To this solution, methanesulfonyl chloride (94  $\mu$ L, 1.21 mmol) was added at -78 °C and the resulting mixture stirred 1 h at -78 °C and 1 h at 0 °C. The reaction mixture was concentrated to dryness in vacuo, and the residue was trit-

 
 Table II. Products from Reaction of Tosylate 32 with t-BuOK under Various Conditions

solvent	temp (°C)	time	cis,erythro-31: byproduct	yield (%)
benzene	10-25	1 h	12:88	85
toluene	-78	4 h	5:95	79
toluene	22	1 h	30:70	80
THF	22	1 h	71:29	82
DME	45	30 min	88:12	83
DME	85	5 min	85:15	73
DMF	35-40	30 min	100:0	82
t-BuOH	22	1 h	80:20	83
t-BuOH	55	30 min	80:20	78

urated with CCl<sub>4</sub> and filtered through a cotton plug. Solvent was removed by rotary evaporation to give the crude mesylate **26e**, which without further purification was dissolved in *t*-BuOH (500  $\mu$ L) and was stirred 1 h at room temperature with potassium *tert*-butoxide (13 mg, 0.12 mmol). The mixture was then filtered through a short dry-packed column of silica gel with 10% EtOAc in hexane. After removal of solvents, the crude product was purified further by HPLC with 5% EtOAc in hexane as eluent to afford the diepoxide *trans,erythro*-**31** (15.3 mg, 83%): <sup>1</sup>H NMR  $\delta$  1.33 (3 H, s), 1.38 (3 H, s), 1.43–1.98 (8 H), 2.44 (H, quin, J = 8 Hz), 2.59 (H, d, J = 6.5 Hz), 2.70 (H, dd, J = 2, 6.5 Hz), 3.36 (H, d, J = 2 Hz), 4.90 (H, t, J = 1.4 Hz), 5.0 (H, d, J = 1Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>13</sub>H<sub>20</sub>O<sub>2</sub> calcd 208.1463, found 208.1485.

Allylic Diepoxide cis, erythro-31 from Threo Epoxy Diol 25t. To a solution of the three epoxy diol 25t (15 mg, 0.083 mmol) in  $CH_2Cl_2$  (1.5 mL) at -78 °C was added pyridine (450 µL, 440 mmol), and the solution was stirred 15 min at that temperature. To this solution was added methanesulfonyl chloride (71 µL, 0.91 mmol) at -78 °C, and the mixture was stirred 1 h at -78 °C and 1 h at 0 °C. Solvents were removed in vacuo, and the residue was triturated with CCl4 and the solution filtered through a cotton plug. Solvent was removed by rotary evaporation, and the residue, crude mesylate 26t, without further purification was dissolved in t-BuOH and stirred 1 h at room temperature with potassium tertbutoxide (10 mg, 0.09 mmol). The mixture was filtered through a short dry-packed column of silica gel with 10% EtOAc in hexane as eluent. Solvent was removed by rotary evporation and the residue purified by HPLC with 5% EtOAc in hexane as eluent to afford the diepoxide cis,erythro-31 (11 mg, 80%): <sup>1</sup>H NMR & 1.31 (3 H, s), 1.43 (3 H, s), 1.4-2.0 (8 H), 2.4-2.6 (2 H, including a doublet at 2.50, J = 8 Hz), 2.90 (H, dd, J = 4.2, 8.1 Hz), 3.55 (H, d, J = 4 Hz); mass spectrum m/z $(M^+)$  for C<sub>13</sub>H<sub>20</sub>O<sub>2</sub> calcd 208.1463, found 208.1479. Tosylate 32. To a solution of 25e (30 mg, 0.13 mmol) in benzene (3

**Tosylate 32.** To a solution of **25e** (30 mg, 0.13 mmol) in benzene (3 mL) at room temperature were added triphenylphosphine (173 mg, 0.66 mmol) and zinc tosylate<sup>10</sup> (28.5 mg, 0.70 mmol). To the resulting suspension was added dropwise 104  $\mu$ L of diethyl azodicarboxylate (DEAD, 115 mg, 0.66 mmol). The resulting clear light yellow solution was stirred 2 h at room temperature. Solvent was removed and the residue was chromatographed rapidly over silica gel, eluting with 40% ethyl acetate in hexane to afford the inverted tosylate **32** (41 mg, 82%). The tosylate is unstable neat but can be stored at 20 °C in solution: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.79 (2 H, d, J = 7.8 Hz), 7.32 (2 H, d, J = 7.8 Hz), 5.12 (1 H, s), 5.05 (1 H, s), 4.74 (1 H, d, J = 6.4 Hz), 3.28 (1 H, dd, J = 6.4, 2.2 Hz), 2.91 (1 H, d, J = 2.2 Hz), 2.44 (3 H, s), 2.37 (1 H, m), 1.91–1.43 (9 H), 1.29 (3 H, s), 1.18 (3 H, s).

cis,erythro-31 from Tosylate 32. To a stirred solution of tosylate 32 (15 mg, 0.039 mmol) in dry 2-methyl-2-propanol (500  $\mu$ l) was added potassium tert-butoxide (8 mg, 0.070 mmol) under a blanket of nitrogen. The reaction mixture immediately turned yellow. Stirring was continued 1 h at room temperature. TLC showed the disappearance of starting material. The solvent was removed, and the residue was flash chromatographed over silica gel eluting with 5% ethyl acetate in hexane, affording an excellent separation of cis,erythro-31 (4.5 mg, 55%) identical by <sup>1</sup>H NMR with a sample prepared from three epoxy diol 25t (vide supra) and an unidentified, UV-active byproduct (1.2 mg): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  6.67 (d, 1 H, J = 13 Hz), 5.83 (1 H, d, J = 13 Hz), 4.84 (1 H, s), 4.80 (1 H, s), 4.69 (1 H, s), 2.58 (1 H, m), 1.92-1.49 (8 H), 1.41 (3 H, s), 1.32 (3 H, s). The influence of reaction conditions on the yield of the desired product, cis,erythro-31, was examined (Table II).

3-Hydroxy-4-isopropyl-1-(trimethylsilyl)pent-4-en-1-yne (33). To a solution of (trimethylsilyl)acetylene (2 g, 0.02 mol) in THF (10 mL) at -78 °C was added dropwise a solution of *n*-BuLi in hexane (12 mL, 0.024 mol, 2.0 M), and the resulting mixture was allowed to stir 1 h at -78 °C and 30 min at 0 °C. Then, the mixture was again cooled to -78 °C, and a solution of 2-isopropylacrolein (2 g, 0.02 mol) in THF (3 mL) was added dropwise, during which time the solution turned light blue. It was

stirred 1 h at the same temperature. Then, saturated aqueous NH<sub>4</sub>Cl was added, the cooling bath was removed, and the resulting mixture was allowed to warm to room temperature and extracted with ether, and the ether extract was washed twice with brine and dried (MgSO<sub>4</sub>). Rotary evaporation of the solvent and flash chromatography of the residue over silica gel with 10% EtOAc in hexane gave alcohol **33** (3.7 g, 92%): <sup>1</sup>H NMR  $\delta$  0.15 (9 H, t, J = 3.5 Hz), 1.07 (3 H, s), 1.11 (3 H, s), 1.86 (1 H, d, J = 6.5 Hz), 2.50 (1 H, quin, J = 8 Hz), 4.84 (1 H, d, J = 6.5 Hz), 4.97 (1 H, s), 5.31 (1 H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>11</sub>H<sub>20</sub>OSi calcd 196.1275, found 196.1283.

3-[(tert-Butyldimethylsilyl)oxy]-4-isopropyl-1-(trimethylsilyl)pent-4en-1-yne (34). To a solution of the alcohol 33 (500 mg, 2.55 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) were added Et<sub>3</sub>N (261 mg, 2.58 mmol, 3.60  $\mu$ L) and DMAP (10 mg, 0.082 mmol) at room temperature, and the solution was stirred for 10 min. Then, tert-butyldimethylsilyl chloride (480 mg, 3.18 mmol) was added, and the resulting mixture was stirred at room temperature overnight. The solvent was removed by rotary evaporation, and the residue was chromatographed over silica gel with hexane as eluent to give the silyl ether 34 (730 mg, 92%) as a volatile liquid: <sup>1</sup>H NMR  $\delta$  0.12–0.16 (15 H, m), 0.90 (9 H, s), 1.05 (3 H, s), 1.09 (3 H, s), 2.52 (1 H, quin, J = 8 Hz), 4.84 (1 H, d, J = 1.0 Hz), 4.90 (1 H, d, J = 1.0Hz), 5.26 (1 H, t, J = 1.0 Hz). Anal. Calcd for C<sub>17</sub>H<sub>34</sub>OSi<sub>2</sub>: C, 63.73; H, 11.03. Found: C, 63.88; H, 10.97.

3-[(tert-Butyldimethylsilyl)oxy]-1-iodo-4-isopropyl-1-(trimethylsilyl)penta-1(E),4-diene (35). To a solution of the acetylene 34 (165 mg, 0.532 mmol) in Et<sub>2</sub>O (660  $\mu$ L) at room temperature was added a solution of triisobutylaluminum in heptane (792 µL, 0.532 mmol, 0.67 M), and the resulting mixture was stirred for 5 min. A solution of diisobutylaluminum hydride in heptane (251 µL of 2.34 M, 0.532 mmol) was added to the reaction mixture, and stirring was continued for 4 h. Then, a solution of iodine (338 mg, 1.33 mmol) in Et<sub>2</sub>O (500  $\mu$ L) was added dropwise to the mixture at -78 °C. The resulting mixture was stirred 1 h at -78 °C and 2 h at 0 °C. Then, crushed ice and 10% HCl were added. The product was extracted into ether, and the ether layer was washed with 10% aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, saturated NaHCO<sub>3</sub>, and finally with brine and dried (MgSO<sub>4</sub>). Removal of solvent afforded an oil that was purified by flash chromatography over silica gel with hexane as eluent to give the iodide 35 (222 mg, 95%): <sup>1</sup>H NMR  $\delta$  0.05 (6 H, d, J = 1.6 Hz), 0.28 (9 H, s), 0.88 (9 H, s), 1.03 (3 H, d, J = 7 Hz), 1.04 (3 H, d, J = 7 Hz), 2.26 (H, quin, J = 7 Hz), 6.67 (H, dd, J = 1.1 and 9 Hz), 4.90 (H, dd, J = 1.0 Hz), 5.06 (H, t, J = 1.0 Hz), 6.97 (H, d, J = 9 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>17</sub>H<sub>35</sub>OSi<sub>2</sub>I calcd 438.1274, found 438.1271.

3-[(tert-Butyldimethylsilyl)oxy]-6-hydroxy-2-isopropyl-6-methyl-5-(trimethylsilyl)hepta-1,4(Z)-diene (36). To a solution of the iodide 35 (212 mg, 0.484 mmol) in THF (2 mL) at -78 °C was added dropwise a solution of n-BuLi in hexane (388  $\mu$ L of 2 M, 0.771 mmol), and the resulting mixture was stirred 30 min at -78 °C. Then, acetone (190  $\mu$ L, 2.58 mmol) was added at -78 °C, and the mixture was stirred 1 h at that temperature. Methanol was added, the cooling bath was removed, and the reaction mixture was allowed to warm to room temperature. Solvent was removed, and the residue was chromatographed over silica gel with 10% EtOAc in hexane as the eluent to afford the alcohol 36 (120 mg, 68%): 'H NMR  $\delta$  0.05 (6 H, d, J = 5 Hz), 0.21-0.24 (9 H), 0.85-0.90 (9 H), 1.03 (6 H, t, J = 7 Hz), 1.36 (6 H, d, J = 3 Hz); 2.39 (H, quin, J = 7 Hz), 4.85-4.97 (3 H), 5.89 (1 H, d, J = 8.6 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>20</sub>H<sub>42</sub>O<sub>2</sub>Si<sub>2</sub> calcd 370.2723, found 370.2723.

3-[(tert-Butyldimethylsilyl)oxy]-4,5-epoxy-6-hydroxy-2-isopropyl-6methyl-5-(trimethylsilyl)-hept-1-enes 37t and 37e. To a solution of the hydroxy diene 36 (16 mg, 0.043 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (500  $\mu$ L) at room temperature was added Mo(CO)<sub>6</sub> (1.2 mg, 0.0043 mmol) followed by tert-butyl hydroperoxide (8.0 mg, 0.090 mmol, 9  $\mu$ L), and the mixture was boiled 3 h under reflux. Solvent was removed, the residue was filtered through silica gel, solvents were removed by rotary evaporation, and the products were isolated by HPLC with 5% EtOAc in hexane as eluent to give the major threo isomer 37t (10.5 mg, 62%) and the minor erythro isomer 37e (3.6 mg, 21%) in a 3:1 ratio. <sup>1</sup>H NMR of the major isomer 37t:  $\delta$  0.05–0.26 (15 H), 0.89 (9 H, s), 1.04 (3 H, d, J = 1.6 Hz), 1.07 (3 H, d, J = 1.6 Hz), 1.14 (3 H, s), 1.28 (3 H, s), 2.46 (H, quin, J = 8 Hz), 3.11 (H, d, J = 8 Hz), 4.09 (H, d, J = 8 Hz), 4.96 (1 H, s), 5.08 (1 H, s).

<sup>1</sup>H NMR of the minor isomer **37e**:  $\delta 0.01-0.06$  (6 H, s), 0.24 (9 H, s), 0.88 (9 H, s), 1.08 (6 H, t, J = 6.5 Hz), 1.16 (3 H, s), 1.29 (4 H, s) for CH<sub>3</sub> and bs for OH at base line), 2.46 (1 H, quin, J = 8 Hz), 3.07 (1 H, d, J = 8 Hz), 4.16 (1 H, d, J = 8 Hz), 4.97 (1 H, s), 5.05 (1 H, s).

threo-3,6-Dihydroxy-4,5-epoxy-2-isopropyl-6-methylhept-1-ene (38t). To a solution of the silyl epoxide 37t (8.5 mg, 0.022 mmol) in THF (400  $\mu$ L) at room temperature was added a solution of Bu<sub>4</sub>NF in THF (110  $\mu$ L of 1 M, 0.11 mmol), and the resulting mixture was stirred overnight.

Solvent was removed by rotary evaporation, and the residue was purified by filtration through silica gel, eluting with 25% EtOAc in hexane to afford the diol **38t** (4.0 mg, 91%): <sup>1</sup>H NMR  $\delta$  1.06 (3 H, d, J = 7 Hz), 1.08 (3 H, d, J = 7 Hz), 1.21 (3 H, bs), 1.30 (3 H, s), 1.74 (1 H, bs), 2.06 (1 H, bs), 2.33 (1 H, quin, J = 7 Hz), 2.98 (1 H, d, J = 2 Hz), 3.16 (1 H, dd, J = 2, 5 Hz), 3.98 (1 H, bs), 5.02 (1 H, s), 5.17 (1 H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>11</sub>H<sub>20</sub>O<sub>3</sub> calcd 200.2802, found 200.1411.

erythro-3,6-Dihydroxy-4,5-epoxy-2-isopropyl-6-methylhept-1-ene (38e). To a solution of the silyl epoxide 37e (5.8 mg, 0.015 mmol) in THF (200  $\mu$ L) at room temperature was added a solution of Bu<sub>4</sub>NF in THF (75  $\mu$ L, 0.075 mmol, 1 M), and the resulting mixture was stirred overnight. Solvent was removed by rotary evaporation, and the residue was purified by filtration through silica gel, eluting with 25% EtOAc in hexane to afford the diol 38e (2.7 mg, 90%): <sup>1</sup>H NMR  $\delta$  1.07 (3 H, d, J = 7 Hz), 1.08 (3 H, d, J = 7 Hz), 1.20 (3 H, s), 1.30 (3 H, s), 1.84 (H, bs), 2.07 (1 H, bs), 2.33 (1 H, quin, J = 7 Hz), 3.10 (1 H, d, J = 2.4 Hz), 3.18 (1 H, t, J = 2.4 Hz), 4.35 (1 H, bs), 5.01 (1 H, s), 5.13 (1 H, s); mass spectrum m/z (M<sup>+</sup>) for C<sub>11</sub>H<sub>20</sub>O<sub>3</sub> calcd 200.2802, found 200.1417.

Bis(silyl ether) 39. A solution of the triol 14 (68 mg, 0.21 mmol), 4-(N,N-dimethylamino)pyridine (2.6 mg, 0.021 mmol), tert-butyldimethylsilyl chloride ((TBDMS)Cl, 90 mg, 0.060 mmol) and triethylamine (118 µL, 0.85 mmol) in anhydrous methylene chloride (5 mL) was heated under gentle reflux, and the reaction was monitored by TLC. After 26 h, (TBDMS)Cl (32 mg, 0.21 mmol) and triethylamine (25  $\mu$ L, 0.21 mmol) were added, and the resulting mixture was boiled under reflux for an additional 22 h where upon TLC analysis showed no remaining starting material. All volatiles were removed by rotory evaporation, and the residue was purified by flash chromatography over silica gel (30 mm  $\times$  7 in). Elution with 15% EtOAc in hexane gave the bis(silyl ether) 39 (110 mg, 94.5%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 5.76-5.56 (2 H, m), 4.93 (1 H, s), 4.80 (1 H, s), 4.37 (1 H, d, J = 5 Hz), 3.61 (1 H, d, J = 3.6 Hz), 3.37 (1 H, m), 2.16–1.38 (11 H), 1.32 (6 H, s), 0.92–0.82 (6 H), 0.87 (s, 9 H), 0.84 (s, 9 H), 0.06 (s, 3 H), 0.01 (s, 3 H), 0.01 (s, 3 H), 0.01 (s, 3 H).

Epoxy Bis(silyl ethers) 40. To a magnetically stirred solution of bis(silyl ether) 39 (90 mg, 0.16 mmol) in dry benzene (10 mL) was added VO(acac)<sub>2</sub> (4.3 mg, 0.016 mmol), and the resulting mixture was stirred 5 min at 40-45 °C under nitrogen. Then, tert-butyl hydroperoxide (50  $\mu$ L, 0.50 mmol) was added dropwise to the reaction mixture whereupon it turned straw yellow, which gradually disappeared during the next 15 min. The reaction mixture was stirred at 40-45 °C with progress monitored by TLC. After 4 h, the mixture still contained starting material. Therefore, 25 µL of tert-butyl hydroperoxide was added dropwise, and stirring was continued at the same temperature. After 2 h, TLC showed no starting material and the reaction was stopped. Rotary evaporation of solvents followed by flash chromatography of the residue over silica gel, using 10% EtOAc in hexane, afforded a diastereomeric mixture of epoxy bis(silyl ethers) 40 (80 mg, 86.3%). According to the vinyl <sup>1</sup>H NMR resonances, the erythro to threo ratio was 6:1: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.19 (1 H, s), 4.93 (3 H, s), 4.03 (1 H, d, J = 3.4 Hz), 3.62 (1 H, d, J = 3.6 Hz), 3.25 (1 H, m), 3.07 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz), 2.94 (1 H, dd, J = 3.6, 2.4 Hz)) d, J = 2.4 Hz), 2.22–1.36 (10 H), 1.28 (3 H, s), 1.24 (3 H, s), 0.87 (3 H, d, J = 6.2 Hz), 0.87 (3 H, s), 0.85 (9 H, s), 0.84 (9 H, s), 0.06 (3 H, s), 0.05 (3 H, s), 0.02 (3 H, s), 0.01 (3 H, s).

**Epoxy Triols 41.** The epoxy bis(silyl ethers) **40** (97 mg, 0.17 mmol) were dissolved in tetrahydrofuran (5 mL), and tetra-*n*-butylammonium fluoride (1.0 mL, 1 M solution in THF) was added. The reaction mixture was stirred at room temperature for 5 days since one of the two silyl ethers was difficult to remove. After 5 days, solvent was removed by rotary evaporation and the residue was flash chromatographed over silica gel with 80% EtOAc in hexane as eluting solvent. Two fractions were separated: the first contained a mono(silyl ether) (18.3 mg, 23.6%), and the second contained a 5.8:1 diastereomeric mixture of epoxy triols **41e** and **41t**, respectively. This mixture was separated by HPLC with 14% isopropanol in hexanes as eluting solvent. Erythro major epoxy triol **41e** (31.8 mg, 54.9%): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.17 (1 H, s), 4.98 (1 H, s), 4.16 (1 H, d, J = 3 Hz), 3.71 (1 H, d, J = 4 Hz), 3.23 (1 H, m), 3.15 (1 H, 1.29 (3 H, s), 1.24 (3 H, s), 0.96 (3 H, s), 0.83 (3 H, d, J = 6.4 Hz).

Three minor epoxy triol 41t (5.5 mg, 9.5%): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.22 (s, 1 H), 4.96 (1 H, s), 3.89 (1 H, d, J = 3.8 Hz), 3.72 (1 H, d, J = 4 Hz), 3.37–3.26 (4 H), 3.23 (1 H, dd, J = 4, 2.4 Hz), 2.94 (1 H, d, J = 2.4 Hz), 2.32–1.91 (4 H), 1.82–1.35 (6 H), 1.28 (3 H, s), 1.21 (3 H, s), 1.01 (3 H, s), 0.82 (3 H, d, J = 6.4 Hz).

Mono(silyl ether): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.19 (1 H, s), 5.00 (1 H, s), 4.27 (1 H, d, J = 2.8 Hz), 3.66 (1 H, d, J = 3.6 Hz), 3.24 (1 H, m), 3.19 (1 H, dd, J = 5.2 and 2.8 Hz), 3.14 (1 H, d, J = 2.8 Hz), 2.26–1.56 (12 H), 1.33 (3 H, s), 1.27 (3 H, s), 0.91 (3 H, s), 0.87 (3 H, d, J = 6.2 Hz), 0.87 (9 H, s), 0.02 (3 H, s), 0.01 (3 H, s).

Rearranged Allylic Diepoxide 42. The epoxy triol 41e (20 mg, 0.059 mmol) and triphenylphosphine (124 mg, 0.47 mmol) were dissolved in anhydrous benzene (2 mL), and zinc tosylate (14.5 mg, 0.035 mmol) was added. To the resulting suspension was added diethyl azodicarboxylate (DEAD, 75 µL, 0.48 mmol) dropwise under nitrogen. After being stirred 10 min, the light yellow reaction mixture became turbid and some solid precipitated. TLC analysis of the reaction mixture showed the complete disappearance of starting triol. The solvent was rotary evaporated, and the residue was flash chromatographed over silica gel and eluted very rapidly with 40% ethyl acetate in hexanes to yield a tosylate (15 mg): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.80 (2 H, d, J = 8.4 Hz), 7.32 (2 H, d, J = 8.4 Hz), 5.49 (1 H, s), 5.12 (1 H, s), 4.72 (1 H, d, J = 6.4 Hz), 4.64 (2 H, s), 3.19 (1 H, dd, J = 6.4, 2 Hz), 3.00 (1 H, d, J = 2 Hz), 2.44 (3 H, Hz)s), 2.44 (1 H, m), 2.17 (1 H, d, J = 4.6 Hz), 2.04–1.94 (3 H), 1.72–1.58 (3 H), 1.49-1.24 (5 H), 1.29 (s, 3 H), 1.17 (s, 3 H), 0.97 (3 H, d, J =6.4 Hz). This tosylate clearly incorporates the rearranged tricyclic skeleton of 42 with the 7-methylidene hydrogens appearing as a singlet at  $\delta$  4.64.

This tosylate (15 mg) was dissolved in anhydrous dimethylformamide (500  $\mu$ L) and treated with solid potassium tert-butoxide (10 mg) under a blanket of nitrogen. The reaction mixture immediately turned yellow. It was stirred 1.5 h at 30-35 °C. Solvent was then removed by azeotropic distillation with n-heptane on the rotary evaporator. Water (4 drops) was added to the residue. The resulting mixture was triturated with 40% ethyl acetate in hexanes, and the solution was filtered and dried by passing through a small bed of dry-packed MgSO4 and silica gel. After evaporation of the solvent, diepoxide 42 (8 mg, 46% overall from 41e) was obtained as an oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.24 (1 H, s), 5.12 (1 H, s), 4.67 (2 H, s), 3.50 (1 H, d, J = 4.2 Hz), 2.94 (1 H, dd, J = 7.8, 3.8 Hz), 2.56 (1 H, m), 2.52 (1 H, d, J = 7.8 Hz), 2.21 (1 H, d, J = 4.8 Hz), 2.08-1.98 (2 H), 1.86-1.58 (3 H), 1.46-1.22 (5 H), 1.43 (3 H, s), 1.30  $(3 \text{ H}, \text{s}), 0.97 (3 \text{ H}, \text{d}, J = 7 \text{ Hz}); {}^{13}\text{C} \text{ NMR} (\text{CDCl}_3) \delta 160.2, 142.2,$ 112.2, 98.7, 58.4, 58.1, 57.9, 56.0, 49.2, 47.5, 44.6, 43.2, 43.1, 38.0, 35.0, 31.8, 24.3, 19.2, 15.6; mass spectrum m/z (M<sup>+</sup>) for C<sub>20</sub>H<sub>28</sub>O<sub>2</sub> calcd 300.2089, found 300.2090.

**Mono(Silyl Ether) (+)-43.** A solution of diol (+)- $7^3$  (200 mg, 0.846 mmol), triethylamine (172 mg, 1.70 mmol), 4-(*N*,*N*-dimethylamino)-pyridine (10.4 mg, 0.845 mmol), and *tert*-butyldimethylsilyl chloride (160 mg, 1.06 mmol) in anhydrous methylene chloride (15 mL) was heated 2.5 h under gentle reflux. The reaction mixture was then cooled and the solvent and volatiles were removed by rotary evaporation. The residue was purified by chromatography over silica gel with ethyl acetate-hexanes (1:4) as eluant to furnish the monosilylated product (+)-43 (270 mg, 91%) as an oil:  $[\alpha]^{22}{}_{\rm D}$  + 23.7° (*c* 1.44, CHCl<sub>3</sub>): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  5.16 (1 H, t, *J* = 1.6 Hz), 4.82 (1 H, t, *J* = 1.6 Hz), 4.00 (1 H, d, *J* = 12 Hz), 3.93 (1 H, d, *J* = 12 Hz), 3.71 (1 H, d, *J* = 4 Hz), 2.94 (1 H, m), 2.27 (1 H, ddd, *J* = 13.5, 12, 4 Hz), 2.06–1.30 (9 H), 0.96 (3 H, s), 0.88 (9 H, s), 0.85 (3 H, d, *J* = 6.5 Hz), 0.03 (3 H, s), 0.02 (3 H, s).

p-Methoxybenzyl Ether (-)-44. To a stirred suspension of NaH (36 mg, 0.75 mmol, from NaH in oil prewashed twice with hexanes) in DME (10 mL) was added the alcohol (250 mg, 0.713 mmol) in DME (2 mL), and then the resulting mixture was stirred 30 min at room temperature. 4-Methoxybenzyl bromide (201 mg, 1.0 mmol) was added dropwise at room temperature. After the resulting solution was boiled 18 h under reflux, TLC analysis showed complete reaction. The mixture was cooled to room temperature. Then,  $H_2O$  (30  $\mu L$ ) was added very carefully, followed by ethyl acetate (25 mL). Anhydrous MgSO4 was added, the solution was filtered, and the solvent was rotary evaporated. The residue was then chromatographed over silica gel with ethyl acetate-hexanes (1:19) as eluant. The benzyl ether (-)-44 was obtained as a colorless oil (270 mg, 81%):  $[\alpha]^{22}_{D} - 37^{\circ}$  (c 1.4, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.22 (2 H, d, J = 8 Hz), 6.83 (2 H, d, J = 8 Hz), 5.17 (1 H, s), 4.83 (1 H, s)s), 4.47 (1 H, d, J = 12 Hz), 4.27 (1 H, d, J = 12 Hz), 3.99 (1 H, d, = 12 Hz), 3.92 (1 H, d, J = 12 Hz), 3.77 (3 H, s), 3.35 (1 H, d, J = 12 Hz), 3.77 (3 H, s), 3.35 (1 H, d, J = 12 Hz) 4 Hz), 2.84 (1 H, m), 2.16-1.14 (10 H), 0.98 (s, 3 H), 0.88 (s, 9 H), 0.84 (3 H, d, J = 6.4 Hz), 0.03 (3 H, s), 0.02 (3 H, s).

Allylic Alcohol (-)-45. To a solution of the silyl ether (-)-44 (270 mg, 0.57 mmol) in THF (10 mL) was added *n*-Bu<sub>4</sub>NF (1.7 mL of 1 M solution in THF), and the resulting mixture was stirred 16 h at room temperature. Then, solvent was rotary evaporated, and the residue was chromatographed over silica gel with ethyl acetate-hexanes (1:3) as eluant to afford the allylic alcohol (-)-45 (190 mg, 93%) as an oil:  $[\alpha]^{22}_D$  -35.5° (c 1.02, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.22 (2 H, d, J = 8.5 Hz), 6.83 (2 H, d, J = 8.5 Hz), 5.13 (1 H, s), 4.88 (1 H, s), 4.47 (1 H, d, J = 12 Hz), 4.28 (1 H, d, J = 12 Hz), 4.00 (1 H, d, J = 12 Hz), 3.94 (1 H, d, J = 12 Hz), 3.77 (3 H, s), 3.37 (1 H, d, J = 4 Hz), 2.96 (1 H, m), 2.39 (1 H, m), 2.21-1.21 (10 H), 0.99 (3 H, s), 0.84 (3 H, d, J = 6.4 Hz). Anal. Calcd for C<sub>23</sub>H<sub>32</sub>O<sub>3</sub>: C, 77.48; H, 9.04. Found: C, 77.47; H, 8.97.

Acrylaldehyde (+)-46. To a solution of the allylic alcohol (-)-45 (150 mg, 0.42 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (25 mL) was added active MnO<sub>2</sub> (225 mg, 2.59 mmol), and the reaction mixture was stirred 24 h at room temperature. It was then filtered through a bed of Celite, which was washed with ethyl acetate (5 × 10 mL). The combined organic solutions were evaporated in vacuo to afford the acrylaldehyde, which was purified by column chromatography over silica gel with an ethyl acetate-hexanes (1:9) eluant. The aldehyde (+)-46 was obtained (140 mg, 94%) as a colorless oil:  $[\alpha]^{22}_{D}$ +12.6° (c 0.79, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  9.40 (1 H, s), 7.15 (2 H, d, J = 7.8 Hz), 6.77 (2 H, d, J = 7.8 Hz), 6.12 (1 H, s), 5.96 (1 H, s), 4.40 (1 H, d, J = 12 Hz), 4.23 (1 H, d, J = 12 Hz), 3.70 (3 H, s), 3.34 (1 H, d, J = 4 Hz), 3.23 (1 H, m), 2.17-1.12 (10 H), 0.94 (3 H, s), 0.57 (3 H, d, J = 6 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>O<sub>3</sub> calcd 354.2195, found 354.2191.

Allylic Alcohols (-)-47a and (-)-47b. To a solution of the vinyl stannane 9 (550 mg, 1.12 mmol) in THF (4 mL) was added *n*-butyl-lithium (450  $\mu$ L, 1.12 mmol, 2.5 M in hexane) at -78 °C under nitrogen over 10 min. After the addition was complete, the mixture was stirred 2 h at -78 °C and 1 h at -10 °C. It was again cooled to -78 °C, and a solution of aldehyde (+)-46 (82 mg, 0.231 mmol) in THF (2 mL) was added dropwise. The reaction mixture then was allowed to warm to room temperature over 4 h, methanol (1.7 mL) was added, and then the volatiles were removed by rotary evaporation. The residue was flash chromatographed, eluting with ethyl acetate-hexanes (1:9) to afford an epimeric mixture of allylic alcohols (-)-47b and (-)-47a. This mixture was separated by HPLC with ethyl acetate-hexanes (1:9) as a eluant to yield (-)-47b (100 mg, 78%):  $[\alpha]^{22}_{D}$  -19.2° (c 7.61, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.25 (2 H, d, J = 8 Hz), 6.87 (2 H, d, J = 8 Hz), 5.81 (1 H, d, J = 16 Hz), 5.66 (1 H, dd, J = 16, 5.4 Hz), 5.11 (1 H, s), 4.95 (1 H, s), 4.51 (1 H, d, J = 12 Hz), 4.47 (1 H, d, J = 5.4 Hz), 4.31 (1 H, d, J = 12 Hz), 3.81 (3 H, s), 3.40 (1 H, d, J = 4 Hz), 3.22 (1 H, m), 2.12-1.38(10 H), 1.32(6 H, s), 1.03-0.87(15 H), 0.59(6 H, q, J = 8Hz).

Also isolated was (-)-47a (7.5 mg, 6%):  $[\alpha]^{22}_D - 28^{\circ}$  (c 0.4, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.26 (2 H, d, J = 8.8 Hz), 6.88 (2 H, d, J = 8.8Hz, 5.85 (1 H, d, J = 16 Hz), 5.56 (1 H, dd, J = 16 and 7 Hz), 5.35 (1 H, s), 4.98 (1 H, s), 4.52 (1 H, d, J = 12 Hz), 4.47 (1 H, d, J = 7Hz), 4.33 (1 H, d, J = 12 Hz), 3.84 (3 H, s), 3.41 (1 H, d, J = 4 Hz), 2.91 (1 H, m), 2.14–1.40 (10 H), 1.33 (6 H, s), 1.04–0.91 (15 H), 0.58 (6 H, q, J = 8 Hz).

**Diol** (-)-48b. To a solution of the silyl ether (-)-(47b) (90 mg, 0.16 mmol) in THF (2 mL) was added *n*-Bu<sub>4</sub>NF (450  $\mu$ L, 1.0 M solution in THF), and the resulting mixture was stirred 20 h at room temperature. The volatiles were removed in vacuo, and the residue was purified by flash chromatography over silica gel. Elution with 1:1 ethyl acetate-hexane afforded the diol (-)-48b (72 mg, 100%):  $[\alpha]^{22}_{D}$ -27.1° (*c* 1.24, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.23 (2 H, d, J = 8.8 Hz), 6.85 (2 H, d, J = 8.8 Hz), 5.89 (1 H, d, J = 16 Hz), 5.65 (1 H, dd, J = 16, 5.6 Hz), 5.09 (1 H, s), 4.94 (1 H, s), 4.48 (1 H, d, J = 12 Hz), 3.38 (1 H, d, J = 4 Hz), 3.19 (1 H, m), 2.10–1.38 (10 H), 1.32 (6 H, s), 1.01 (3 H, s), 0.86 (3 H, d, J = 6.5 Hz).

**Triethylsilyl Ether** (-)-49b. A solution of diol (-)-48b (72 mg, 0.16 mmol), DMAP (3.6 mg, 0.029 mmol), triethylamine (33 mg, 0.34 mmol), and triethylsilyl chloride (42 mg, 0.28 mmol) in methylene chloride (6 mL) was stirred at room temperature. After 4 h, TLC analysis showed complete silylation of the secondary hydroxyl group. Solvents and volatiles were removed by rotary evaporation, and the residue was chromatographed over silica gel, eluting with a mixture of ethyl acetate and hexane (1:4) to afford the silyl ether (-)-49b (90 mg, 100%):  $[\alpha]^{22}_D - 18.1^\circ$  (c 1.5, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.26 (2 H, d, J = 8.8 Hz), 6.88 (2 H, d, J = 8.8 Hz), 5.79 (1 H, d, J = 16 Hz), 5.61 (1 H, dd, J = 16, 6.4 Hz), 5.11 (1 H, s), 4.92 (1 H, s), 4.61 (1 H, d, J = 12 Hz), 4.48 (1 H, d, J = 6.4 Hz), 4.31 (1 H, d, J = 12 Hz), 3.83 (3 H, s), 3.41 (1 H, d, J = 3 Hz), 3.20 (1 H, m), 2.16-1.40 (10 H), 1.36 (6 H, s), 1.03-0.90 (15 H), 0.63 (6 H, q, J = 8 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>34</sub>H<sub>54</sub>O<sub>4</sub>Si calcd 554.3793, found 554.3791.

Epoxy Alcohols erythro-50b and threo-50b. To a solution of the allylic alcohol (-)-49b (75 mg, 0.135 mmol) in benzene (7.5 mL) was added VO(acac)<sub>2</sub> (3.6 mg, 0.014 mmol), and the mixture was stirred 5 min at 40-45 °C. The temperature was then raised to 60 °C, and tert-butyl hydroperoxide (27  $\mu$ L, 0.28 mmol) was added during which time the mixture turned straw yellow, which disappeared in the next 15 min. After the solution was stirred 2 h at 60 °C, more tert-butyl hydroperoxide (13.5  $\mu$ L, 0.13 mmol) was added and the resulting mixture stirred for another 1 h. It was then cooled to room temperature. Volatiles were removed by rotary evaporation, and the residue was chromatographed over silica gel. Elution with ethyl acetate-hexane (1:9) furnished a diastereomeric mixture of threo and erythro epoxy alcohol silyl ethers (58.5 mg, 76%) in a ratio of 3.7:1 as judged from the ratio of vinyl H

resonances at  $\delta$  5.22 and 4.99 (major) versus  $\delta$  5.20 and 4.96 (minor).

To a solution of the above diastereomeric mixture of epoxy alcohol silyl ethers (58.5 mg, 0.102 mmol) in THF (6 mL) was added *n*-Bu<sub>4</sub>NF (450  $\mu$ L of 1.0 M in THF), and the resulting mixture was stirred at room temperature. After 16 h, TLC analysis indicated that the reaction was complete. Solvents were removed in vacuo, and the residue was purified by flash chromatography over silica gel. Elution with ethyl acetate-hexane (1:1) afforded a mixture of epoxy diols that was separated by HPLC with 2-propanol-hexane (1:9) as eluting solvent to give *erythro***50b** (33.6 mg, 72.1%):  $[\alpha]^{22}_D$ -34.2° (*c* 0.5, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.23 (2 H, d, *J* = 8.8 Hz), 6.85 (2 H, d, *J* = 8.8 Hz), 5.21 (1 H, s), 5.04 (1 H, s), 4.48 (1 H, d, *J* = 12 Hz), 4.30 (1 H, d, *J* = 8.6 Hz), 4.29 (1 H, d, *J* = 12 Hz), 3.80 (3 H, s), 3.39 (1 H, d, *J* = 4 Hz), 3.28-3.13 (3 H), 2.12-1.38 (10 H), 1.32 (3 H, s), 1.24 (3 H, s), 1.02 (3 H, s), 0.86 (3 H, d, *J* = 6.5 Hz); mass spectrum *m/z* (M<sup>+</sup>) for C<sub>28</sub>H<sub>40</sub>O<sub>5</sub> calcd 456.2877, found 456.2875.

Also isolated are *threo*-**50b** (9.3 mg, 20.0%): <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$ 7.23 (2 H, d, J = 8.6 Hz), 6.86 (2 H, d, J = 8.6 Hz), 5.27 (1 H, s), 5.02 (1 H, s), 4.50 (1 H, d, J = 11.6 Hz), 4.31 (1 H, d, J = 11.6 Hz), 3.91 (1 H, br s), 3.80 (3 H, s), 3.40 (1 H, d, J = 4 Hz), 3.33–3.18 (2 H), 2.96 (1 H, d, J = 2.4 Hz), 2.06–1.27 (10 H), 1.32 (3 H, s), 1.23 (3 H, s), 1.02 (3 H, s), 0.85 (3 H, d, J = 6.4 Hz).

Levulinate Ester 51. A solution of the epoxy diol erythro-50b (5.0 mg, 0.011 mmol) and 4-(N,N-dimethylamino)pyridine (2.8 mg, 0.023 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (500 µL) was stirred 10 min at 0 °C. Then, methanesulfonic anhydride (4.0 mg, 0.023 mmol, freshly crystallized from anhydrous ether and dried under vacuum) was added. The resulting mixture was then stirred 5 h at 0-5 °C. Solvents were removed by rotary evaporation, and the residue was triturated with CCl4 and the solution filtered. After rotary evaporation of the solvent, the resulting crude mesylate was dissolved in 1,2-dimethoxyethane (500  $\mu$ L), and tetramethylammonium levulinate (10 mg, 0.053 mmol) and sodium bicarbonate (1 mg) were added. The resulting mixture was stirred 7 h at 85-90 °C and then cooled to room temperature. The solvents were rotary evaporated, and the residue was chromatographed over silica gel, eluting with ethyl acetate-hexane (2:3) to furnish levulinate ester 51 (4.8 mg, 79%):  $[\alpha]^{22}$  $-8.8^{\circ}$  (c 0.3, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.21 (2 H, d, J = 8.6 Hz), 6.85 (2 H, d, J = 8.6 Hz), 5.26 (1 H, s), 5.05 (1 H, d, J = 7 Hz), 5.04(1 H, s), 4.47 (1 H, d, J = 11.4 Hz), 4.29 (1 H, d, J = 11.4 Hz), 3.80(3 H, s), 3.37 (1 H, d, J = 3.4 Hz), 3.16 (1 H, dd, J = 7, 2.2 Hz), 3.01(1 H, d, J = 2.2 Hz), 2.94 (1 H, m), 2.80-2.61 (4 H), 2.18 (3 H, s),2.16-1.34 (10 H), 1.28 (3 H, s), 1.16 (3 H, s), 1.02 (3 H, s), 0.85 (3 H, d. J = 6.2 Hz)

**Epoxy Diol threo-50a.** To a solution of the levulinate ester **51** (4.0 mg, 0.007 mmol) in dioxane (500  $\mu$ L) and water (25  $\mu$ L) was added sodium borohydride (1.0 mg, 0.026 mmol). The resulting mixture was stirred 3 h at room temperature. Solvents were then rotary evaporated, and the residue was chromatographed on a small column of silica gel. Elution with ethyl acetate-hexane (1:1) furnished the diol *threo-50a* (2.8 mg, 88%): [ $\alpha$ ]<sup>22</sup><sub>D</sub> - 8.9° (c 0.28, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.21 (2 H, d, J = 8.6 Hz), 6.85 (2 H, d, J = 8.6 Hz), 5.39 (1 H, s), 5.05 (3 H, s), 4.47 (1 H, d, J = 11.4 Hz), 4.29 (1 H, d, J = 11.4 Hz), 3.83 (1 H, d, J = 5.8, 2.4 Hz), 3.02 (1 H, d, J = 2.4 Hz), 2.94 (1 H, m), 2.20-1.34 (10 H), 1.29 (3 H, s), 1.17 (3 H, s), 1.02 (3 H, s), 0.85 (3 H, d, J = 6.2 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>28</sub>H<sub>40</sub>O<sub>5</sub> calcd 456.2877, found 456.2876.

Allylic Diepoxide (-)-52 from erythro-50b. To a solution of epoxy diol erythro-50b (10 mg, 0.022 mmol) in 2 mL of benzene were added triphenylphosphine (35 mg, 0.13 mmol) and zinc tosylate (6 mg, 0.0147 mmol). Diethyl azodicarboxylate (21  $\mu$ L, 0.33 mmol) was then added dropwise to the resulting mixture to give a pale yellow clear solution. After the solution was stirred 2 h at room temperature, TLC analysis of the reaction mixture showed the presence of starting material. Triphenylphosphine and DEAD (2 equiv of each) were added again to complete the reaction. After 1 h more, TLC analysis showed no remaining starting material. The solvent was rotary evaporated, and the residue was immediately filtered rapidly through a short silica gel column with 30% ethyl acetate in hexanes as eluant. Removal of solvents in vacuo afforded a tosylate intermediate.

The tosylate intermediate was immediately dissolved in anhydrous N,N-dimethylformamide (1 mL), and to the resulting solution was added potassium *tert*-butoxide (10 mg, 0.089 mmol) under nitrogen. The reaction mixture was stirred 1.5 h at 35 °C until TLC analysis showed no unreacted tosylate. The solvent was removed by repeated azeotropic distillation with *n*-heptane by rotary evaporation. The residue was then triturated with 10% ethyl acetate in hexanes and passed through a short silica gel column to yield the benzyl ether (-)-**52** (6.5 mg, 68% from epoxy diol *erythro*-**50b**):  $[\alpha]^{22}_{D}-29^{\circ}$  (c 0.1, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.22 (2 H, d, J = 8.6 Hz), 6.84 (2 H, d, J = 8.6 Hz), 5.13 (1 H, d, J

= 1.2 Hz), 5.09 (1 H, s), 4.48 (1 H, d, J = 11.6 Hz), 4.29 (1 H, d, J = 11.6 Hz), 3.77 (3 H, s), 3.49 (1 H, d, J = 4 Hz), 3.38 (1 H, d, J = 2.6 Hz), 3.06 (1 H, m), 2.90 (1 H, dd, J = 7.4 and 4 Hz), 2.54 (1 H, d, J = 7.4 Hz), 2.15–1.64 (10 H), 1.42 (3 H, s), 1.01 (3 H, s), 1.29 (3 H, s), 1.01 (3 H, s), 0.91 (3 H, d, J = 6.4 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>28</sub>H<sub>38</sub>O<sub>4</sub> calcd 438.2771, found 438.2757.

15,16,17-Trisepispatol (-)-53. The benzyl ether (-)-52 (11.5 mg, 0.026 mmol) from erythro-50b was dissolved in 1 mL of methylene chloride. Water (55  $\mu$ L) was added, and to the resulting stirred suspension at 0 °C was added 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ, 12 mg, 0.052 mmol). After 1 h, TLC showed complete disappearance of the starting material. The solution was filtered through a pipette containing a little MgSO<sub>4</sub>, which was washed repeatedly with methylene chloride. The filtrate was rotary evaporated and the brown residue was immediately subjected to HPLC with 24% ethyl acetate in hexanes as eluant to deliver 15,16,17-trisepispatol (-)-53 (7 mg, 84.5%) as a white crystalline solid: mp 69–71 °C;  $[\alpha]^{22}_{D}$  –10.0° (c 0.70, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  5.14 (1 H, d, J = 1.3 Hz), 5.09 (1 H, s), 3.74 (1 H, d, J = 4.5 Hz), 3.50 (1 H, d, J = 4.0 Hz), 3.09 (1 H, ddd, J = 13.2, 5.5, 5.5 Hz), 2.90 (1 H, dd, J = 7.5, 4.1 Hz), 2.52 (1 H, d, J= 7.5), 2.25 (1 H, ddd, J = 13.2, 13.2, 4.3 Hz), 2.15 (1 H, t, J = 5.0Hz), 2.08-2.02 (1 H, m), 1.97 (1 H, t, J = 7.4 Hz), 1.88-1.80 (2 H), 1.76-1.66 (2 H), 1.46-1.32 (6 H), 1.30 (3 H, s), 0.98 (3 H, s), 0.92 (3 H, d, J = 6.7 Hz); <sup>13</sup>C NMR (100.607 MHz, CDCl<sub>3</sub>, for APT spectra a (+) indicates 0 or 2 attached protons and a (-) indicates 1 or 3 attached protons) δ 141.8 (+), 112.9 (+), 80.13 (-), 58.92 (-), 58.75 (+), 57.32 (-), 55.55 (-), 47.18 (+), 44.27 (-), 43.62 (-), 43.44 (-), 38.03 (-), 36.89 (+), 36.63 (-), 35.26 (+), 27.78 (+), 24.23 (-), 19.34 (-), 14.16 (-), 12.96 (-).

Allylic Diepoxide (-)-52 from threo-50a. A solution of the diol threo-50a (2.0 mg, 0.0044 mmol) and 4-(N,N-dimethylamino)pyridine (2.0 mg, 0.016 mmol) in methylene chloride (500  $\mu$ L) was stirred 10 min at 0 °C, and then methanesulfonic anhydride (3.0 mg, 0.017 mmol, freshly crystallized from anhydrous ether and dried under vacuum) was added. The resulting mixture was stirred 10 h at 0-5 °C. The solvents were removed by rotary evaporation, and the residue was triturated with CCl4 and the solution filtered. After removal of CCl4 in vacuo, the residue was dissolved in 2-methyl-2-propanol (500  $\mu$ L), and a 1% solution of potassium tert-butoxide in 2-methyl-2-propanol (26  $\mu$ L) was added. The resulting mixture was stirred 1 h at room temperature. Solvents were rotary evaporated, and the residue was dissolved in ethyl acetatehexane (1:1) and passed through a small bed of silica gel. After rotary evaporation of the solvent, the residue was subjected to HPLC with ethyl acetate-hexanes (1:6) as eluant with a Partisil PXS 10/25 column to afford the allylic diepoxide (-)-52 (1.4 mg, 72.5%):  $[\alpha]^{22}_{D}$  -29° (c 0.1, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.22 (2 H, d, J = 8.8 Hz), 6.84 (2 H, d, J = 8.8 Hz), 5.13 (1 H, s), 5.09 (1 H, s), 4.48 (1 H, d, J = 11.6 Hz), 4.29 (1 H, d, J = 11.6 Hz), 3.77 (3 H, s), 3.49 (1 H, d, J = 4 Hz), 3.38 (1 H, d, J = 2.8 Hz), 3.03 (1 H, m), 2.91 (1 H, dd, J = 7.6 and 4 Hz),2.54 (1 H, d, J = 7.6 Hz), 2.15-1.46 (10 H), 1.29 (3 H, s), 1.23 (3 H, s), 1.01 (3 H, s), 0.91 (3 H, d, J = 6.6 Hz); mass spectrum m/z (M<sup>+</sup>) for C<sub>28</sub>H<sub>38</sub>O<sub>4</sub> calcd 438.2771, found 438.2757.

**15,16,17-Trisepispatol** (-)-**53.** To a solution of *p*-methoxybenzyl ether (-)-**52** (1.1 mg, 0.0025 mmol) from *threo*-**50a** in a mixture of methylene chloride (500  $\mu$ L) and water (28  $\mu$ L) was added DDQ (1 mg, 0.0044 mmol, 1.7 equiv) at 0 °C, and the resulting mixture was stirred 1 h at the same temperature. Solvents were then rotary evaporated, and the residue was passed through a small bed of silica gel with ethyl acetatehexanes (1:2). The eluate was concentrated, and the residue was purified by HPLC on a Partisil PXS 10/25 column with EtOAc-hexanes (1:3) as eluant to provide trisepispatol (-)-**53** (0.6 mg, 75%), which was identical by <sup>1</sup>H NMR spectroscopy with the sample prepared previously from *erythro*-**50b**.

Allylic Diepoxide (+)-54. To a solution of *threo*-50b (6.0 mg, 0.013 mmol) in anhydrous methylene chloride (1.5 mL) were added DMAP (3.0 mg) and dry pyridine (20  $\mu$ L). The resulting mixture was stirred 5 min at 0 °C under N<sub>2</sub>. Methanesulfonic anhydride (35 mg, 0.20 mmol, freshly crystallized from anhydrous ether and dried under vacuum) was then added, and stirring was continued. After 3 h, TLC analysis showed no starting material remaining. Solvents were rotary evaporated, the residue was triturated with carbon tetrachloride several times, and the resulting solution was passed through a small cotton plug in a pipette. The solvent was rotary evaporated, and the mesylate thus obtained was immediately used for the next step.

The mesylate was dissolved in anhydrous 2-methyl-2-propanol (1.5 mL) and potassium *tert*-butoxide (15 mg) was added. The reaction mixture was stirred 1 h at room temperature under a nitrogen atmosphere. The solvent was then rotary evaporated, the residue was triturated with 25% ethyl acetate in hexanes, and the resulting solution was passed through a short silica gel column. The residue obtained after

rotary evaporation of solvent was subjected to HPLC with 15% ethyl acetate in hexanes as eluting solvent. Data for spatol benzyl ether (+)-54 (3.6 mg, 62.2%) thus obtained:  $[\alpha]^{22}{}_D + 8^\circ$  (c 0.12, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.22 (2 H, d, J = 8.2 Hz), 6.83 (2 H, d, J = 8.2 Hz), 5.09 (1 H, s), 5.02 (1 H, s), 4.48 (1 H, d, J = 11.8 Hz), 4.29 (1 H, d, J = 11.8 Hz), 3.77 (3 H, s), 3.43 (1 H, d, J = 4.4 Hz), 3.38 (1 H, d, J = 3 Hz), 2.93 (1 H, m), 2.87 (1 H, dd, J = 7.8, 4.4 Hz), 2.52 (1 H, d, J = 7.8 Hz), 2.11–1.61 (10 H), 1.41 (3 H, s), 1.30 (3 H, s), 1.01 (3 H, s), 0.89 (3 H, d, J = 6.4 Hz).

(+)-Spatol (3). To a magnetically stirred solution of spatol benzyl ether (+)-54 (3.6 mg, 0.0081 mmol) in methylene chloride (1.5 mL) and water (84  $\mu$ L) was added DDQ (6.0 mg) at 0 °C, and stirring was continued at the same temperature. After 1 h, TLC analysis showed no unreacted starting material. Solvents were rotary evaporated, and the residue was passed through a short column of silica gel with 30% ethyl acetate in hexanes as eluant. The residue obtained after rotary evaporation was purified by HPLC with 24% ethyl acetate in hexanes as eluant to deliver (+)-3 (2.1 mg, 81%):  $[\alpha]^{22}_{D}$  +44.2° (c 0.66, CHCl<sub>3</sub>) (reported<sup>2</sup>  $[\alpha]_{D}$  +45.6° (c 1.56, CHCl<sub>3</sub>)): <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  5.13 (1 H, dd, J = 3.0, 1.5 Hz), 5.03 (1 H, s), 3.74 (1 H, d, J = 4.4 Hz), 3.44 (1 H, d, J = 3.8 Hz), 3.03 (1 H, ddd, J = 14.5, 5.5, 5.5 Hz), 2.87 (1 H, dd, J = 7.9, 4.3 Hz), 2.49 (1 H, d, J = 7.9 Hz), 2.28 (1 H, ddd, J = 13.2, 13.2, 4.3 Hz), 2.12–2.05 (2 H), 1.97 (1 H, t, J = 6.8 Hz), 1.89–1.80 (2 H), 1.78–1.65 (2 H), 1.47–1.18 (3 H), 1.41 (3 H, s), 1.29 (3 H, s), 0.99 (3 H, s), 0.91 (3 H, d, J = 6.7 Hz); <sup>13</sup>C NMR (100.607 MHz, CDCl<sub>3</sub>, for APT spectra a (+) indicates 0 or 2 attached protons and a (-) indicates 1 or 3 attached protons)  $\delta$  141.5 (+), 111.0 (+), 79.96 (-), 58.42 (+), 58.12 (+), 57.06 (-), 55.14 (-), 47.21 (+), 43.79 (-), 43.40 (-), 43.25 (-), 37.82 (-), 36.87 (+), 36.56 (-), 35.18 (+), 27.87 (+), 24.23 (-), 19.26 (-), 14.47 (-), 12.97 (-).

Acknowledgment. This research was assisted financially by

Grant CA31595 from the National Cancer Institute of the National Institutes of Health. We thank M. Jirousek, B. Banik, K. Murthy, and A. Valeriu for technical assistance and Professor Fenical for copies of NMR spectra and for an authentic sample of natural spatol.

**Registry No.** (±)-1, 89196-20-3; 2, 132199-63-4; 3, 76520-52-0; 5, 930-30-3; (±)-6, 89165-79-7; (±)-7, 89165-80-0; (+)-7, 127182-28-9; (±)-8, 89177-59-3; 9, 132101-35-0; (+)-11, 127182-29-0; (+)-12, 132101-36-1; (+)-13, 132199-64-5; (+)-14, 132199-65-6; 19, 132101-37-2; 19 alcohol, 26639-84-9; (±)-20, 132199-66-7; 21, 132101-38-3;  $(\pm)$ -22, 132101-39-4;  $(\pm)$ -23, 132101-40-7;  $(\pm)$ -24e, 132101-41-8; (±)-24t, 132199-69-0; (±)-25e, 132101-42-9; (±)-25t, 132199-70-3; (±)-26e, 132101-43-0; (±)-26t, 132199-71-4; (±)-27, 132101-44-1; (±)-28, 132101-45-2; (±)-29, 132101-46-3; (±)-trans, erythro-31,  $132101-47-4; (\pm)-cis, erythro-31, 132199-72-5; (\pm)-32, 132101-48-5;$ (±)-33, 132101-49-6; (±)-34, 132101-50-9; (±)-35, 132101-51-0; (±)-36, 132101-52-1; (±)-37e, 132101-53-2; (±)-37t, 132101-65-6; (±)-38e, 132101-54-3; (±)-38t, 132199-73-6; 39, 132101-55-4; 40t, 132199-75-8; 40e, 132101-56-5; 41t, 132199-74-7; 41e, 132101-57-6; 42, 132101-58-7; 42 (tosylate precursor), 132101-66-7; (+)-43, 132101-59-8; (-)-44, 132101-60-1; (-)-45, 132101-61-2; (+)-46, 127156-24-5; (-)-47a, 132199-67-8; (-)-47b, 127135-70-0; (-)-48a, 127135-76-6; (-)-49b, 127135-71-1; threo-50a, 132199-68-9; threo-50b, 127182-31-4; erythro-50b, 127135-72-2; threo-50b TES ether, 127182-32-5; erythro-50b TES ether, 127135-73-3; 51, 132101-62-3; (-)-52, 127135-75-5; (-)-53, 127304-32-9; (+)-54, 127182-33-6; HC=C(CH<sub>3</sub>)<sub>2</sub>OH, 115-19-5; HC=  $C(CH_3)_2OTES$ , 17963-41-6; (±)-HC= $C(CH_3)_2OTHP$ , 132101-64-5; HC==CSiMe<sub>3</sub>, 1066-54-2; (CH<sub>3</sub>)<sub>2</sub>CHC(CHO)==CH<sub>2</sub>, 4417-80-5; diethyl cyclopentylmalonate, 18928-91-1; monoethyl (±)-cyclopentylmalonate, 132101-63-4; ethyl 2-cyclopentylacrylate, 81143-91-1.

# Synthesis and Chemistry of Dynemicin A Models

## K. C. Nicolaou,\* A. L. Smith,<sup>†</sup> S. V. Wendeborn,<sup>‡</sup> and C.-K. Hwang

Contribution from the Department of Chemistry, Research Institute of Scripps Clinic, 10666 North Torrey Pines Road, La Jolla California 92037, and Department of Chemistry, University of California, San Diego, La Jolla, California 92093. Received September 25, 1990

Abstract: The synthesis of the model systems 10 and 22 of dynemicin A (2) containing the nitrogen, enediyne, and epoxide functionalities has been achieved. These models are shown to undergo acid-induced triggering to give the corresponding Bergman-cyclized products in the presence of suitable H atom donors. Removal of the N protecting group from 22 gave the unstable free amine 30, which was shown to cause double-stranded-DNA cleavage, presumably in a manner similar to that of dynemicin A (2) itself. Some interesting chemistry related to dicobalt complexes of the enediynes is also presented.

### Introduction

A number of years ago, a new series of highly active antibiotics, the esperamicins<sup>1</sup> and calicheamicins,<sup>2</sup> was isolated. These antibiotics, containing a unique 1,5-diyn-3-ene bridging ring, displayed extremely potent antitumor activity with  $IC_{50}$  values in the nanogram per milliliter range against a number of murine and human cell lines.<sup>3</sup> The autitumor activity of these compounds has been ascribed to DNA damage resulting from H atom abstraction from the sugar phosphate backbone by a benzenoid diradical. This benzenoid diradical is generated by Bergman cyclization<sup>4</sup> of the enediyne bridge upon triggering by conformational changes brought about by bioreductive cleavage of the trisulfide moiety and 1,4-addition of the resulting thiol. The potency of these molecules has spawned considerable interest in the synthetic community, with the synthesis of a number of model

<sup>•</sup>To whom correspondence should be addressed at the Research Institute of Scripps Clinic.

<sup>&</sup>lt;sup>†</sup>Recipient of a NATO (SERC, U.K.) Postdoctoral Fellowship, 1990–1992. <sup>‡</sup>Recipient of a Verband der Chemischen Industrie Doctoral Fellowship, 1989–1990.

<sup>(1) (</sup>a) Golik, J.; Clardy, J.; Dubay, G.; Groenewold, G.; Kawaguchi, H.; Konishi, M.; Krishnan, B.; Doyle, T. W. J. Am. Chem. Soc. **1987**, 109, 3461-3462. (b) Golik, J.; Dubay, G.; Groenewold, G.; Kawaguchi, H.; Konishi, M.; Krishnan, B.; Ohjuma, H.; Doyle, T. W. J. Am. Chem. Soc. **1987**, 109, 3462-3464.

<sup>(2) (</sup>a) Lee, M. D.; Dunne, T. S.; Siegel, M. M.; Chang, C. C.; Morton,
G. O.; Borders, D. B. J. Am. Chem. Soc. 1987, 109, 3464–3466. (b) Lee, M.
D.; Dunne, T. S.; Chang, C. C.; Ellestad, G. A.; Siegel, M. M.; Morton, G.
O.; McGahren, W. J.; Borders, D. B. J. Am. Chem. Soc. 1987, 109, 3466–3468.

<sup>(3)</sup> Long, B. H.; Golik, J.; Forenza, S.; Ward, B.; Rehfuss, R.; Dabrowiak, J. C.; Catino, J. J.; Musial, S. T.; Brookshire, K. W.; Doyle, T. W. Proc. Natl. Acad. Sci. U.S.A. 1989, 86, 2-6.

 <sup>(4) (</sup>a) Bergman, R. G. Acc. Chem. Res. 1973, 6, 25-31. Jones, R. R.;
 Bergman, R. G. J. Am. Chem. Soc. 1972, 94, 660-661. Lockhart, T. P.;
 Gomita, P. B.; Bergman, R. G. J. Am. Chem. Soc. 1981, 103, 4091-4096. (b)
 Darby, N.; Kim, C. V.; Salaun, J. A.; Shelton, K. W.; Takadar, S.; Masamune,
 S. J. Chem. Soc., Chem. Commun. 1971, 1516-1517. (c) Wong, H. N. C.;
 Sondheimer, F. Tetrahedron Lett. 1980, 21, 217-220.