Acta Crystallographica Section C Crystal Structure Communications

ISSN 0108-2701

Diphenylmercury, redetermined at 120 K: sheets built from a single C— $H \cdots \pi$ (arene) hydrogen bond

Christopher Glidewell,^a* John N. Low^b and James L. Wardell^c

^aSchool of Chemistry, University of St Andrews, Fife KY16 9ST, Scotland,
^bDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, and ^cInstituto de Química, Departamento de Química Inorgânica, Universidade Federal do Rio de Janeiro, 21945-970 Rio de Janeiro, RJ, Brazil

Correspondence e-mail: cg@st-andrews.ac.uk

Received 21 December 2004 Accepted 22 December 2004 Online 31 January 2005

At 120 K, the molecules of the title compound, $[Hg(C_6H_5)_2]$, lie across centres of inversion in space group $P2_1/n$ and are linked by a single $C-H\cdots\pi(arene)$ hydrogen bond into (101) sheets. The same supramolecular structure is found at 298 K.

Comment

The structure of diphenylmercury, (I), was first successfully analysed only in 1977 (Grdenić et al., 1977), despite numerous earlier attempts (Kitaigorodsky & Grdenić, 1948; Ziólkowska, 1962; Ziólkowska et al., 1964), which had been hampered by a combination of inadequate absorption corrections and the occurrence of twinning. Using diffraction data collected at ambient temperature, an R value of 0.023 was achieved with 725 unweighted data and with H atoms included in the calculations with C-H distances in the range 0.93-1.02 Å (Grdenić et al., 1977). The structure consisted of nearly planar centrosymmetric molecules; although a number of fairly short $C \cdots C$, $C \cdots H$ and $H \cdots H$ intermolecular contacts were recorded, the consequences of these were not analysed. Our reanalysis, using PLATON (Spek, 2003), of the published structure using atomic coordinates retrieved from the Cambridge Structural Database (Allen, 2002; refcode DIPHHG11) found no direction-specific intermolecular interactions.



We have now reinvestigated the structure of (I) using a larger data set collected at 120 K; in addition, we collected a data set at 298 K, and the same phase was found. Although, for the sake of convenience and the much lower β angle, we have chosen to refine the structure in space group $P2_1/n$,

rather than the alternative $P2_1/c$, as employed by Grdenić *et al.* (1977), we also solved the structure in $P2_1/c$. It was evident, from both the cell dimensions and the atom coordinates in $P2_1/c$ that the same phase was employed in this study as in the earlier one. The structure of (I) determined at 120 K has better precision than that reported at ambient temperatures. Thus, the s.u. values on the Hg–C and C–C distances (Table 1) are less than half of those reported previously (0.007 Å for Hg–C and 0.010–0.014 Å for C–C). Despite the larger data set used here, at 120 K, the conventional *R* value is significantly lower than the previous ambient-temperature value (0.023).

We find at 120 K, centrosymmetric molecules (Fig. 1) with an interplanar spacing of 0.222 (4) Å, consistent with the





The molecule of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level and atoms labelled with the suffix A are at the symmetry position (1 - x, 1 - y, 1 - z).



Figure 2

Part of the crystal structure of (I), showing a (101) sheet formed from the action of a C-H··· π (arene) hydrogen bond. Hg atoms marked with an asterisk (*), a hash (#) or an ampersand (&) are at the positions (1, 0, 0), $(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$ and (0, 0, 1), respectively. For the sake of clarity, H atoms not involved in the motif shown have been omitted.

structure previously reported at ambient temperature. However, these molecules are linked into sheets by a single nearly linear $C-H\cdots\pi(\text{arene})$ hydrogen bond (Table 2). With the reference molecule centred at $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, atoms C6 at (x, y, z)and (1 - x, 1 - y, 1 - z) act as hydrogen-bond donors to the aryl rings at $(\frac{3}{2} - x, -\frac{1}{2} + y, \frac{1}{2} - z)$ and $(-\frac{1}{2} + x, \frac{3}{2} - y, \frac{1}{2} + z)$, respectively, which form parts of the molecules centred at (1, 0, 0) and (0, 1, 1), respectively. Similarly, the aryl rings at (x, y, z) and (1 - x, 1 - y, 1 - z) accept hydrogen bonds from atoms C6 at $(\frac{3}{2} - x, \frac{1}{2} + y, \frac{1}{2} - z)$ and $(-\frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z)$, respectively, which form parts of the molecules centred at (1, 1, 0) and (0, 0, 1). In this way, the molecules are linked into (101) sheets (Fig. 2), although there are no direction-specific interactions between adjacent sheets.

We find precisely the same supramolecular aggregation at 298 K, and the question then arises why this aggregation is not apparent from the coordinates reported by Grdenić et al. (1977). The explanation appears to lie in the location of the H atoms in the earlier structure. The authors stated that 'the H atoms were included in the structure-factor calculations with the isotropic thermal parameters of the bonded C atoms, but the parameters were not refined' (Grdenić et al., 1977); however, at no point did the authors specify how the H atoms were actually located or what constraints were applied to their positions during the refinement. In fact, analysis of their Hatom coordinates shows that many of these atoms are significantly displaced from the plane of the aryl ring; in particular, the two calculated C-C-C-H torsion angles for the H atom bonded to atom C6 are 168 and -169° . In addition, the two exocyclic C-C-H angles at atom C6 are 116 and 123°, and it seems probable that the erroneous location of the H atom bonded to atom C6 has previously obscured the occurrence of the C-H··· π (arene) hydrogen bond.

Experimental

The title compound was isolated from the reaction between mercury(II) chloride and methyltriphenyltin(IV) (2:1 molar ratio) in ethanol. Crystals suitable for single-crystal X-ray diffraction were grown by slow evaporation of a solution in ethanol.

Crystal data

$[Hg(C_6H_5)_2]$	$D_x = 2.458 \text{ Mg m}^{-3}$
$M_r = 354.79$	Mo $K\alpha$ radiation
Monoclinic, $P2_1/n$	Cell parameters from 1098
a = 5.6222 (3) Å	reflections
b = 8.0912 (4) Å	$\theta = 3.2-27.5^{\circ}$
c = 10.5852 (5) Å	$\mu = 16.00 \text{ mm}^{-1}$
$\beta = 95.485(3)^{\circ}$	T = 120 (2) K
V = 479.32 (4) Å ³	Plate, colourless
Z = 2	$0.28\times0.12\times0.04$ mm
Data collection	
Nonius KappaCCD diffractometer	837 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\rm int} = 0.034$
Absorption correction: multi-scan	$\theta_{\rm max} = 27.5^{\circ}$
(SADABS; Sheldrick, 2003)	$h = -7 \rightarrow 7$
$T_{\min} = 0.088, T_{\max} = 0.527$	$k = -10 \rightarrow 10$
7117 measured reflections	$l = -13 \rightarrow 13$
1098 independent reflections	

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_a^2) + (0.0209P)^2$
$R[F^2 > 2\sigma(F^2)] = 0.017$	+ 0.2975P]
$wR(F^2) = 0.045$	where $P = (F_o^2 + 2F_c^2)/3$
S = 1.09	$(\Delta/\sigma)_{\rm max} < 0.001$
1098 reflections	$\Delta \rho_{\rm max} = 0.71 \ {\rm e} \ {\rm \AA}^{-3}$
52 parameters	$\Delta \rho_{\rm min} = -0.87 \ {\rm e} \ {\rm \AA}^{-3}$
H-atom parameters constrained	Extinction correction: SHELXL97
	Extinction coefficient: 0.0133 (8)

Table 1

Selected geometric parameters (Å).

Hg1-C1	2.088 (3)	C4-C5	1.383 (5)
C1-C2	1.393 (5)	C5-C6	1.398 (5)
C2-C3	1.384 (5)	C6-C1	1.400 (5)
C3-C4	1.402 (5)		

Table 2

Hydrogen-bond geometry (Å, °).

$D - H \cdot \cdot \cdot A$	D-H	$H \cdot \cdot \cdot A$	$D \cdot \cdot \cdot A$	$D-\mathrm{H}\cdots A$
$C6-H6\cdots Cg^{i}$	0.95	2.84	3.759 (4)	164

Symmetry code: (i) $-x + \frac{3}{2}$, $y - \frac{1}{2}$, $-z + \frac{1}{2}$. Note: Cg is the centroid of the C1–C6 ring.

All H atoms were located from difference maps and then treated as riding, with C-H = 0.95 Å and $U_{iso}(H) = 1.2U_{eq}(C)$.

Data collection, cell refinement and data reduction: *COLLECT* (Hooft, 1999) and *DENZO* (Otwinowski & Minor, 1997); structure solution: *OSCAIL* (McArdle, 2003) and *SHELXS97* (Sheldrick, 1997); structure refinement: *OSCAIL* and *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97* and *PRPKAPPA* (Ferguson, 1999).

X-ray data were collected at the EPSRC X-ray Crystallographic Service, University of Southampton, England; the authors thank the staff for all their help and advice. JLW thanks CNPq and FAPERJ for financial support.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK1803). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. (2002). Acta Cryst. B58, 380-388.
- Ferguson, G. (1999). PRPKAPPA. University of Guelph, Canada.
- Grdenić, D., Kamenar, B. & Nagl, A. (1977). Acta Cryst. B33, 587-589.
- Hooft, R. (1999). COLLECT. Nonius BV, Delft, The Netherlands.
- Kitaigorodsky, A. I. & Grdenić, D. (1948). Izv. Akad. Nauk SSSR Otd. Khim. Nauk, p. 262.
- McArdle, P. (2003). OSCAIL for Windows. Version 10. Crystallography Centre, Chemistry Department, NUI Galway, Ireland.
- Otwinowski, Z. & Minor, W. (1997). Methods in Enzymology, Vol. 276, Macromolecular Crystallography, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.
- Sheldrick, G. M. (2003). SADABS. Version 2.10. University of Göttingen, Germany.
- Spek, A. L. (2003). J. Appl. Cryst. 36, 7-13.
- Ziólkowska, B. (1962). Rocz. Chem. 36, 1341-1347.
- Ziólkowska, B., Myasnikova, R. M. & Kitaigorodsky, A. I. (1964). Zh. Strukt. Khim. 5, 737–742.