

Crystallographic Approach to the Origin of "Syn-Effect"¹⁾

Katsuhiko INOMATA,^{*} Takaki HIRATA, Yoshihiro SASADA, Takahiro ASADA,
Hitoshi SENDA,[†] and Hideki KINOSHITA

Department of Chemistry, Faculty of Science, Kanazawa University, Kanazawa 920

[†]Department of Chemistry, College of Liberal Arts, Kanazawa University, Kanazawa 920

It was revealed by X-ray crystallography that α -unsubstituted (E)-vinyl sulfones have syn-conformation which seems to be the cause of "syn-effect" found in the conversion of (E)-vinyl sulfones to the corresponding allyl sulfones with a base under mild conditions. Syn-conformation of a terminal olefin in solid state was also confirmed.

In the previous papers,^{2a,b)} we reported the regio- and stereoselective syntheses of (E)- and (Z)-vinyl sulfones and their conversion to the corresponding allyl sulfones under the mild basic conditions, and it was found that (E)-vinyl sulfone preferentially affords (Z)-allyl sulfone as a kinetically-controlled product, while (Z)-vinyl sulfone and α -substituted vinyl sulfone give (E)-allyl sulfones exclusively. The latter fact may be due to the steric congestion which precludes the possibility of a stabilizing syn-interaction between the α - and δ -positions.^{3d)}

The former experimental results were rationalized by the new concept "conformational acidity" (a sort of kinetic acidity),^{2a,b)} which essentially depends on a "syn-effect"³⁾ within the cases of the conversion discussed so far. The relative degree of the "syn-effect" was further determined by the stereochemical investigation on the conversion of various kinds of γ -substituted (E)-vinyl sulfones with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) to the corresponding allyl sulfones as follows: $\text{CH}_3\text{O}- \geq \text{AcO}- > \text{CH}_3 > -\text{CH}_2- \gg \text{t-Bu}- \text{ and } \text{Ph}-$.^{2c)}

Several explanations for the "syn-effect" have been proposed,^{3b)} namely (1) 6π -electron homoaromaticity, (2) σ -orbital interactions, (3) dipole-dipole interactions, (4) chelations, and (5) hydrogen bonding.^{3e)} Regarding these problems, we have performed X-ray crystallography for 2-ethyl-1-tosyl-1-butene (1), which afforded exclusively (Z)-2-ethyl-1-tosyl-2-butene by treatment with DBU, and found that "syn-effect" worked in 1 itself as shown in Fig. 1.^{2c)} Herein we wish to report the new results of the crystallographic studies on the other α -unsubstituted (E)-vinyl sulfones and a crystalline terminal olefin derivative.

Since methoxy group has been proven to be the most effective one for "syn-effect" as described above,^{2c)} the conversion of the similar vinyl sulfones, (E)-3-phenoxy- and (E)-3-ethoxy-1-tosyl-1-propenes (2a,b), to the corresponding allyl sulfone derivatives (3a,b) was examined by using DBU as a base. The time-course of the reaction shown in Table 1 revealed that 2a,b afford exclusively (Z)-3a,b, especially at higher selectivity at the initial step of the reaction. These




Fig. 1. Stereoscopic view of 2-ethyl-1-tosyl-1-butene (1).

results strongly suggested the possibility that 2a,b have the syn-conformation as well as 1. In order to confirm the structure of 2a, X-ray crystallography was performed (Fig. 2).⁴⁾ It was found that H1, C1, C2, C3, O3 and H5, C17, C18, C19, O6 exist on a plane, respectively, and that "syn-effect" works in the solid state of 2a. The syn-conformation seems to arise from the intramolecular hydrogen bonding between the rather acidic α -hydrogens (H1 and H5), neighboring to the electron withdrawing tosyl group, and oxygens (O3 and O6) using the electron pairs of their sp^2 orbitals (O3-H1 2.36 Å, O6-H5 2.39 Å) or dipole-dipole interaction between $C_{sp^2}+H$ and $C\rightarrow OR$. However, 6π -electron homoaromaticity should not be excluded in spite of the long distances of O3-C1 (2.72(2) Å) and O6-C17 (2.78(2) Å), because there are p-orbitals on oxygen atoms (O3 and O6) conjugating with a phenyl group, which appear to correspond to the pseudo-p-orbital of the methyl group useful to stabilize syn-conformation of 1 in which the intramolecular hydrogen bonding is impossible.

Though the comparison of the structure of 2b with 2a was therefore required, 2b was oil at

Table 1. Conversion of (E)-3-Phenoxy- and (E)-3-Ethoxy-1-tosyl-1-propenes (2a,b) to the Corresponding 1-Tosyl-2-propene Derivatives (3a,b) with DBU


DBU (2 equiv.)
in CH₃CN, 25 °C

R=PhO, 2a
R=EtO, 2b


R=PhO, 3a
R=EtO, 3b

<u>2</u>	Time	Products ratio ^{a)}			Isolated total yield of <u>3</u> /%
		(E)- <u>2</u>	(E)- <u>3</u>	(Z)- <u>3</u>	
<u>2a</u>	0 min	100	0	0	-
	30 min	0	3	97	quant.
	2 h	0	3	97	-
	12 h	0	4	96	-
	96 h	0	8	92	-
<u>2b</u>	0 min	100	0	0	-
	30 min	27	2	71	-
	1 h	6	4	90	-
	2 h	0	5	95	-
	3 h	0	5	95	97
	6 h	0	4	96	-
	24 h	0	4	96	-
	72 h	0	4	96	-

a) Determined by 400 MHz ^1H -NMR spectra.



However, the question why the "syn-effect" was observed for (E)-vinyl sulfones such as 1 having no phenoxy or alkoxy group on γ -position remains. Therefore, X-ray crystallography of the terminal olefin (5) having 2-naphthalenesulfonyl group as a crystallizing auxiliary was performed (Fig. 4).⁶⁾ The syn-conformation was surprisingly observed (C13-C16 2.94(1) Å, C13-H17 2.67(10) Å, C16-H12 2.87(5) Å, C16-H13 2.93(4) Å) again, contrary to the reported preferable structure of 1-butene, CH₃-skew (83% of rotamer population).⁷⁾ Although the structure of 5 is only of the solid state [its packing diagram is shown in Fig. 4(c)] till now, it is very suggestive regarding the syn-conformation of the olefinic compounds. Related works including the structure of 5 in solution are further in progress in our laboratory.






Fig. 4. (a) Stereoscopic view and (b) numbering of 6-(2-naphthalenesulfonyl)-1-hexene (**5**) and (c) stereoscopic view of its packing diagram. Selected bond distances (Å) and dihedral angles (ϕ°): C13-C14 1.509(8), C14-C15, 1.51(1), C15-C16 1.27(1), C16-H18 1.00(7); C12-C13-C14-C15 174.8(5), C13-C14-C15-C16 5(1).

References

- 1) It means here the effects to stabilize syn-conformations against the steric hindrance.
- 2) a) K. Inomata, S. Sasaoka, T. Kobayashi, Y. Tanaka, S. Igarashi, T. Ohtani, H. Kinoshita, and H. Kotake, *Bull. Chem. Soc. Jpn.*, **60**, 1767 (1987); b) T. Kobayashi, Y. Tanaka, T. Ohtani, H. Kinoshita, K. Inomata, and H. Kotake, *Chem. Lett.*, **1987**, 1209; c) K. Inomata, T. Hirata, H. Suhara, H. Kinoshita, H. Kotake, and H. Senda, *Chem. Lett.*, **1988**, 2009.
- 3) a) D. Cremer, *J. Am. Chem. Soc.*, **101**, 7199 (1979); b) K. N. Houk, R. W. Strozier, N. G. Rondan, R. R. Fraser, and N. Chuaqui-Offermanns, *ibid.*, **102**, 1426 (1980) and references cited therein; c) E. Block, R. E. Penn, A. A. Bazzi, and D. Cremer, *Tetrahedron Lett.*, **22**, 29 (1981); d) E. Block, M. Aslam, V. Eswarakrishnan, K. Gebreyes, J. Hutchinson, R. Iyer, J.-A. Laffitte, and A. Wall, *J. Am. Chem. Soc.*, **108**, 4568 (1986); e) J. R. Larson, N. D. Epiotis, and F. Bernardi, *J. Am. Chem. Soc.*, **100**, 5713 (1978) and references cited therein.
- 4) **2a**, Mp 70 °C (from i-PrOH). Crystal data: $C_{16}H_{16}O_3S$, FW=288.4, Z=8, monoclinic, space group $P2_1/n$, $a=16.630(4)$, $b=7.669(2)$, $c=24.046(7)$ Å, $\beta=92.88(2)^\circ$, $U=3063(2)$ Å³, $D_c=1.25$ g/cm³, $F(000)=1216$, $\mu(\text{Mo-K}\alpha)=2.15$ cm⁻¹. Intensities were measured on a Rigaku AFC-5R diffractometer using Mo-K α radiation within $2\theta=45^\circ$ and θ - 2θ scan method. Observed independent reflections of 982 with $I>3\sigma(I)$ were used in the structure analysis and refinement applying TEXSAN program system. The final R factor was 0.070.
- 5) **4**, Mp 107 °C (from i-PrOH). This compound could not be converted to the allylsulfone by treatment with DBU, probably due to the unstable ketene acetal structure of the product. Crystal data: $C_{12}H_{14}O_4S$, FW=254.3, Z=2, monoclinic, space group $P2_1$, $a=5.489(1)$, $b=7.824(2)$, $c=13.877(2)$ Å, $\beta=90.95(1)^\circ$, $U=595.8(2)$ Å³, $D_c=1.42$ g/cm³, $F(000)=268$, $\mu(\text{Mo-K}\alpha)=2.71$ cm⁻¹; Rigaku AFC-5R, $2\theta=55^\circ$, θ - 2θ scan method, number of observation=1077 ($I>3\sigma(I)$), TEXSAN program system, R=0.034.
- 6) **5**, Mp 70.5-71.0 °C (from EtOH/H₂O). Crystal data: $C_{16}H_{18}O_2S$, FW=274.4, Z=4, monoclinic, space group $P2_1/c$, $a=9.817(4)$, $b=5.770(6)$, $c=25.97(1)$ Å, $\beta=100.34(3)^\circ$, $U=1447(2)$ Å³, $D_c=1.26$ g/cm³, $F(000)=584$, $\mu(\text{Mo-K}\alpha)=2.09$ cm⁻¹; Rigaku AFC-5R, $2\theta=55^\circ$, θ - 2θ scan method, number of observation=1309 ($I>3\sigma(I)$), TEXSAN program system, R=0.049.
- 7) D. Van Hemelrijk, L. Van den Enden, H. J. Geise, H. L. Sellers, and L. Schaefer, *J. Am. Chem. Soc.*, **102**, 2189 (1980).

(Received August 13, 1990)