## Synthesis, Resolution, and Crystallographic Characterization of a C<sub>2</sub>-Symmetric Diphosphaferrocene

Shuang Qiao, Diego A. Hoic, and Gregory C. Fu\*

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Received September 29, 1997

Summary: Treatment of  $FeCl_2$  with potassium 2-phenyl-3,4-dimethylphospholide provides a new  $C_2$ -symmetric diphosphaferrocene,  $(\pm)$ -3,3,4,4-tetramethyl-2,2-diphenyl-1,1'-diphosphaferrocene (1). Separation by chiral HPLC furnishes both antipodes of complex 1, the first diphosphaferrocene to be prepared in enantiopure form. The absolute configuration of (-)-1 has been determined by X-ray crystallography.

We have initiated a program directed at the development of applications of planar-chiral heterocycles in asymmetric catalysis. Our early work focused on the chemistry of  $\pi$ -bound nitrogen heterocycles, which we established serve as effective chiral nucleophilic catalysts<sup>1</sup> and as useful chiral ligands.<sup>2</sup> More recently, we have begun to explore the corresponding chemistry of  $\pi$ -bound *phosphorus* heterocycles.<sup>3</sup> At the time that we started this investigation, there were no reports of resolutions of this family of planar-chiral complexes. However, very recently Ganter has described the preparation of an enantiopure phosphaferrocene derivative through the use of a chiral auxiliary.<sup>4</sup> In this note, we establish, in the context of obtaining enantiopure diphosphaferrocene complex 1, that chiral HPLC represents a viable alternate method for the resolution of planarchiral phosphorus heterocycles. We have determined the absolute configuration of (-)-1 by X-ray crystallography.



In light of Cowley's report that 1,1'-diphosphaferrocenes can serve as bidentate ligands for transition metals,<sup>5</sup> we decided to prepare an enantiopure  $C_2$ symmetric diphosphaferrocene.<sup>6</sup> Complex **1** was an appealing target since phospholide **3** is readily available by a two-step route developed by Mathey.<sup>7</sup> We have



**Figure 1.** HPLC trace for ( $\pm$ )-**1** on a DAICEL CHIRACEL OD column ( $\alpha = 1.70$ ).

found that treatment of **3** with  $FeCl_2$  provides the desired diphosphaferrocene **1** and its meso isomer in 73% yield (eq 1). The diastereomers can be separated



by HPLC on silica, and the enantiomers of **1** can then be resolved by HPLC on a chiral stationary phase (DAICEL CHIRALCEL OD column;  $\alpha = 1.70$ ; Figure 1).<sup>8</sup>

<sup>(1) (</sup>a) Ruble, J. C.; Latham, H. A.; Fu, G. C. *J. Am. Chem. Soc.* **1997**, *119*, 1492–1493. (b) Ruble, J. C.; Fu, G. C. *J. Org. Chem.* **1996**, *61*, 7230–7231.

<sup>(2)</sup> Dosa, P. I.; Ruble, J. C.; Fu, G. C. J. Org. Chem. 1997, 62, 444-445.

<sup>(3)</sup> Garrett, C. E.; Fu, G. C. J. Org. Chem. 1997, 62, 4534–4535.
(4) Ganter, C.; Brassat, L.; Ganter, B. Tetrahedron: Asymmetry 1997, 8, 2607–2611.

<sup>(5)</sup> Atwood, D. A.; Cowley, A. H.; Dennis, S. M. *Inorg. Chem.* **1993**, *32*, 1527–1528.

<sup>(6) (</sup>a) For the synthesis of 1, 1'-diphosphaferrocene, see: De Lauzon,
G.; Mathey, F.; Simalty, M. J. Organomet. Chem. 1978, 156, C33–C36. (b) For the synthesis of a racemic C<sub>2</sub>-symmetric 1,1'-diphosphaferrocene derivative, see: de Lauzon, G.; Deschamps, B.; Fischer, J.; Mathey, F.; Mitschler, A. J. Am. Chem. Soc. 1980, 102, 994–1000.
(7) Holand, S.; Jeanjean, M.; Mathey, F. Angew. Chem., Int. Ed.

<sup>(7)</sup> Holand, S.; Jeanjean, M.; Mathey, F. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 98–100. Phosphole **2** can be synthesized in one step from commercially available PhPCl<sub>2</sub> and 2,3-dimethyl-1,3-butadiene. See: Breque, A.; Mathey, F.; Savignac, P. *Synthesis* **1981**, 983–985.

<sup>(8)</sup> For precedent for the use of chromatography to resolve planarchiral transition metal complexes, see: (a) Falk, H.; Schlögl, K. *Tetrahedron* **1966**, *22*, 3047–3053. (b) Falk, H.; Schlögl, K.; Steyrer, W. *Monatsh. Chem.* **1966**, *97*, 1029–1044.



**Figure 2.** ORTEP illustration, with thermal ellipsoids drawn at the 35% probability level, of **1**.

Table 1.	Crystallogra	phic D	ata for	1
----------	--------------	--------	---------	---

empirical formula	C <sub>24</sub> H <sub>24</sub> FeP <sub>2</sub>		
formula weight	430.22		
cryst color, habit	red block		
cryst dimens (mm <sup>3)</sup>	0.15 imes 0.27 imes 0.45		
cryst syst	monoclinic		
space group	C2		
a (Å)	22.9303(8)		
b (Å)	11.3262(4)		
c (Å)	16.6256(8)		
$\beta$ (deg)	105.5510(10)		
$V(Å^3)$	4159.8(3)		
Z	8		
Pcalc	1.374 Mg/m <sup>3</sup>		
radiation, monochromator	Mo K $\alpha$ ( $\lambda = 0.710$ 69 Å), graphite		
temp (K)	183(2)		
μ(Μο Κα)	$0.885 \text{ mm}^{-1}$		
diffractometer	Siemens SMART/CCD		
	(3-circle, $\chi$ fixed at 54.78°		
scan type; limiting indices	$\omega$ ; $-19 \leq h \leq 25$ ,		
	$-11 \leq k \leq 12$ ,		
	$-17 \leq l \leq 18$		
$\theta$ range for collection	1.27° to 23.23°		
total no. of reflns	8372		
no. of unique reflns	5422 ( $R_{int} = 0.0342$ )		
corrections	Lorentz-polarization; absorption		
	(semiempirical from $\psi$ -scans);		
	max and min trans.,		
	0.6199 and 0.5557		
structure solution,	direct methods, full-matrix		
refinement	least-squares on $F^2$		
data/restraints/params	5415/1/488		
$R_1$ ; w $R_2$ (data with $I > 2\sigma I$ )	0.0396; 0.1020		
$R_1$ ; w $R_2$ (all data)	0.0432; 0.1143		
goodness of fit	1.057		
ext coeff.; largest peak, hole	0.0007(2); 0.348 e Å <sup>-3</sup> ,		
	$-0.559 \mathrm{~e~\AA^{-3}}$		
absolute structure	-0.01(2)		
(Flack) param			
programs used	Siemens software package:		
	SMART, SAINT, XPREP,		
	SHELXTL 5.0		

Crystals of (–)-1 grown from  $CH_2Cl_2$ /pentane proved to be suitable for an X-ray diffraction study, which established the absolute configuration of this enantiomer (Figure 2; Table 1). The structure resembles that of 3,3',4,4'-tetramethyl-1,1'-diphosphaferrocene<sup>9</sup> —the two phospholyl rings are nearly parallel (6° deviation from coplanarity) and adopt an eclipsed geometry relative to one another (Figure 3).<sup>10</sup> The Fe–centroid distance is 1.67 Å.

Thus, we have developed a straightforward method for the synthesis and resolution of a new  $C_2$ -symmetric



Figure 3. View of 1 from above.

diphosphaferrocene (1), the first diphosphaferrocene to be prepared in enantiopure form. HPLC on a chiral stationary phase provided an effective means for separating the enantiomers of 1. We anticipate that complex 1 will find application in asymmetric catalysis.

## **Experimental Section**

Synthesis of 3,3',4,4'-Tetramethyl-2,2'-diphenyl-1,1'diphosphaferrocene (1). In a closed Schlenk tube, a mixture of 3,4-dimethyl-1-phenylphosphole (1.88 g, 10.0 mmol) and t-BuOK (1.35 g, 12.0 mmol) in THF (10 mL) was heated (140 °C) with stirring under N<sub>2</sub> for 12 h.<sup>7</sup> The resulting yellow solution was transferred by cannula to a stirred slurry of FeCl<sub>2</sub> (Strem; 0.89 g, 7.0 mmol) in THF (20 mL) at room temperature, immediately forming a dark red mixture. The reaction was stirred at room temperature for 12 h, and then the solvent was removed in vacuo. The resulting dark brown residue was purified by column chromatography (20% benzene/hexane), which afforded 1.58 g (73%) of complex 1 and its meso isomer (0.95:1) as a red solid. The diastereomers were separated by HPLC (Alltech Econosphere 10  $\mu$  silica; 250 mm  $\times$  22 mm; CH<sub>2</sub>Cl<sub>2</sub>:hexane 8:92; 20 mL/min; the racemate eluted from 11.5 to 12.6 min).

(±)-1: <sup>1</sup>H NMR (300 MHz, CD<sub>2</sub>Cl<sub>2</sub>)  $\delta$  7.30 (m, 4H), 7.14 (m, 6H), 3.44 (m, 2H), 2.15 (s, 6H), 2.09 (s, 6H); <sup>13</sup>C NMR (75 MHz, CD<sub>2</sub>Cl<sub>2</sub>)  $\delta$  139.6 (apparent t,  $J_{C-P}$  = 8.6 Hz), 131.0 (apparent t,  $J_{C-P}$  = 4.8 Hz), 128.1, 126.5, 100.8, 94.5, 83.4 (d,  $J_{C-P}$  = 6.1 Hz), 82.6 (d,  $J_{C-P}$  = 6.2 Hz), 15.5, 13.9; <sup>31</sup>P NMR (CD<sub>2</sub>Cl<sub>2</sub>, 122 MHz): - 63.7; IR (KBr) 3448, 3025, 2922, 1595, 1492, 1441, 1374, 1028, 839, 749; HRMS calcd for C<sub>24</sub>H<sub>24</sub>P<sub>2</sub>Fe 430.0703, found 430.0704. Anal. Calcd for C<sub>24</sub>H<sub>24</sub>P<sub>2</sub>Fe: C, 67.00; H, 5.62. Found: C, 67.30; H: 5.89.

The enantiomers of complex **1** were separated by chiral HPLC (14 mg per injection; DAICEL CHIRALCEL OD; 25 cm  $\times$  1 cm; chloroform:hexane 14:86; 2.5 mL/min). Enantiomer (+)-**1** ([ $\alpha$ ]<sup>20</sup><sub>D</sub> = +441°, *c* = 0.10, THF) was collected from 5.8 to 6.8 min, and (-)-**1** was collected from 7.6 to 10.0 min.

Acknowledgment. Support has been provided by the Alfred P. Sloan Foundation, the American Cancer Society, the Camille and Henry Dreyfus Foundation, Eli Lilly, Firmenich, Glaxo Wellcome, the National Science Foundation (Young Investigator Award to G.C.F., with funding from Procter & Gamble, Merck, Bristol-Myers Squibb, DuPont, Pfizer, Rohm & Haas, Pharmacia & Upjohn, and Bayer), and the Research Corporation. Acknowledgment is made to the donors of the Petroleum Research Fund, administered by the ACS, for partial support of this research.

**Supporting Information Available:** Text giving the experimental details of the X-ray crystal structure determination and tables of crystal data, atomic coordinates, bond lengths and angles, anisotropic displacement parameters, and hydrogen coordinates (13 pages). Ordering information is given on any current masthead page.

## OM970845S

<sup>(9)</sup> de Lauzon, G.; Deschamps, B.; Fischer, J.; Mathey, F.; Mitschler, A. J. Am. Chem. Soc. **1980**, *102*, 994–1000.

<sup>(10)</sup> For the structural characterization of 2,2',5,5'-tetraphenyl-1,1'diphosphaferrocene, see: Hitchcock, P. B.; Lawless, G. A.; Marziano, I. *J. Organomet. Chem.* **1997**, *527*, 305–308.