

A NEW APPROACH TO THE SYNTHESIS OF OPTICALLY ACTIVE *PSEUDO*-SUGAR AND *PSEUDO*-NUCLEOSIDE —SYNTHESIS OF *PSEUDO*- α -D-ARABINOFURANOSE, (+)-CYCLARADINE, AND (+)-1-*PSEUDO*- β -D-ARABINOFURANOSYLURACIL FROM D-ARABINOSE —

Masayuki YOSHIKAWA, ^{*}, ^a Nobutoshi MURAKAMI, ^a Yasunao INOUE, ^a Shoko HATAKEYAMA, ^a and Isao KITAGAWA ^b

Kyoto Pharmaceutical University, ^a 5 Nakauchi-cho, Misasagi, Yamashina-ku, Kyoto 607, Japan and Faculty of Pharmaceutical Sciences, Osaka University, ^b 1-6, Yamada-oka, Suita, Osaka 565, Japan


An optically active *pseudo*-sugar, *pseudo*- α -D-arabinofuranose, was synthesized from D-arabinose in favorable overall yield by using a stereoselective formation of branched nitropyranose as the key step. Furthermore, two biologically active *pseudo*- β -D-arabinofuranosyl nucleosides, (+)-cyclaradine and (+)-1-*pseudo*- β -D-arabinofuranosyluracil, were synthesized via Michael-type reactions of nucleic acid bases and a nitrocyclopentene derivative which was prepared from a synthetic intermediate of *pseudo*-D-arabinofuranose.

KEYWORDS *pseudo*- α -D-arabinofuranose; (+)-cyclaradine; (+)-1-*pseudo*- β -D-arabinofuranosyluracil; *pseudo*-nucleoside; *pseudo*-sugar; D-arabinose

Recently, we have developed a method for transforming D-glucose into *pseudo*-sugars and successfully synthesized optically active *pseudo*-hexopyranoses,¹⁾ *pseudo*-pentofuranoses,²⁾ and *pseudo*-aminosugars.³⁾ Furthermore, we have synthesized *pseudo*-nucleosides such as (-)-9-*pseudo*- β -D-glucopyranosyladenine,⁴⁾ (-)-9-*pseudo*- β -L-idopyranosyladenine,⁴⁾ (+)-9-*pseudo*- β -L-xylofuranosyladenine,⁵⁾ (+)-cyclaradine,⁵⁾ and (-)-aristeromycin⁶⁾ from nitro-cyclohexenes and nitro-cyclopentenones which were readily prepared from the synthetic intermediates of those *pseudo*-sugars. As an extension of these studies on synthesis of *pseudo*-sugar and *pseudo*-nucleoside, we have developed a new versatile conversion method starting from D-arabinose leading to *pseudo*- α -D-arabinofuranose (10). In addition, two biologically active *pseudo*- β -D-arabinofuranosyl nucleosides, (+)-cyclaradine (15) and (+)-1-*pseudo*- β -D-arabinofuranosyluracil (18), have been synthesized by use of a Michael-type addition reaction of nucleic acid base and a nitro-cyclopentene derivative which was prepared from the synthetic intermediate of *pseudo*-D-arabinofuranose.

Benzylation of methyl 3,4-*O*-isopropylidene- β -D-arabinopyranoside (1)⁷⁾ quantitatively gave 2, colorless oil, $[\alpha]_D -109.1^\circ$ (CHCl₃), C₁₆H₂₂O₅,^{8a)} which was treated with 80% aq. AcOH followed by selective benzylation⁹⁾ of the product to furnish methyl 2,3-di-*O*-benzyl- β -D-arabinopyranoside (3), colorless oil, $[\alpha]_D -64.2^\circ$ (CHCl₃), C₂₀H₂₄O₅.^{8b)} Swern oxidation¹⁰⁾ of 3 gave an unstable ketone 4, colorless oil, $[\alpha]_D -78.2^\circ$ (CHCl₃), C₂₀H₂₂O₅.^{8c)} which was treated with CH₃NO₂ in DMF in the presence of KF and 18-crown-6 to provide 5, colorless oil, $[\alpha]_D -20.8^\circ$ (CHCl₃), C₂₁H₂₅NO₇.^{8d)} The nitromethane adduct 5 was acetylated and subsequently subjected to deacetoxyhydrogenation with NaBH₄ in EtOH to furnish 6, colorless oil, $[\alpha]_D -54.0^\circ$ (CHCl₃), C₂₁H₂₅NO₆.^{8e)} Acidic hydrolysis of 6 gave a branched nitropyranose 7^{8f)} which was subjected to an intramolecular condensation reaction with CsF in DMF (25°C, 3h) to furnish the desired cyclization product 8 (a mixture of 5 α -nitro and 5 β -nitro epimers).^{8g)}

Ethoxyethylation of 8 followed by denitrohydrogenation with n-Bu₃SnH in toluene in the presence of α , α' -azobis-*iso*-butyronitrile (AIBN) gave 9.^{8h)} Finally, removal of the benzyl and ethoxyethyl groups of 9 furnished *pseudo*- α -D-arabinofuranose (10).²⁾ The present conversion method for *pseudo*-D-arabinose seems to be significant due not only to the simplicity of the procedure but also to the improved overall yield (14% from D-arabinose) as compared with yields in most of the previous methods.^{2, 11)} Next, the cyclization product 8 was treated with Ac₂O in the presence of *p*-TsOH·H₂O to provide a nitro-cyclopentene 11, colorless oil, $[\alpha]_D -82.5^\circ$ (CHCl₃), C₂₂H₂₃NO₆.⁸ⁱ⁾ Treatment of 11 with N⁶-benzoyladenine in DMF in the presence of CsF (0°C, 1h) provided 12, white powder, $[\alpha]_D +65.7^\circ$ (CHCl₃), C₃₄H₃₂N₆O₇.^{8j)} which was subjected to denitrohydrogenation, as described above for the conversion from 8 to 9, to furnish 13, white powder, $[\alpha]_D +47.7^\circ$ (CHCl₃), C₃₄H₃₃N₅O₅.^{8k)} Removal of the acetyl and N-benzoyl groups of 13 yielded 14, white powder, $[\alpha]_D +60.2^\circ$ (CHCl₃), C₂₅H₂₇N₅O₃.^{8l)} from which the benzyl groups were finally removed to furnish (+)-cyclaradine (15).⁵⁾ On the other hand, treatment of 11 with a silylated uracil¹²⁾ in CsF-DMF (25°C, 1h) gave 16, white powder, $[\alpha]_D +91.3^\circ$ (CHCl₃), C₂₆H₂₇N₃O₈.^{8m)} Reductive elimination of the nitro group in 16 gave 17, white powder,



Bn : $\text{C}_6\text{H}_5\text{CH}_2-$, EE : ethoxyethyl, Bz : $\text{C}_6\text{H}_5\text{CO}-$, PPTS : pyridinium *p*-toluenesulfonate

a) BnCl / NaH / DMF, **b)** 1) 80% aq. AcOH (50°C) 2) Bu_2SnO / toluene (110°C) 3) BnBr / CsF / DMF, **c)** Swern oxid., **d)** CH_3NO_2 / KF / 18-crown-6 / DMF, **e)** 1) Ac_2O / *p*-TsOH $\cdot$ H_2O 2) NaBH_4 / EtOH, **f)** c-HCl / AcOH (30°C), **g)** CsF / DMF (25°C), **h)** 1) ethyl vinyl ether / PPTS / CH_2Cl_2 2) $n\text{-Bu}_3\text{SnH}$ / AIBN / toluene (110°C), **i)** 1) H_2 / Pd-black / AcOEt 2) PPTS / 80% aq. acetone (40°C), **j)** 1) Ac_2O / *p*-TsOH $\cdot$ H_2O 2) pyridine, **k)** CsF / DMF (0°C), **l)** $n\text{-Bu}_3\text{SnH}$ / AIBN / toluene (110°C), **m)** 10% NaOMe-MeOH, **n)** H_2 (6 atm) / Pd-black / 10% AcOH-EtOH, **o)** 1) 1% NaOMe-MeOH 2) H_2 / Pd-black / 10% AcOH-EtOH.

$[\alpha]_D^{+120.5^\circ}$ (CHCl_3), $\text{C}_{26}\text{H}_{28}\text{N}_2\text{O}_6$ (8n), which was subjected to successive deprotection to furnish the final product, 1-(+)-pseudo- β -D-arabinofuranosyluracil (**18**), white powder, $[\alpha]_D^{+112.3^\circ}$ (CH_3OH), $\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_5$.^{8o)}

REFERENCES AND NOTES

- 1) M. Yoshikawa, B. C. Cha, T. Nakae, and I. Kitagawa, *Chem. Pharm. Bull.*, **36**, 3714 (1988).
- 2) M. Yoshikawa, B. C. Cha, Y. Okaichi, and I. Kitagawa, *Chem. Pharm. Bull.*, **36**, 3718 (1988).
- 3) M. Yoshikawa, B. C. Cha, Y. Okaichi, Y. Takinami, Y. Yokokawa, and I. Kitagawa, *Chem. Pharm. Bull.*, **36**, 4236 (1988).
- 4) I. Kitagawa, B. C. Cha, T. Nakae, Y. Okaichi, Y. Takinami, and M. Yoshikawa, *Chem. Pharm. Bull.*, **37**, 542 (1989).
- 5) M. Yoshikawa, T. Nakae, B. C. Cha, Y. Yokokawa, and I. Kitagawa, *Chem. Pharm. Bull.*, **37**, 545 (1989).
- 6) a) M. Yoshikawa, Y. Okaichi, B. C. Cha, and I. Kitagawa, *Chem. Pharm. Bull.*, **37**, 2555 (1989); b) *Idem*, *Tetrahedron*, **46**, 7459 (1990).
- 7) S. D. Gero, D. Horton, A. M. Sepulchre, and D. Wander, *Tetrahedron*, **29**, 2963 (1973).
- 8) All new compounds were fully characterized by IR (neat), ^1H -NMR (270 MHz, CDCl_3) and MS spectra (unless specified otherwise), a) **2**: IR: 2930, 1460 cm^{-1} , ^1H -NMR: δ 3.52 (dd, $J=3$, 8Hz, 2-H), 3.91 (br s, 5-H₂), 4.21 (m, 4-H), 4.33 (dd, $J=6$, 8Hz, 3-H), 4.64 (d, $J=3$ Hz, 1-H), EI-MS (%): m/z 294 (M^+ , 0.4); b) **3**: IR: 3480, 2930 cm^{-1} , ^1H -NMR: δ 3.71 (m, 5-H₂), 3.87 (m, 2, 3-H), 3.98 (m, 4-H), 4.68 (d, $J=3$ Hz, 1-H), EI-MS (%): m/z 312 (M^+ -MeOH, 2); c) **4**: IR: 3030, 1730 cm^{-1} , ^1H -NMR: δ 3.80 (dd, $J=3$, 10Hz, 2-H), 3.90, 4.14 (ABq, $J=15$ Hz, 5-H₂), 4.44 (d, $J=10$ Hz, 3-H), 4.76 (d, $J=3$ Hz, 1-H), EI-MS (%): m/z 342 (M^+ , 0.1); d) **5**: IR: 3470, 1560, 1380 cm^{-1} , ^1H -NMR: δ 3.47 (dd, $J=3$, 8Hz, 2-H), 3.60, 3.70 (ABq, $J=12$ Hz, 5-H₂), 3.89 (d, $J=8$ Hz, 3-H), 4.60 (d, $J=3$ Hz, 1-H), 4.70 (br s, 6-H₂), NOEs: 1-H & 2-H, 3-H & 6-H₂, EI-MS (%): m/z 342 (M^+ -CH₃NO₂, 0.1); e) **6**: IR: 1560, 1380 cm^{-1} , ^1H -NMR: δ 2.91 (m, 4-H), 3.36 (dd, $J=4$, 10Hz, 2-H), 3.51 (dd, $J=3$, 12Hz), 3.84 (dd, $J=2$, 12Hz) (5-H₂), 4.01 (dd, $J=6$, 10Hz, 3-H), 4.52 (dd, $J=10$, 13Hz), 4.69 (dd, $J=5$, 13Hz) (6-H₂), 4.61 (d, $J=4$ Hz, 1-H), EI-MS (%): m/z 355 (M^+ -MeOH, 0.1); f) **7**: colorless oil, IR: 3400, 3030, 1560, 1380 cm^{-1} , ^1H -NMR: 2.93-2.98 (m, 4-H), 3.39-3.91 (m, 2,3-H, 5-H₂), EI-MS (%): m/z 282 (3); g) **8**: colorless oil, IR (KBr): 3400, 3030, 1560, 1380 cm^{-1} , ^1H -NMR: 2.70-3.11 (m, 4-H), 3.57-4.05 (m, 2,3-H, 6-H₂), 4.47-4.95 (m, 1,5-H), EI-MS (%): m/z 355 (M^+ -H₂O, 0.1); h) **9**: colorless oil, IR: 2980, 1130, 1060 cm^{-1} , ^1H -NMR: δ 1.78-2.40 (m, 4,5-H), 3.35-4.16 (m, 1,2,3-H, 6-H₂), EI-MS (%): m/z 399 (M^+ -CH₃CHOEt, 1); i) **11**: IR: 3030, 1740, 1530 cm^{-1} , ^1H -NMR: δ 3.38 (m, 4-H), 4.12 (m, 3-H), 4.38 (d, $J=5$ Hz, 6-H₂), 4.57 (m, 2-H), 6.95 (br s, 1-H), EI-MS (%): m/z 397 (M^+ , 0.1); j) **12**: IR (KBr): 1740, 1700, 1610, 1590, 1560, 1370, 1240 cm^{-1} , ^1H -NMR: δ 2.02 (COCH₃), 3.20 (m, 4'-H), 3.89 (dd, $J=1$, 3Hz, 3'-H), 4.15 (br d, 2'-H), 4.33 (d, $J=7$ Hz, 6'-H₂), 5.46 (dd, $J=8$, 10Hz, 5'-H), 5.93 (dd, $J=5$, 10Hz, 1'-H), 8.24 (s, 8-H), 8.77 (s, 2-H), 9.10 (br s, NH), NOEs: 1'-H & 2'-H, 1'-H & 8-H, 5'-H & 6'-H₂, 5'-H & 8-H, EI-MS (%): m/z 398 (0.2); k) **13**: IR (KBr): 1740, 1700, 1610, 1580 cm^{-1} , ^1H -NMR: δ 2.05 (COCH₃), 2.13 (m, 5'-H), 2.48 (m, 4', 5'-H), 3.85 (br s, 3'-H), 4.08 (d, $J=5$ Hz, 2'-H), 4.19 (d, $J=7$ Hz, 6'-H), 5.30 (m, 1'-H), 8.25 (s, 8-H), 8.77 (s, 2-H), 9.12 (br s, NH), NOEs: 1'-H & 2'-H, 1'-H & 8-H, EI-MS (%): m/z 591 (M^+ , 2); l) **14**: IR (KBr): 3300, 1650, 1600 cm^{-1} , ^1H -NMR: δ 2.20 (m, 5'-H), 2.40 (m, 4', 5'-H), 3.72 (dd, $J=6$, 11Hz), 3.79 (dd, $J=5$, 11Hz) (6'-H₂), 3.93 (br s, 3'-H), 4.08 (dd, $J=1$, 5Hz, 2'-H), 5.21 (m, 1'-H), 5.65 (br s, NH₂), 8.04 (s, 8-H), 8.33 (s, 2-H), EI-MS (%): m/z 445 (M^+ , 0.7); m) **16**: UV (CHCl_3): 263 nm (ϵ 14000), IR (KBr): 3060, 3030, 1740, 1690, 1560, 1370 cm^{-1} , ^1H -NMR: δ 2.99 (m, 4'-H), 3.69 (dd, $J=2$, 4Hz, 3'-H), 4.00 (br d, 2'-H), 4.18 (d, $J=8$ Hz, 6'-H₂), 5.01 (dd, $J=8$, 11Hz, 5'-H), 5.59 (dd, $J=5$, 11Hz, 1'-H), 5.63 (d, $J=8$ Hz, 5-H), 7.35 (d, $J=8$ Hz, 6-H), 8.36 (br s, NH), EI-MS (%): m/z 397 (4); n) **17**: UV (CHCl_3): 266nm (ϵ 8500), IR (KBr): 3460, 3220, 1740, 1690, 1250 cm^{-1} , ^1H -NMR: δ 2.02 (COCH₃), 1.84 (dd, $J=10$, 13Hz), 2.15 (dd, $J=7$, 12Hz) (5'-H₂), 2.39 (m, 4'-H), 3.70 (dd, $J=1$, 4Hz, 3'-H), 3.99 (br d, 2'-H), 4.14 (d, $J=7$ Hz, 6'-H₂), 5.10 (ddd, $J=5$, 7, 13Hz, 1'-H), 5.65 (dd, $J=2$, 8Hz, 5-H), 7.46 (d, $J=8$ Hz, 6-H), 8.53 (br s, NH), NOEs: 1'-H & 2'-H, 1'-H & 4'-H, 1'-H & 6-H, EI-MS (%): m/z 373 (M^+ -CH₂Ph, 9); o) **18**: UV (MeOH): 268 nm (ϵ 10000), IR (KBr): 3350, 1690 cm^{-1} , ^1H -NMR (CD_3OD): δ 1.88, 2.12 (both m, 5'-H₂), 1.99 (m, 4'-H), 3.63 (dd, $J=7$, 11Hz), 3.74 (dd, $J=5$, 11Hz) (6'-H₂), 3.77 (dd, $J=3$, 5Hz, 3'-H), 3.99 (dd, $J=3$, 6Hz, 2'-H), 4.96 (ddd, $J=6$, 7, 12Hz, 1'-H), 5.62 (d, $J=8$ Hz, 5-H), 7.72 (d, $J=8$ Hz, 6-H), EI-MS (%): m/z 242 (M^+ , 5).
- 9) N. Nagashima and M. Ohno, *Chem. Lett.*, **1987**, 141.
- 10) K. Omura and D. Swern, *Tetrahedron*, **34**, 1651 (1978).
- 11) K. Tadano, H. Kimura, M. Hoshino, S. Ogawa, and T. Suami, *Bull. Chem. Soc. Jpn.*, **60**, 3673 (1987).
- 12) C. Chavis, F. Dumont, R. H. Wightman, J. C. Ziegler, and J. L. Imbach, *J. Org. Chem.*, **47**, 202 (1982).

(Received January 18, 1993)