

A new class of anomeric spironucleosides


Thanasis Gimisis,^a Carlo Castellari^b and Chrysostomos Chatgililoglu^{*a}

^a I.Co.C.E.A., Consiglio Nazionale delle Ricerche, Via P. Gobetti 101, I-40129 Bologna, Italy

^b Dipartimento di Chimica 'G. Ciamician', Università di Bologna, Via Selmi 2, I-40126 Bologna, Italy

A 1,5-hydrogen migration of a conveniently situated alkoxy radical affords spironucleosides which possess an unusual orthoamide structure at the anomeric position; X-ray crystallography establishes the configuration of the C-1' position.

Anomeric spironucleosides are useful modifications of natural nucleosides in that they contain the base unit in a fixed conformation around the *N*-glycosidic bond.¹ In recent years, they have gained in importance with the discovery of hydantocidin² **1**, a natural spironucleoside with herbicidal and plant growth regulatory activities, and generally, with the notion that important pharmaceutical leads can be found among modified nucleoside analogues.



The first anomeric spironucleosides to be synthesized were derivatives of psiconucleosides.^{3,4} All recent developments on the synthesis of anomeric spironucleosides are based on free-radical chemistry.^{3b,5,6}


Modification of the anomeric position of nucleosides can also be attained *via* the intermediacy of C-1' radicals.⁷ We recently reported the first example of this kind of transformation for the preparation of compound **2**.⁸ We report herein the synthesis of a new class of anomeric spironucleosides *via* a [1,5]-radical translocation of alkoxy radicals to the anomeric position.

When the protected 6-hydroxymethylribouridine **5**, prepared from the corresponding protected uridine **4** by fine tuning of literature procedures,⁹ was treated under the standard Suárez conditions¹⁰ a major product was obtained in moderate (36%) yield after flash column chromatography (Scheme 1). The structure of the product could be assigned to compound **6** based on its ¹H NMR spectrum, in which 2'-H appeared as a doublet and the 7-Hs appeared as a well resolved AB quartet. This, coupled with the lack of a 1'-H or 7-OH signal, corroborated the assignment of an orthoamide function to the product nucleoside, although the configuration of the anomeric centre remained ambiguous. Deprotection of the silyl groups provided the water soluble compound **7**. A single crystal of this material grown from MeOH–H₂O with mp 172–173 °C was subjected to X-ray diffraction analysis in order to determine unambiguously the stereochemistry of the C-1' position.

The independent part of the cell contains two formula units. An ORTEP diagram of **7** (molecule 1) is shown in Fig. 1. All atoms in molecule 1 were well defined, while atom O(5A) in molecule 2 exhibited a split image, as a consequence of two-fold rotational disorder around the C(4A)–C(5A) axis. An occupation factor for the latter was refined for O(5A) and the accompanying hydrogens, the final value being 0.57(1). The two independent molecules have strictly comparable param-

eters. In both molecules the stereogenic centre C(1A) is in the *R* configuration. The oxazolo[3,4-*c*]pyrimidine moiety is flat [maximum deviation from the average plane is 0.137(3) and 0.171(3) Å for atom O(7) in molecules **1** and **2**, respectively]. The dihedral angle between the five membered ring and the fused pyrimidine moiety is 82.3(2) and 86.5(2)° for molecules **1** and **2**, respectively.

When the Suárez conditions,¹⁰ were applied to the 2'-deoxynucleoside analogue **8**, prepared from the corresponding protected 2-deoxyuridine,⁹ two spironucleoside products were isolated in 71% combined yield and in a 1.1 : 1 ratio after flash column chromatography (Scheme 1). The structures of the products were assigned to compounds **9** and **10** based on their ¹H NMR spectra, in which the two diastereotopic 2'-Hs appeared as well resolved doublets of doublets and the 7-Hs appeared as an AB quartet. The determined absolute configuration of the anomeric centre in the spiroribonucleoside **7** was linked with that of the 2'-deoxy analogues through a straightfor-



Scheme 1 Reagents and conditions: i, Bu^tMe₂SiCl (5.0 equiv.) imidazole (7.0 equiv.), DMF, room temp., overnight, 98%; ii, LDA, THF, –70 °C, 3 h; HCO₂Et, –60 °C, 2 h; iii, NaBH₄, MeOH, room temp., 30 min, 68% based on consumption of **4**; iv, PhI(OAc)₂, I₂, cyclohexane, hv, 28 °C, 5 h, 36% for **6**, 71% for **9** and **10**; v, Bu₄NF on SiO₂, THF, room temp., 2 h, 90%; vi, (Pr₂SiCl)₂O, pyridine, room temp., overnight, 61%; vii, PhOC(S)Cl, DMAP, CH₂Cl₂, 1 h, room temp., 84%; viii, (Me₃Si)₂SiH, AIBN, toluene, 80 °C, 6 h, 95%





Fig. 1 ORTEP drawing of one of the independent molecules of **7** (molecule 1) in the unit cell

ward, three-step chemical transformation, as outlined in Scheme 1. Standard regioselective protection of the 3'- and 5'-positions with the 1,1,3,3-tetraisopropylidisiloxane-1,3-diyl group to afford **11**, followed by transformation of the 2'-hydroxy group to give the thiocarbonate **12**,¹¹ and finally Barton–McCombie radical deoxygenation in the presence of $(\text{Me}_3\text{Si})_3\text{SiH}$ ¹² produced the chromatographically less polar stereoisomeric 2'-deoxyspironucleoside **9**. These transformations, apart from establishing unequivocally the configuration of the C-1' anomeric centre in **9** and **10**, also demonstrated the inherent stability of the orthoamide structure.

The mechanism for the formation of the spironucleoside involves generation, under the Suárez conditions, of the alkoxy radical intermediate **13** which undergoes a [1,5]-radical translocation reaction¹³ to yield the anomeric C-1' radical intermediate **14** [eqn. (1)], which in turn produces the observed



orthoamide **6** after oxidation and cyclization. It is worth pointing out that the steric hindrance induced by the bulky 2'-substituent in the ribo series is likely responsible for the

stereospecificity of the cyclization. The absence of the epimeric spironucleoside with *S* configuration in the reaction product can be attributed to the steric interaction between the carbonyl in the 2-position of the base and the bulky-2' substituent as can be envisaged by inspection of the crystal structure in Fig. 1.

The spironucleosides reported herein correspond to the first spiro[tetrahydrofuran-2,3'-(1,5,6,7-tetrahydro-3*H*-[1,3]-oxazolo[3,4-*c*]pyrimidine)] nucleosides, possessing a remarkably stable orthoamide modification of the C-1' anomeric position. Further work on the preparation and chemical transformations of this class of compounds is in progress.

We are grateful to the European Commission for a 'Marie Curie' post-doctoral fellowship to T. G.

Footnote and References

* E-mail: chrys@area.bo.cnr.it

- 1 M. H. el Kouni, F. N. M. Naguib, R. P. Panzica, B. A. Otter, S.-H. Chu, G. Gosselin, C. K. Chu, R. F. Schinazi, Y. F. Shealy, N. Goudgaon, A. A. Ozerov, T. Ueda and M. H. Iltzch, *Biochem. Pharmacol.*, 1996, **51**, 1687.
- 2 M. Nakajima, K. Itoi, Y. Tamamatsu, T. Kinoshita, T. Okasaki, K. Kawakubo, M. Shindou, T. Honma, M. Tohjigamori and T. Haneishi, *J. Antibiot.*, 1991, **44**, 293.
- 3 (a) S. G. Zavgorodny, *Tetrahedron Lett.*, 1981, **22**, 3003; (b) for a recent approach to 6,1'-anhydro-6-hydroxypsicouridine, see M. S. P. Sarma, S. Megati, R. S. Klein and B. A. Otter, *Nucleosides Nucleotides*, 1995, **14**, 393.
- 4 H. Hrebabesky and J. Farkas, *Collect. Czech. Chem. Commun.*, 1974, **39**, 1098; A. Grouiller and J. Chattopadhyaya, *Acta Chem. Scand., Ser. B*, 1984, **338**, 367.
- 5 Y. Yoshimura, B. A. Otter, T. Ueda and A. Matsuda, *Chem. Pharm. Bull.*, 1992, **40**, 1761.
- 6 A. Kittaka, N. Yamada, H. Tanaka, K. T. Nakamura and T. Miyasaka, *Nucleosides Nucleotides*, 1996, **15**, 1447.
- 7 C. Chatgililoglu and T. Gimisis, in *Free Radicals in Biology and Environment*, ed. F. Minisci, Kluwer, Dordrecht, 1997, pp. 281–292.
- 8 (a) T. Gimisis and C. Chatgililoglu, *J. Org. Chem.*, 1996, **61**, 1908; (b) A. Kittaka, H. Tanaka, N. Yamada and T. Miyasaka, *Tetrahedron Lett.*, 1996, **37**, 2801.
- 9 M. P. Groziak, A. Koohang, W. C. Stevens and P. D. Robinson, *J. Org. Chem.*, 1993, **58**, 4054.
- 10 P. de Armas, C. G. Francisco and E. Suárez, *J. Am. Chem. Soc.*, 1993, **115**, 8865; R. L. Dorta, A. Martin, J. A. Salazar, E. Surárez and T. Prangé, *Tetrahedron Lett.*, 1996, **37**, 6021.
- 11 M. J. Robins and J. S. Wilson, *J. Am. Chem. Soc.*, 1981, **103**, 932.
- 12 C. Chatgililoglu and C. Ferreri, *Res. Chem. Intermed.*, 1993, **19**, 755.
- 13 A. L. J. Beckwith and K. U. Ingold, in *Rearrangements in the Ground and Excited States Vol. 1*, ed. P. de Mayo, Academic, New York, 1980, pp. 161–310.

Received in Glasgow, UK, 5th August 1997; 7/05742C