

New C,P-Cyclometalated Compounds of Platinum(II) and Platinum(IV) Derived from Tri-*o*-tolylphosphine

J. Forniés,* A. Martín, R. Navarro, V. Sicilia, and P. Villarroja

Departamento de Química Inorgánica, Instituto de Ciencia de Materiales de Aragón,
Universidad de Zaragoza-CSIC, 50009 Zaragoza, Spain

Received November 3, 1995[®]


The reaction between [*trans*-PtCl₂(PhCN)₂] (PhCN = benzonitrile) and tri-*o*-tolylphosphine rendered [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)]₂ (**1**), a new C,P-cyclometalated compound of Pt(II). The treatment of **1** with AgClO₄ (1: 2) in NCMe gave the mononuclear compound [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(NCMe)₂]ClO₄ (**2**). Solutions probably containing the cation [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(THF)₂]⁺ (THF = tetrahydrofuran) reacted with sodium dimethyldithiocarbamate (NaS₂CNMe₂) and potassium ethyl xanthate (KS₂COEt) in 1:1 molar ratio yielding the mononuclear complexes [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CR)] (R: NMe₂ (**3**), OEt (**4**)). The molecular structure of **3** was determined by X-ray diffraction. The reactions of **3** and **4** with Cl₂, Br₂ and I₂ gave the mononuclear Pt(IV) complexes [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CR)X₂] (X: Cl, Br, I) (**5–10**). In each reaction a single product can be detected spectroscopically, indicating that the oxidative addition of halogens (X₂) to **3** and **4** proceeds stereoselectively. The molecular structure of compound **7**, determined by X-ray diffraction, revealed that the obtained isomer has the OC-6-42 structure with both halogen atoms in *cis* positions.

Introduction

Cyclometalation reactions with tertiary phosphines give organometallic compounds with a C-P-M structure. Two important factors which promote internal metalation in such complexes are (a) the presence of bulky substituents on the tertiary phosphine and (b) the possibility of forming five-membered rings on metalation.¹ P(*o*-tolyl)₃ is an adequate ligand for such processes since the value of the cone angle, crystallographically determined, is 198°² and the activation of one H of the CH₃ in the *o*-tolyl substituent leads to a five-membered ring.

Phosphines with one [P^tBu₂(*o*-tolyl)] or two [P^tBu(*o*-tolyl)₂] *o*-tolyl groups undergo cyclopalladation and cycloplatination involving the methyl fragment of one *o*-tolyl group.^{3–5} Although tri-*o*-tolylphosphine undergoes side chain metalation on Mn⁶ and Rh⁷ complexes, as far as we know, only two palladium derivatives, [Pd{CH₂C₆H₄P(*o*-tolyl)₂}(μ-X)]₂ (X = I,^{8a} CH₃COO^{8b}), and one platinum derivative, [Pt{CH₂C₆H₄P(*o*-tolyl)₂}(P(*o*-tolyl)₃)(Br)],⁹ with cyclometalated P(*o*-tolyl)₃ have been

Chart 1



reported, [Pd{CH₂C₆H₄P(*o*-tolyl)₂}(μ-I)]₂ being the initial product formed during arylation reactions of conjugated polyenes using palladium (II) acetate, tri-*o*-tolylphosphine, and triethylamine as catalyst. Recently Herrmann et al. have reported that these C,P-cyclometalated Pd(II) complexes are very active catalysts for the Heck or Suzuki reactions.^{8b,c}

As an extension of our interest in cyclometalated derivatives,¹⁰ we decided to explore the reactions of cycloplatination of P(*o*-tolyl)₃. In this paper we describe the synthesis of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)]₂ (**1**), which we have used for the preparation of other Pt(II) and Pt(IV) complexes containing the "Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}" fragment [Pt(C⁺P)] in the following; see Chart 1A].

Results and Discussion

(A) Synthesis and Characterization of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)]₂ (**1**) and [Pt{CH₂

(9) Chappell, S. D.; Cole-Hamilton, D. J. *J. Chem. Soc., Dalton Trans.* **1983**, 1051.

(10) Forniés, J.; Navarro, R.; Sicilia, V.; Tomás, M. *Inorg. Chem.* **1993**, 32, 3675 and references given therein.

[®] Abstract published in *Advance ACS Abstracts*, March 1, 1996.

(1) (a) Omae, I. *Organometallic Intramolecular-Coordination Compounds*; J. Organomet. Chem. Libr. 18; Elsevier: Oxford, U.K., 1986. (b) Omae, I. *Coord. Chem. Rev.* **1980**, 32, 235.

(2) Ferguson, G.; Roberts, P. J.; Aleya, E. C.; Khan, M. *Inorg. Chem.* **1978**, 17, 2965.

(3) (a) Cheney, A. J.; Shaw, B. L. *J. Chem. Soc., Dalton Trans.* **1972**, 860. (b) *Ibid.* **1972**, 754.

(4) Gainsford, G. J.; Mason, R. *J. Organomet. Chem.* **1974**, 80, 395.

(5) Cheney, A. J.; McDonald, W. S.; O'Flynn, K.; Shaw, B. L.; Turtle, B. L. *J. Chem. Soc., Chem. Commun.* **1973**, 128.

(6) McKinney, R. J.; Hoxmeir, R.; Kesz, H. D. *J. Am. Chem. Soc.* **1975**, 97, 3059.

(7) (a) Bennett, M. A.; Longstaff, P. A. *J. Am. Chem. Soc.* **1969**, 91, 6266. (b) Mason, R.; Towl, A. D. C. *J. Chem. Soc. A* **1970**, 1601.

(8) (a) Rheingold, A. L.; Fultz, W. C. *Organometallics* **1984**, 3, 1414.


(b) Herrmann, W. A.; Brossmer, C.; Öfele, K.; Reisinger, C. P.; Priemeier, T.; Beller, M.; Fischer, H. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 1844. (c) Beller, M.; Fischer, H.; Herrmann, W. A.; Öfele, K.; Brossmer, C. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 1844.

Table 1. $^{31}\text{P}\{^1\text{H}\}$ NMR^a Data for Complexes 2–10

complex	$\delta(\text{P})$ (ppm)	$1J_{\text{Pt-P}}$ (Hz)
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{NCMe})_2](\text{ClO}_4)^b$ (2)	17.19 (s)	4673.0
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{CNMe}_2)]^b$ (3)	24.70 (s)	3969.4
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{COEt})]^b$ (4)	23.73 (s)	4083.8
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{CNMe}_2)\text{Cl}_2]^b$ (5)	24.20 (s)	2609.5
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{CNMe}_2)\text{Br}_2]^b$ (6)	21.87 (s)	2607.9
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{CNMe}_2)\text{I}_2]^b$ (7)	19.60 (s)	2620.7
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{COEt})\text{Cl}_2]^c$ (8)	24.91 (s)	2691.6
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{COEt})\text{Br}_2]^c$ (9)	23.13 (s)	2692.5
$[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{COEt})\text{I}_2]^c$ (10)	19.60 (s)	2620.5


^a $\text{C}^{\wedge}\text{P} = \text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2$ -C,P; external reference 85% H_3PO_4 ; CDCl_3 , room temperature. ^b Recorded on a Varian XL-200. ^c Recorded on a Varian Unity 300.

$\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2$ -C,P $\{(\text{NCMe})_2\}\text{ClO}_4$ (2**).** Equimolar quantities of $[\text{trans-PtCl}_2(\text{PhCN})_2]$ and tri-*o*-tolylphosphine react, in refluxing 2-methoxyethanol, giving the intramolecular coordination complex $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\mu\text{-Cl})_2]$ (**1**) (Chart 1B) (eq 1).



Complex **1** is insoluble in the usual organic solvents. Its IR spectrum shows two absorptions due to $\nu_{\text{st}}(\text{Pt}-\text{Cl})$ at 249 and 255 cm^{-1} , consistent with a dinuclear formulation, $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\mu\text{-Cl})_2]$ (**1**). Absorptions assignable to the $\text{C}^{\wedge}\text{P}$ ligand ($\text{C}^{\wedge}\text{P}:\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}$) are also observed (see Experimental Section).

Halide abstraction with AgClO_4 (1:2 molar ratio) in a donor solvent (L), at room temperature, affords the cationic solvento complexes $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{L})_2]^+$ (eq 2) (Chart 1C), which contain two solvent




molecules, such as THF (tetrahydrofuran) or NCMe (acetonitrile) in *cis* positions. For $\text{L} = \text{NCMe}$, a solid stable at room temperature, $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{NCMe})_2]\text{ClO}_4$ (**2**) was obtained. When $\text{L} = \text{THF}$, no solid has been isolated, but solutions containing presumably the $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{THF})_2]^+$ species have been used as starting materials for the synthesis of new complexes.

The IR spectrum of **2** (see Experimental Section) shows the following: (a) two absorptions assignable to the ClO_4^- (T_d) anion,¹¹ (b) several absorptions assignable to the $\text{C}^{\wedge}\text{P}$ ligand, and (c) two absorptions due to $\nu_{\text{st}}(\text{C}-\text{N})$ (2288 (m), 2298 (m) cm^{-1}) of the acetonitrile groups, denoting their *cis* positions. The shift to higher energies of these frequencies with respect to that in the free ligand (2254 cm^{-1}) suggests that the acetonitrile groups are N-coordinated, acting as pure σ -donors.¹² In the ^1H NMR spectrum (Table 2) the two inequivalent NCMe ligands give two singlets at 2.22 and 2.55 ppm. The methylene group of the $\text{C}^{\wedge}\text{P}$ fragment gives one

singlet at 3.39 ppm with the corresponding ^{195}Pt satellites ($^2J_{\text{Pt-H}}$: 101.25 Hz) which confirms the existence of the Pt–C σ -bond. The equivalence of these two diastereotopic hydrogen atoms agrees with the existence of a mirror plane in the molecule. Accordingly, the methyl groups of the two *o*-tolylphosphine's substituents appear as only one singlet at 2.55 ppm.

(B) Neutral C,P-Cyclometalated Pt(II) Complexes: $[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{CR})]$ ($\text{C}^{\wedge}\text{P} = -\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2$; $\text{R} = \text{NMe}_2$ (3**), OEt (**4**)).** Solutions of $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{THF})_2]\text{ClO}_4$ in methanol or THF react with sodium dimethyldithiocarbamate ($\text{NaS}_2\text{CNMe}_2$) and potassium ethyl xanthate (KS_2COEt) in 1:1 molar ratio to give the mononuclear complexes $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{CNMe}_2)]$ (**3**) and $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{COEt})]$ (**4**), in high yields, as air stable solids (Chart 1D) (eq 3).



In spite of the similarity of both complexes, the isolation procedure is different in each case (see Experimental Section) due to the solubility of complex **4** in the most common solvents.

In the IR spectrum of **3**, the presence of only one absorption in the 950–1050 cm^{-1} region (975 cm^{-1}) due to $\nu(\text{CS})$ indicates the bidentate binding of the “ S_2CNMe_2 ” group.¹³ An absorption at 1543 cm^{-1} corresponds to the $\nu_{\text{st}}(\text{C}-\text{N})$, and suggests a pronounced degree of C–N double bond.¹³ In the IR spectrum of **4**, the “ S_2COEt ” group is characterized by the presence of absorptions at 1239, 1123, and 1031 cm^{-1} .¹³

Structure of $[\text{Pt}(\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\{(\text{S}_2\text{CNMe}_2)\}]$ (3**).** Figure 1 shows the molecular structure of **3** with the atom-numbering scheme. General crystallographic information is collected in Table 3. Final atomic positional parameters are listed in Table 4. Relevant bond distances and angles are listed in Table 5.

As expected, the platinum atom is located in a distorted square-planar environment formed by the two sulfur atoms of the dimethyldithiocarbamate group and the phosphorus and an alkyl carbon (C(10)) atom of the C,P-cyclometalated phosphine ligand.

The five-membered metallocycle is not planar. C(4) and C(10) are 0.176 and 0.252 Å apart from the best least-squares plane through Pt–P–C(4)–C(9)–C(10).¹⁴ The small bite angle of the chelate ligands makes the corresponding angles smaller than 90°; S(1)–Pt–S(2) = 74.1(1)° and P–Pt–C(10) = 84.2(2)°. These parameters and the bond distances related with the platinum environment are similar to those found for other complexes containing these types of chelating groups.^{3–5,8}

The Pt–S(1) bond length (2.392(2) Å) is slightly longer than the Pt–S(2) one (2.354(1) Å) because of the

(13) Coucouvanis, D. *Progress in Inorganic Chemistry*; John Wiley & Sons: New York, 1979; Vol. 26, p 424.

(14) Nardelli, M. *Comput. Chem.* **1983**, 7, 95.

(15) Chen, H.-W.; Fackler, J. P., Jr.; Masters, A. F.; Pan, W.-H. *Inorg. Chim. Acta* **1979**, 35, L-333.


(11) Hathaway, B. J.; Underhill, A. E. *J. Chem. Soc.* **1961**, 3091.

(12) (a) Storhoff, B. N.; Lewis, H. C., Jr. *Coord. Chem. Rev.* **1977**, 23, 1. (b) Ford, P. C.; Clarke, R. E. *Chem. Commun.* **1968**, 1109.

Table 2. ^1H NMR^a Data for Complexes 2–10

compd	Pt–CH ₂			C ₆ H ₄ Me δ (ppm)	C ₆ H ₄ δ (ppm)	NCMe δ (ppm)	S ₂ CNMe ₂ δ (ppm)	S ₂ COEt	
	δ (ppm)	$^2J_{\text{H-H}}$ (Hz)	$^2J_{\text{Pt-H}}$ (Hz)					δ (ppm)	$^3J_{\text{H-H}}$ (Hz)
2	3.39 (s)		101.25	2.55 (s)	6.9–7.3 (m)	2.22 (s) 2.55 (s)			
3	3.38 (s)		85.70	2.42 (s)	6.9–7.3 (m)		3.23 (s)		
	3.44 (s)		94.30	2.78 (s)			3.24 (s)		
4	3.54 (s)		90.00	2.40 (s)	6.8–7.6 (m)			4.64 (q)	7.14
	3.55 (s)		98.20	2.74 (s)				1.47 (t)	
5	4.28 (d)	12.63	65.40	2.02 (s)	7.0–7.6 (m)		2.95 (s)		
	4.69 (d)		80.00	2.31 (s)			3.20 (s)		
6	4.38 (d)	12.45	68.00	2.00 (s)	7.0–7.6 (m)		2.92 (s)		
	4.82 (d)		79.00	2.32 (s)			3.19 (s)		
7	4.60 (d)	12.64	70.90	1.96 (s)	7.0–7.6 (m)		2.90 (s)		
	4.95 (d)		73.60	2.33 (s)			3.20 (s)		
8	4.23 (d)	12.25	64.40	2.04 (s)	7.0–7.6 (m)			4.58 (q)	7.04
	4.69 (d)		80.04	2.30 (s)				1.45 (t)	
9	4.32 (d)	12.01	66.10	2.02 (s)	7.0–7.6 (m)			4.57 (q)	6.93
	4.82 (d)		78.81	2.32 (s)				1.45 (t)	
10	4.52 (d)	12.25	85.00	1.97 (s)	7.0–7.6 (m)			4.54 (q)	7.17
	4.95 (d)		72.50	2.35 (s)				1.44 (t)	

^aSolvent CDCl₃, room temperature, 300 MHz; s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet.

**Figure 1.** Molecular structure of [Pt(CH₂C₆H₄P(*o*-tolyl)₂-C,P)(S₂CNMe₂)] (**3**).

different *trans* influence of the P and C donor atoms of the C,P-cyclometalated group.¹⁵

The [S₂CNMe₂][−] group is approximately planar. Bond angles around the N and C(1) atoms average 120°, as expected considering an sp² hybridization of both atoms; however the S(1)–C(1)–S(2) angle is only 112.8(3)° due to the coordination to the platinum center. The N–C(1) distance (1.336(7) Å) is not far from the C=N double bond distance (1.265 Å);¹⁶ these structural features are common in chelating dithiocarbamate complexes.¹⁷

NMR Spectra of Complexes 3 and 4. The ^1H NMR spectra of **3** and **4** in CDCl₃ show a similar pattern due to the cyclometalated group "Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}" (see Table 2). At room temperature (20 °C) the following is observed: (a) the –CH₂– group appears as an AX system with the corresponding ¹⁹⁵Pt satellites, $^2J_{\text{Pt-H}}$ being different and the H–H coupling constant not being observed. (b) The CH₃ groups of the *o*-tolylphosphine's substituents appear as two singlets indicating their inequivalence. Variable-temperature ^1H NMR spectra reveal the coalescence of both methyl groups and both methylenic protons of the "[CH₂C₆H₄P(*o*-tolyl)₂-C,P]" ligand at 45 °C. At 50 °C both methyl

groups are equivalent (**3**, 2.62 (s) ppm; **4**, 2.58 (s) ppm) and both methylenic protons are equivalent as well (**3**, 3.4 (s) ppm, $^2J_{\text{Pt-H}}$ = 93.6 Hz; **4**, 3.54 (s) ppm, $^2J_{\text{Pt-H}}$ = 94.92 Hz). These data indicate that at room temperature the molecular plane does not act as a mirror plane, i.e. the hindered rotation of the P–C_{ipso} bonds. Analysis of the proton ortho methyl resonances using the approximation to Eyring's equation¹⁸

$$\Delta G^\ddagger = 19.14 T_c (9.97 + \log T_c / \Delta \nu)$$

leads to the ΔG^\ddagger at the coalescence temperature for the P–C_{ipso} bond rotation process in complexes **3** and **4** (**3**, $\Delta G^\ddagger_{318} = 63.53 \text{ kJ mol}^{-1}$, $\Delta \nu = 108.34 \text{ Hz}$; **4**, $\Delta G^\ddagger_{318} = 63.79 \text{ kJ mol}^{-1}$, $\Delta \nu = 98.15 \text{ Hz}$). Only a few cases in which restricted P–C_{phenyl} bond rotation is also evident have been reported,^{9,19} and in most of them this process is hindered at low temperature but not at room temperature as in our cases.

The ^1H NMR spectra of **3** and **4** in CDCl₃ also show the signals due to the chelate S-donor ligands (see Table 2). The S₂CNMe₂[−] ligand gives two singlets at 3.23 and 3.24 ppm (**3**) even at 50 °C, indicating a high free energy of activation (ΔG^\ddagger) associated with C–N bond rotation in accord with its pronounced double-bond character in the dithiocarbamate ligand. This behavior is usual in dithiocarbamate Pt(II) and Pd(II) complexes.^{17,20} The S₂COEt[−] ligand (**4**) gives its two characteristic signals: 4.64 (q), 1.50 (t); $^3J_{\text{H-H}} = 7.14 \text{ Hz}$.

(C) Neutral C,P-Cyclometalated Pt(IV) Complexes: [Pt(C[†]P)(S₂CR)X₂] (C[†]P = –CH₂C₆H₄P(*o*-tolyl)₂, R = NMe₂, OEt; X = Cl, Br, I). The reactions of **3** and **4** with X₂ (X = Cl, Br, I) (1:1 molar ratio) in dichloromethane at room temperature afford the air-stable complexes [Pt(C[†]P)(S₂CR)X₂] (R = NMe₂, X = Cl (**5**), X = Br (**6**), X = I (**7**); R = OEt, X = Cl (**8**), X = Br (**9**), X = I (**10**)) in high yields.

Oxidative addition reactions of halogens to square-planar d⁸ organometallic complexes often result in the

(18) Günther, H. *NMR Spectroscopy: An Introduction*; John Wiley and Sons: Chichester, U.K., 1980.

(19) Howell, J. A. S.; Palin, M. G.; McArdle, P.; Cunningham, D.; Goldschmidt, Z.; Gottlieb, H. E.; Hezroni-Langerman, D. *Organometallics* **1993**, *12*, 1694.

(20) Fackler, J. P., Jr.; Lin, I. J. B.; Andrews, J. *Inorg. Chem.* **1977**, *16*, 450.

(16) Pauling, L. *The Nature Of The Chemical Bond*, 2nd ed.; Cornell University Press: Ithaca, NY, 1960.

(17) Chan, L. T.; Chen, H.-W.; Fackler, J. P., Jr.; Masters, A. F.; Pan, W.-H. *Inorg. Chem.* **1982**, *21*, 4291.

Table 3. Summary of Crystallographic Data

	3	7
formula	C ₂₄ H ₂₆ NPptS ₂ ·CHCl ₃	C ₂₄ H ₂₆ I ₂ NPptS ₂ ·1/2CH ₂ Cl ₂
<i>M_r</i>	738.043	914.941
cryst system	monoclinic	monoclinic
space group	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ / <i>c</i>
systematic absences	0 <i>k</i> 0, <i>k</i> = 2 <i>n</i> + 1; <i>h</i> 0 <i>l</i> , <i>l</i> = 2 <i>n</i> + 1	0 <i>k</i> 0, <i>k</i> = 2 <i>n</i> + 1; <i>h</i> 0 <i>l</i> , <i>l</i> = 2 <i>n</i> + 1
<i>a</i> , Å	12.2980(10)	10.961(2)
<i>b</i> , Å	12.9748(10)	14.431(3)
<i>c</i> , Å	18.2293(18)	18.587(4)
β, deg	105.766(10)	91.39(3)
<i>V</i> , Å ³	2799.3(4)	2939(1)
<i>Z</i>	4	4
<i>d</i> _{calc} , g cm ⁻³	1.75	2.07
cryst size, mm	0.45 × 0.26 × 0.19	0.38 × 0.38 × 0.45
μ(Mo Kα), cm ⁻¹	55.2	71.8
transm factors (min, max)	0.675, 0.965	0.485, 1.000
diffractometer	Siemens/Stoe AED2	Nicolet (Siemens) Autodiff
radiation (graphite monochromated)	Mo Kα (λ _α = 0.710 73 Å)	Mo Kα (λ _α = 0.710 73 Å)
<i>T</i> , °C	20 ± 1	20 ± 1
scan method	ω	ω
scan range, deg	4 < 2θ < 50	3 < 2θ < 51
no. of unique data	4858	5396
no. of unique data with <i>F</i> _o ² ≥ 3σ(<i>F</i> _o ²)	3996	3494
abs corr	ψ scans	ψ scans
no. of refined params	326	293
<i>R</i> ^b	0.029	0.038
<i>R</i> _w ^c	0.044	0.049
quality of fit indicator ^d	0.95	0.96
weighting param <i>g</i> ^e	0.0015	0.0013
largest shift/esd final cycle	0.001	0.001
final diff Fourier max peak, trough, e/Å ³	0.83, -0.96	1.39, -1.06

^a **3**, [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,*P*}(S₂CNMe₂)]·CHCl₃; **7**, [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,*P*}(S₂CNMe₂)I₂]^{1/2}·CH₂Cl₂. ^b *R* = Σ||*F*_o| - |*F*_c||/Σ|*F*_o|. ^c *R*_w = [Σw(|*F*_o| - |*F*_c|)²/Σw|*F*_o|²]^{1/2}. ^d Quality of fit = [Σw(|*F*_o| - |*F*_c|)²/(N_{observns} - N_{parameters})]^{1/2}. ^e *w*⁻¹ = [σ²(|*F*_o|) + *g*|*F*_o|²].

formation of octahedral complexes.²¹ Assuming that the C⁺P and S₂CR⁻ groups maintain their chelating function, these octahedral Pt(IV) complexes can exhibit three different structures (Chart 2). The dichloro complexes [Pt(C⁺P)(S₂CR)Cl₂] (R: NMe₂ (**5**), OEt (**8**)) show two absorptions at about 310 and 270 cm⁻¹ (see Experimental Section) assignable to the ν(Pt-Cl) stretching vibrations, but this feature does not allow us to distinguish between any of the three possible isomers because two IR-active absorptions should be expected for each one of them (Chart 2).

In the IR spectra of complexes **5–7** the ν_{st}(C-N) absorptions appear at higher frequencies than in their precursor (complex **3**), showing a correlation between increasing oxidation state and the C-N stretching frequency as has been observed previously.²²

The ³¹P{¹H} NMR spectra of complexes **5–10** (see Table 1) show one singlet at about 20.0 ppm with the corresponding ¹⁹⁵Pt satellites, which indicates that only one diastereomer is present in each case. The smaller Pt-P coupling constant in these complexes (¹*J*_{Pt-P} = 2600–2700 Hz) with respect to that in their Pt(II) precursors (**3** and **4**) is consistent with a higher oxidation state of the metal center.²³

The ¹H NMR signals corresponding to the C⁺P fragment show, for all complexes, the same pattern (see Table 2): (a) two signals at ca. 2.0 ppm due to the

methyl groups; (b) two signals for the diastereotopic H atoms of the methylene group, which appear as an AB system with the corresponding ¹⁹⁵Pt satellites, and (c) some multiplets corresponding to the aromatic hydrogens. The inequivalence of both methyl groups and both H atoms of -CH₂- can be explained in two different ways: (a) complexes **5–10** show either an OC-6-42 or OC-6-32 geometry, which excludes the existence of any symmetry plane, or (b) these complexes show the geometry OC-6-14 but a hindered P-C_{ipso} bond rotation process precludes the equatorial plane of the molecule to act as a mirror plane. In the last case, temperature-dependent ¹H NMR spectra should be expected as for complexes **3** and **4**. However we have observed that the ¹H NMR spectrum of complex [Pt(C⁺P)(S₂CNMe₂)I₂] (**7**) remain invariable even at 60 °C, suggesting the structure OC-6-42 or OC-6-32 as more probable. The X-ray study on complex [Pt(C⁺P)(S₂CNMe₂)I₂] (**7**) indicates an OC-6-42 geometry for it. The similarity of the IR and NMR spectra of these Pt(IV) complexes suggests the same geometry for all of them; i.e., the oxidative addition takes place in a stereoselective mode in all cases.

The most common mechanism in oxidative addition reactions of halogens (X₂) on square-planar platinum(II) complexes is the S_N2 type,²⁴ the process being initiated by end-on coordination of X₂ formed by donation of electron density from the metal to the X₂ molecule (**A**, Scheme 1). The basicity of the platinum center in [Pt(C⁺P)(S₂CR)] (**3**, **4**) becomes apparent considering their ability to form donor-acceptor Pt→M bonds.²⁵ The subsequent steps very often result in a *trans* addition of the halogen (Scheme 1, path 2) yielding

(21) (a) Forniés, J.; Fortuño, C.; Gómez, M. A.; Menjón, B. *Organometallics* **1993**, *12*, 4368. (b) van Beek, J. A. M.; van Koten, G.; Wehman-Ooyevaar, I. C. M.; Smeets, W. J. J.; van der Sluis, P.; Spek, A. L. *J. Chem. Soc., Dalton Trans.* **1991**, 883.

(22) Edgar, B. L.; Duffy, D. J.; Palazzotto, M. C.; Pignolet, L. H. *J. Am. Chem. Soc.* **1973**, *95*, 1125.

(23) (a) Parish, R. V. *NMR, NQR, EPR and Mössbauer Spectroscopy in Inorganic Chemistry*; Ellis Horwood Ltd: New York, 1990. (b) Pregosin, P. S. *³¹P and ¹³C NMR of Transition Metal Phosphine Complexes*; Springer-Verlag: Berlin, 1979.

(24) Skinner, C. E.; Jones, M. M. *J. Am. Chem. Soc.* **1969**, *91*, 4405.

(25) Sicilia, V. Unpublished results.

Table 4. Atomic Coordinates ($\times 10^4$) and Equivalent Isotropic Displacement Coefficients ($\text{\AA}^2 \times 10^3$) for $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{CNMe}_2)]\cdot\text{CHCl}_3$


	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> (eq) ^a
Pt	2446(1)	3060(1)	3161(1)	43(1)
P	3531(1)	2037(1)	4040(1)	40(1)
S(1)	1669(1)	4361(1)	3800(1)	61(1)
S(2)	1318(1)	4104(1)	2194(1)	62(1)
N	556(4)	5772(4)	2779(3)	66(2)
C(1)	1100(4)	4871(4)	2909(3)	54(2)
C(2)	434(6)	6419(6)	3407(5)	86(3)
C(3)	36(7)	6132(6)	1998(4)	88(3)
C(4)	4384(4)	1356(4)	3522(3)	44(2)
C(5)	5344(5)	794(4)	3862(3)	53(2)
C(6)	5919(5)	309(5)	3411(3)	60(2)
C(7)	5531(6)	394(5)	2630(4)	68(3)
C(8)	4585(5)	961(5)	2299(3)	62(2)
C(9)	3999(4)	1469(4)	2746(3)	48(2)
C(10)	2997(6)	2154(5)	2404(3)	62(2)
C(11)	2745(4)	1036(4)	4385(3)	41(2)
C(12)	2814(5)	12(4)	4148(3)	52(2)
C(13)	2220(5)	-763(5)	4369(4)	69(3)
C(14)	1517(6)	-547(6)	4831(4)	78(3)
C(15)	1432(5)	449(6)	5063(3)	67(2)
C(16)	2029(4)	1272(5)	4843(3)	51(2)
C(17)	1870(6)	2326(6)	5106(4)	71(3)
C(18)	4566(4)	2487(4)	4893(3)	46(2)
C(19)	4732(5)	1962(4)	5580(3)	54(2)
C(20)	5561(6)	2250(6)	6222(4)	72(3)
C(21)	6264(6)	3057(6)	6185(5)	80(3)
C(22)	6121(5)	3594(6)	5511(5)	80(3)
C(23)	5278(5)	3331(4)	4855(4)	60(2)
C(24)	5190(6)	3916(5)	4131(4)	84(3)
C(25)	1334(5)	6370(4)	6304(2)	158(7)
Cl(1)	1665(10)	5913(10)	7251(4)	120(4)
Cl(2)	2227(11)	7367(7)	6156(9)	255(11)
Cl(3)	1643(20)	5315(13)	5782(9)	153(6)
Cl(4)	2335(11)	7382(9)	6470(7)	161(5)
Cl(5)	2026(11)	5780(14)	7179(5)	181(6)
Cl(6)	1424(23)	5546(17)	5551(10)	223(11)

^a Equivalent isotropic *U* defined as one-third of the trace of the orthogonalized U_{ij} tensor.

Table 5. Selected Bond Lengths (\AA) and Angles (deg) for $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{CNMe}_2)]$ (3)


Pt-C(10)	2.063(7)	C(1)-S(1)	1.717(6)
Pt-P	2.223(1)	C(1)-S(2)	1.718(6)
Pt-S(1)	2.392(2)	C(1)-N	1.336(7)
Pt-S(2)	2.354(1)		
S(1)-Pt-S(2)	74.1(1)	S(2)-C(1)-N	123.1(5)
S(1)-Pt-P	108.2(1)	S(1)-C(1)-N	124.1(5)
P-Pt-C(10)	84.2(2)	C(1)-N-C(2)	121.2(5)
S(2)-Pt-C(10)	93.5(2)	C(1)-N-C(3)	121.0(6)
S(1)-C(1)-S(2)	112.8(3)	C(2)-N-C(3)	117.8(5)

Chart 2



the isomer OC-6-14. In our case, however, the couple of enantiomers OC-6-42 were solely obtained in all cases. These results mean that either the halogen molecule adds at once in a concerted fashion at *cis* positions or that isomerization must follow some of the proposed steps for the S_N2 mechanism.

Scheme 1



We tried to find some experimental evidence about the possible reaction mechanism by focusing our study on the iodination reaction of $[\text{Pt}(\text{C}^{\text{P}})(\text{S}_2\text{CNMe}_2)]$ (3). Assuming that the oxidative addition of halogen to $[\text{Pt}(\text{C}^{\text{P}})(\text{S}_2\text{CNMe}_2)]$ (3) takes place in a concerted fashion, iodination of 3 in the presence of chloride anions should exclusively give products derived from the already formed complex $[\text{Pt}(\text{C}^{\text{P}})(\text{S}_2\text{CNMe}_2)\text{I}_2]$ (7) by simple halide substitution (C or C'). In the reaction of $[\text{Pt}(\text{C}^{\text{P}})(\text{S}_2\text{CNMe}_2)\text{I}_2]$ (7) with NEt_3BzCl in a 1:1 molar ratio ($[\text{I}^2\text{H}]$ chloroform, room temperature) we observe the appearance of a single new product (C or C'; $^{31}\text{P}\{^1\text{H}\}$ NMR: δ 20.69, $^1J_{\text{Pt-P}} = 2634.4$ Hz) which means that halide substitution on 7 proceeds specifically at a certain position. This substitution reaction depends on the temperature and does not take place at -60°C even in a 1:3 molar ratio. Because of that, we performed the iodination of 3 in the presence of NEt_3BzCl (1:3) at -78°C in CD_2Cl_2 . The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the reaction mixture at this temperature shows signals corresponding to 7 and C (or C') in the integrated ratios 0.926:1. The presence of C (or C') as a reaction product allows us to accept the S_N2 mechanism including an additional isomerization step to explain the *cis* arrangement of the halogen atoms in the final product.

Aiming to know which intermediate species (OC-6-14 isomer or B⁺) undergoes further isomerization, we performed the reaction of $[\text{Pt}(\text{C}^{\text{P}})(\text{S}_2\text{CNMe}_2)]$ (3) with I_2 at -78°C in CD_2Cl_2 in a NMR tube monitoring the progress of the reaction at $-78, -65, -50, -35, -20, 0$, and 20°C by $^{31}\text{P}\{^1\text{H}\}$ NMR. At -78°C , the reaction mixture show the presence of two different products, the final one (7) and an intermediate (X) ($^{31}\text{P}\{^1\text{H}\}$: δ 16.04, $^1J_{\text{Pt-P}} = 2376$ Hz. ^1H : δ Pt-CH₂, 3.18 (d), 5.43 (d),




Figure 2. Molecular structure of $[\text{Pt}(\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P})(\text{S}_2\text{CNMe}_2)\text{I}_2]$ (**7**).

$^2J_{\text{H-H}} = 12.5$ Hz, $^2J_{\text{Pt-H}} = 75$ Hz; *o*-tolyl, 1.78 (s), 1.85 (s); S_2CNMe_2 , 3.25 (s), 3.28 (s)). As temperature increases, the amount of **7** grows meanwhile the amount of (**X**) decreases with **7** being the only product in the reaction mixture at room temperature (20 °C). The possible conversion of **X** to **7** at low temperatures ($T < -60$ °C) even in the presence of NEt_3BzCl and the very different δ values observed for the signals corresponding to both H atoms of the Pt-CH₂ group in this species (**X**) suggest that this intermediate (**X**) can be either **A**, **B**⁺, or **B'**⁺ (Scheme 1) rather than the Pt(IV) isomer OC-6-14 whose isomerization probably would need higher temperatures²⁶ and in which both H atoms of the Pt-CH₂- group would have very similar chemical environments. However, we cannot determine the exact nature of the intermediate (**X**). From this observation we suggest that, in all likelihood, isomerization occur at the Pt(IV) cation (**B**⁺, Scheme 1, path 1) rather than at the coordinatively saturated product OC-6-14.

In conclusion from our study on oxidation reactions of $[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{CNMe}_2)]$ and $[\text{Pt}(\text{C}^{\wedge}\text{P})(\text{S}_2\text{COEt})]$ with chlorine, bromine, and iodine we can say that these reactions do not proceed in a concerted fashion but by an $\text{S}_{\text{N}}2$ mechanism including an isomerization step probably at the five-coordinated Pt(IV) cation intermediate to give a unique final product in each case with a *cis* arrangement of the halide ligands.

Molecular Structure of $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{CNMe}_2)\text{I}_2]$ (7**).** The stereochemistry of this platinum complex, as determined by single-crystal experiments, is illustrated in Figure 2 together with the atomic numbering scheme. General crystallographic information is collected in Table 3. Final atomic positional parameters are listed in Table 6. Relevant bond distances and angles are listed in Table 7.

The molecule shows a slightly distorted octahedral geometry around the platinum atom. One iodine atom [I(1)] and the carbon of the C[∧]P group are in the axial sites, and the I(2), the bidentate dimethyldithiocarbamate ligand, and the P of the C[∧]P group are located in the equatorial positions, with the angle between the line 1 (I(1), Pt, C(10)) and the perpendicular to the equatorial plane defined by the atoms I(2), S(1), S(2), Pt, and P

Table 6. Atomic Coordinates ($\times 10^4$) and Equivalent Isotropic Displacement Coefficients ($\text{\AA}^2 \times 10^3$) for $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{CNMe}_2)\text{I}_2] \cdot \frac{1}{2}\text{CH}_2\text{Cl}_2$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> (eq) ^a
Pt	2572(1)	1196(1)	1372(1)	29(1)
I(1)	197(1)	1623(1)	1715(1)	47(1)
I(2)	2670(1)	2632(1)	472(1)	45(1)
P	3714(3)	1924(2)	2276(1)	30(1)
S(1)	2359(3)	-240(2)	1954(1)	40(1)
S(2)	1716(3)	149(2)	493(1)	40(1)
N	1153(9)	-1470(6)	1109(5)	39(3)
C(11)	3255(10)	2973(7)	2735(5)	31(3)
C(18)	4106(11)	1086(7)	2974(6)	39(4)
C(9)	5920(11)	2856(9)	1971(7)	46(4)
C(23)	5265(13)	657(9)	2945(7)	53(5)
C(6)	6270(10)	1694(8)	795(6)	40(4)
C(4)	5084(10)	2155(8)	1804(5)	37(4)
C(1)	1656(10)	-643(7)	1178(6)	37(4)
C(15)	1755(12)	4215(9)	2754(7)	49(4)
C(12)	3791(11)	3316(8)	3377(6)	38(4)
C(19)	3276(12)	854(8)	3515(6)	43(4)
C(10)	4306(11)	831(9)	1026(7)	47(4)
C(2)	458(14)	-1755(9)	468(7)	59(5)
C(5)	5254(10)	1567(8)	1215(6)	38(4)
C(17)	4890(12)	2878(9)	3766(7)	53(5)
C(16)	2239(10)	3443(7)	2431(6)	36(3)
C(14)	2255(13)	4527(9)	3382(8)	57(5)
C(7)	7113(12)	2386(10)	979(7)	56(5)
C(24)	2040(12)	1255(10)	3560(7)	58(5)
C(8)	6939(12)	2969(10)	1547(8)	54(5)
C(3)	1143(15)	-2109(9)	1728(7)	69(6)
C(13)	3266(12)	4082(8)	3694(7)	49(4)
C(20)	3661(16)	235(9)	4046(7)	63(5)
C(22)	5595(15)	19(10)	3478(8)	69(6)
C(21)	4816(19)	-173(11)	4028(8)	79(7)
Cl(1)	-1099(20)	4673(16)	-721(13)	214(9)
Cl(2)	809(18)	5576(14)	108(12)	190(7)
C(50)	-27(35)	5579(26)	-709(17)	142(18)

^a See footnote a in Table 4.

Table 7. Selected Bond Lengths (Å) and Angles (deg) for $[\text{Pt}\{\text{CH}_2\text{C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-C,P}\}(\text{S}_2\text{CNMe}_2)\text{I}_2]$ (**7**)

Pt-I(1)	2.764(1)	Pt-P	2.323(3)
Pt-I(2)	2.668(1)	Pt-S(1)	2.353(3)
Pt-C(10)	2.089(12)	Pt-S(2)	2.400(3)
I(1)-Pt-I(2)	91.4(1)	S(1)-Pt-S(2)	73.6(1)
I(1)-Pt-P	103.1(1)	P-Pt-S(1)	97.1(1)
I(1)-Pt-S(1)	89.2(1)	I(2)-Pt-P	94.2(1)
I(1)-Pt-S(2)	86.7(1)	I(2)-Pt-S(2)	94.7(1)
I(1)-Pt-C(10)	174.9(3)	P-Pt-C(10)	81.9(3)

being 7.50°. ¹⁴ Both P and S(2) atoms are located outside plane 1 at 0.1368 and 0.1740 Å, respectively.

Angles around the platinum are rather different from 90° because of the small bite angles of the chelate ligands. The Pt-I distances are similar to the observed ones in other Pt(IV) iodo complexes²⁷ and slightly different from each other, showing the stronger *trans* influence of the C(sp³) σ -bond.¹⁵ The Pt-C, Pt-P, and Pt-S distances are similar to the corresponding ones in the starting material (complex **3**). The metallacycle formed by the atoms Pt, C(10), C(5), C(4), and P is not planar, the atoms C(4) and C(10) being above and below the best least-squares-calculated plane 0.3826 and 0.5218 Å, respectively. The two *o*-tolyl groups are planar, and they are oriented by forming an angle of 112.43° between them and 57.115 and 84.80°, respectively, with the equatorial plane.

The S_2CNMe_2 ligand is basically planar and almost coplanar with the equatorial plane (interplanar angle 5.77°). Like in complex **3**, the structural parameters

(26) Wehman-Ooyevaar, I. C. M.; Drenth, W.; Grove, D. M.; van Koten, G. *Inorg. Chem.* **1993**, *32*, 3347.

(27) (a) Cheetman, A. K.; Puddephatt, R. J.; Zalkin, A.; Templeton, D. H.; Templeton, L. K. *Inorg. Chem.* **1976**, *15*, 2997. (b) Casalone, G.; Mason, R. *Inorg. Chim. Acta* **1973**, *7*, 429.

concerning the $\text{Me}_2\text{NCS}_2^-$ ligand point out to a high degree of C–N double-bond character.

Experimental Section

General Procedures and Materials. Elemental analyses were determined by using a Perkin-Elmer 240-B microanalyzer. IR spectra were recorded on a Perkin-Elmer 599 spectrophotometer (Nujol mulls between polyethylene plates in the range 4000–200 cm^{-1}). NMR spectra were recorded on either a Varian XL-200 or a Varian Unity 300 NMR spectrometer using the standard references. [*trans*-PtCl₂(PhCN)₂] was prepared by the literature method.²⁸

Safety Note. Perchlorate salts are potentially explosive. Only small amounts of material should be prepared, and these should be handled with great caution!

[Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)₂] (1). A mixture of [*trans*-PtCl₂(PhCN)₂] (2.0 g, 4.235 × 10⁻³ mol) and tri-*o*-tolylphosphine (1.29 g, 4.235 × 10⁻³ mol) in 2-methoxyethanol (25 mL) was refluxed for 30 min. The initial suspension dissolved to give a clear pale yellow solution, and then a white crystalline solid was deposited. After cooling, the solid was filtered off, washed with methanol and diethyl ether, and identified as complex **1** (yield 1.9 g, 84%). Anal. Calcd for C₄₂Cl₂H₄₀P₂Pt₂: C, 47.26; H, 3.78. Found: C, 47.54; H, 3.86. IR (cm⁻¹): C⁻P, 467 (vs), 479 (vs), 486 (vs), 527 (vs, sh), 564 (vs), 585 (vs), 750 (vs), 759 (vs), 784 (m), 806 (m), 1568 (m), 1584 (m), 1590 (m); ν(Pt–Cl), 249 (s), 255 (s).

[Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(NCMe)₂]ClO₄ (2). To a suspension of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)₂] (**1**) (0.6407 g, 0.6 × 10⁻³ mol) in acetonitrile (20 mL) was added AgClO₄ (0.2488 g, 1.2 × 10⁻³ mol), and the mixture was stirred for 4 h at room temperature. The AgCl precipitated was filtered off and washed with 5 mL of acetonitrile. Evaporation of the resulting solution to dryness followed by addition of OEt₂ to the residue afforded a white solid which was identified as **2** (yield 0.71 g, 87%). Anal. Calcd for C₂₅ClH₂₆N₂O₄PtP: C, 44.16; H, 3.85; N, 4.12. Found: C, 43.79; H, 3.66; N, 3.95. IR (cm⁻¹): C⁻P, 466 (s), 479 (s), 487 (s), 516 (m), 531 (s), 566 (s), 589 (s), 759 (s), 772 (s), 793 (s), 1568 (w), 1589 (w); NCMe, 2288 (m), 2298 (m), 2324 (m); ClO₄⁻, 1083 (vs, br), 623 (s).

[Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)] (3). To a stirred suspension of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)₂] (**1**) (0.8544 g, 0.8 × 10⁻³ mol) in THF (40 mL) was added AgClO₄ (0.3317 g, 1.6 × 10⁻³ mol), and the mixture was stirred at room temperature for 4 h. The AgCl precipitated was filtered off, and the resulting solution was evaporated almost to dryness. The residue was treated with 30 mL of methanol, and the addition of NaS₂CNMe₂·2H₂O (0.2867 g, 1.6 × 10⁻³ mol) to the solution gave rise to a white solid, which was filtered off, recrystallized from CH₂Cl₂/*n*-hexane, and identified as **3** (yield 0.88 g, 89%). Anal. Calcd for C₂₄H₂₆NPtS₂: C, 46.59; H, 4.24; N, 2.26. Found: C, 46.71; H, 3.89; N, 2.24. IR (cm⁻¹): C⁻P, 465 (s), 476 (s), 486 (s), 514 (m), 529 (m), 562 (s), 583 (s), 754 (vs), 765 (vs), 1583 (w); -S₂CNMe₂, 1543 (vs) (ν(C–N)); 975 (vs) (ν(C–S)).

Synthesis of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂COEt)] (4). To the resulting solution of the treatment of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(μ-Cl)₂] (**1**) (0.5447 g, 0.51 × 10⁻³ mol) with AgClO₄ (0.2116 g, 1.02 × 10⁻³ mol) in THF (20 mL), after elimination of the AgCl, was added KS₂COEt (0.1636 g, 1.02 × 10⁻³ mol) and the mixture stirred for 15 min at room temperature. Then, the solution was evaporated to dryness and *n*-pentane (100 mL) was added to the residue. After filtration, the resulting solution was concentrated to 10 mL and cooled to ca. 5 °C. The yellow solid precipitated was filtered off, heated at 70 °C for 24 h, and identified as **4** (yield 0.474 g, 75%). Anal. Calcd for C₂₄H₂₅OPtS₂: C, 46.52; H,

4.07. Found: C, 46.66; H, 4.08. IR (cm⁻¹): C⁻P, 458 (m), 477 (s), 487 (m), 505 (m), 586 (s), 758 (vs), 1567 (w), 1586 (w); -S₂COEt, 1031 (vs), 1123 (vs), 1239 (vs).

[Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)X₂] (X = Cl (5**), Br (**6**), I (**7**)). X = Cl.** To a pale yellow solution of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)] (**3**) (0.6 g, 0.9698 × 10⁻³ mol) in CH₂Cl₂ (20 mL) was added a slight excess of Cl₂ in CCl₄ (1.1 × 10⁻³ mol), and immediately the solution turned yellow. After 10 min of stirring at room temperature the solution was evaporated to dryness and OEt₂ (20 mL) was added to the residue to give a yellow solid, which was filtered off, dried, and identified as **5** (yield 0.515 g, 77%). Anal. Calcd for C₂₄Cl₂H₂₆NPtS₂: C, 41.80; H, 3.80; N, 2.03. Found: C, 41.87; H, 3.87; N, 1.98. IR (cm⁻¹): C⁻P group, 455 (s), 475 (s), 487 (s), 506 (m), 522 (s), 536 (s), 551 (s), 567 (m), 583 (m), 751 (vs), 759 (s), 767 (m), 1588 (w); -S₂CNMe₂, 1561 (vs) (ν(C–N)), 978 (w) (ν(C–S)); Pt–Cl, 272 (m), 311 (m).

Complexes 6 (X = Br) and 7 (X = I) were prepared similarly.

Complex 6 (X = Br). **3** (0.2877 g, 0.465 × 10⁻³ mol) and Br₂ in CCl₄ (0.558 × 10⁻³ mol) were used. Yield: 0.323 g, 89%. Anal. Calcd for Br₂C₂₄H₂₆NPtS₂: C, 37.03; H, 3.37; N, 1.80. Found: C, 37.36; H, 3.27; N, 1.71. IR (cm⁻¹): C⁻P group, 455 (s), 475 (s), 487 (s), 507 (m), 523 (s), 536 (s), 551 (s), 567 (m), 583 (m), 751 (vs), 757 (s), 765 (m), 1636 (w); -S₂CNMe₂, 1558 (vs) (ν(C–N)), 977 (w) (ν(C–S)).

Complex 7 (X = I). **3** (0.3036 g, 0.4907 × 10⁻³ mol) and I₂ (0.137 g, 0.54 × 10⁻³ mol) were used. Yield: 0.385 g, 90%. Anal. Calcd for C₂₄H₂₆I₂NPtS₂: C, 33.04; H, 3.00; N, 1.60. Found: C, 32.93; H, 3.38; N, 1.46. IR (cm⁻¹): C⁻P group, 454 (m), 472 (m), 487 (w), 505 (w), 519 (m), 536 (m), 550 (w), 567 (w), 580 (w), 748 (vs), 762 (s), 1584 (w); -S₂CNMe₂, 1549 (vs) (ν(C–N)), 976 (w) (ν(C–S)).

[Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂COEt)X₂] (X = Cl (8**), Br (**9**), I (**10**)). X = Cl.** To a pale yellow solution of [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂COEt)] (**4**) (0.2497 g, 0.403 × 10⁻³ mol) in CH₂Cl₂ (25 mL) was added a little excess of Cl₂ in CCl₄ (0.462 × 10⁻³ mol). The mixture was stirred at room temperature for 5 min, and then the solution was evaporated to dryness. Addition of OEt₂ (20 mL) to the residue afforded a solid, which was identified as **8** (yield 0.223 g, 80%). Anal. Calcd for C₂₄Cl₂H₂₅OPtS₂: C, 41.74; H, 3.65. Found: C, 41.71; H, 3.66. IR (cm⁻¹): C⁻P group, 453 (s), 475 (s), 488 (s), 503 (w), 522 (s), 535 (s), 552 (m), 567 (w), 583 (m), 749 (s), 761 (vs), 1567 (w), 1591 (m); -S₂COEt, 1030 (s), 1298 (vs); Pt–Cl, 276 (m), 306 (m).

Complexes 9 (X = Br) and 10 (X = I) were prepared similarly.

Complex 9 (X = Br). **4** (0.2010 g, 0.32 × 10⁻³ mol) and Br₂ in CCl₄ (0.372 × 10⁻³ mol) were used. Yield: 0.2 g, 80%. Anal. Calcd for Br₂C₂₄H₂₅OPtS₂: C, 36.98; H, 3.23. Found: C, 36.73; H, 3.50. IR (cm⁻¹): C⁻P group, 453 (m), 474 (m), 487 (m), 502 (w), 521 (m), 535 (m), 552 (w), 567 (w), 582 (w), 758 (m, sh), 1565 (w), 1591 (m); -S₂COEt, 1032 (s), 1291 (vs).

Complex 10 (X = I). **4** (0.1554 g, 0.25 × 10⁻³ mol) and I₂ (73 mg, 0.287 × 10⁻³ mol) were used. Yield: 0.157 g, 72%. Anal. Calcd for C₂₄H₂₅I₂OPtS₂: C, 33.00; H, 2.88. Found: C, 32.62; H, 2.96. IR (cm⁻¹): C⁻P group, 454 (m), 474 (m), 486 (m), 503 (m), 521 (m), 536 (m), 551 (m), 567 (m), 580 (m), 744 (m), 1567 (w), 1591 (w); -S₂COEt, 1032 (s), 1288 (vs).

Crystallographic Studies. [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)]·CHCl₃ (**3**). Important crystal data and data collection parameters for complex **3** are listed in Table 3. The complex [Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)]·CHCl₃ crystallizes in space group *P*2₁/*c* with *Z* = 4. A parallelepiped-shaped crystal mounted on the tip of a glass fiber with epoxy cement was used for geometric and intensity data collection. Four circle diffractometer data were taken at 293 ± 1 °C. Lattice dimensions and type were determined by routine procedures and verified by oscillation photography. Cell constants were refined from 2θ values of 48 reflections including Friedel pairs (27.7 < 2θ < 29.5°). During intensity

(28) (a) Hartley, F. R. *Organomet. Chem. Rev. A* **1970**, 6, 119. (b) Uchiyama, T.; Toshiyasu, Y.; Nakamura, Y.; Miwa, T.; Kawaguchi, S. *Bull. Chem. Soc. Jpn.* **1981**, 54, 181.

data collection, three monitor reflections were measured at regular intervals. These check reflections did not vary appreciably in intensity during the course of data collection. For the diffractometer data a measured absorption correction was applied, based on 10 complete Ψ -scans of reflections with diffractometer angle χ near 90° .

The structure of compound **3** was solved by direct methods and developed and refined in series of alternating difference Fourier maps and least-squares analyses using all data and the program SHELXTL-PLUS.²⁹ The chlorine atoms of the chloroform solvent were disordered over two sites (occupancy 0.5) sharing the same C atom. The C–Cl distances were fixed at 1.77 Å. All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included in calculated positions and refined as riding atoms [C–H = 0.96 Å and $U = 0.0872 \text{ Å}^3$]. Residuals and other final refinement parameters are listed in Table 3.

[Pt{CH₂C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)I₂]} \cdot $\frac{1}{2}$ CH₂Cl₂ (7**).** Important crystal data and data collection parameters for complex **7** are listed in Table 3. Diffraction data were measured by Crystallogics Co.³⁰ The complex [Pt{CH₂-C₆H₄P(*o*-tolyl)₂-C,P}(S₂CNMe₂)I₂]} \cdot $\frac{1}{2}$ CH₂Cl₂ crystallizes in space group $P2_1/c$ with $Z = 4$. A parallelepiped-shaped crystal mounted on the tip of a glass fiber with epoxy cement was used for geometric and intensity data collection. Four-circle diffractometer data were taken at $293 \pm 1^\circ\text{C}$. Lattice dimensions and type were determined by routine procedures and verified by oscillation photography. Cell constants were refined from 2θ

values of 15 reflections including Friedel pairs ($20 < 2\theta < 30^\circ$). During intensity data collection, six monitor reflections were measured at regular intervals. These check reflections did not vary appreciably in intensity during the course of data collection. For the diffractometer data a measured absorption correction was applied, based on 6 complete Ψ -scans of reflections with diffractometer angle χ near 90° .

The structure of compound **7** was solved by direct methods and developed and refined in series of alternating difference Fourier maps and least-squares analyses using all data and the program SHELXTL-PLUS.²⁹ The solvent molecule lays near an inversion center and is disordered over two positions. The occupancy of the atoms of the solvent molecule was fixed at 0.5. All non-hydrogen atoms except for those of the solvent were refined anisotropically. All hydrogen atoms except for the solvent one were included in calculated positions and refined as riding atoms [C–H = 0.96 Å and $U = 0.0832 \text{ Å}^3$]. A difference map following convergence showed two peaks higher than 1 e/Å^3 (1.39 and 1.21 e/Å^3). These two peaks lay very close to the platinum atoms (0.91 and 0.95 Å) and had no chemical significance. Residuals and other final refinement parameters are listed in Table 3.

Acknowledgment. We thank the Dirección General de Investigación Científica y Técnica (Spain) for financial support (Proyect PB92-0364) and Dr. B. Menjón for his very valuable comments.

Supporting Information Available: Tables of anisotropic thermal parameters, H positional and thermal parameters, and complete bond distances and bond angles (8 pages). Ordering information is given on any current masthead page.

OM950861U

(29) SHELXTL-PLUS: *Software Package for the Determination of Crystal Structures*, Release 4.0; Siemens Analytical X-ray Instruments, Inc.: Madison, WI, 1990.

(30) Crystallogics Co., P.O. Box 82286, Lincoln, NE.