Inorganic Chemistry

Construction of an M₃L₂A₆ Cage with Small Windows from a Flexible Tripodal Ligand and Cu(hfac)₃

Mari Ikeda,[†] Keiji Ohno,[‡] Yuki Kasumi,[‡] Shunsuke Kuwahara,^{†,‡} and Yoichi Habata^{*,†,‡}

[†]Research Centre for Materials with Integrated Properties and [‡]Department of Chemistry, Faculty of Science, Toho University, 2-2-1 Miyama, Funabashi, Chiba 274-8510, Japan

Supporting Information

ABSTRACT: An $M_3L_2A_6$ cage has been prepared with small windows from a tripodal ligand, L, and Cu(hfac)₂. Cold spray ionization mass spectrometry of a mixture of L and Cu(hfac)₂ revealed the formation of a Cu₃L₂hfac₆ cage in solution. X-ray crystallography showed that the Cu₃L₂hfac₆ cage included neutral molecules such as THF and CHCl₃. Furthermore, the six hfac anions have been shown to play an important role in holding neutral guest molecules securely in place.

T he formation of molecular cages by self-assembly has attracted considerable attention from many researchers,¹ and various tripodal ligands have been reported in the literature in this particular context.² For example, Fujita et al.^{2u} reported that the combination of 1,3,5-tris(4-pyridylmethyl)benzene and enPd(NO₃)₂ led to the formation of a cagelike complex, and Kaim et al.^{2v} reported that different combinations of AgBF₄ or CuI and 1,3,5-tris(benzimidazol-1-ylmethyl)-2,4,6-trimethylbenzene led to the formation of metallocages containing BF₄⁻ or CuI₃⁻ anions in their cavity. In these cases, the counteranions of the metal salts (e.g., NO₃⁻, BF₄⁻, and I⁻) did not participate in the formation of cage structures, indicating that these cages contained large windows through which the anions could readily pass.

Hexafluoroacetylacetonate (hfac) salts have been used in a number of studies to form supramolecular systems such as coordination polymers,³ metallomacrocycles,⁴ and clusters.⁵ In these supramolecular structures, the hfac anions bind tightly to the metal cations, and Cu(hfac)₂ is about 280 Å^{3.6} On the basis of these observations, it was envisaged that the combination of hfac salts with a tripodal ligand would lead to the formation of a new type of cage with small windows, where the hfac anions would effectively form a barrier, preventing escape of the guest molecules and leading to enhanced binding properties (Figure 1a). Herein, we report the combination of a flexible tripodal ligand and Cu(hfac)₂ to construct a new type of three-dimensional cage structure with small windows.

A new tripodal ligand (Figure 1b, L) was prepared in 45% yield by the reaction of tris(4-bromobenzeyl)amine (2) with 3pyridineboronic acid.

Single crystals of the $Cu(hfac)_2$ complex with L were obtained as follows. A solution of L in CHCl₃ was mixed with a solution of $Cu(hfac)_2$ in CH₃OH to give a green powder, which was recrystallized from a mixture of tetrahydrofuran (THF)/hexane (1:1, v/v) to give single crystals of the desired material.



Figure 1. Side view (a) and top view (b) of a cage structure with small windows and our new flexible tripodal ligand (c). Red, green, and blue tubes mean L, Cu^{2+} , and hfac anions, respectively.

Recrystallization of the green powder from a mixture of CHCl₃ and hexane (1:1, v/v) gave the [CHCl₃ \subset Cu₃L₂hfac₆] complex. The X-ray crystal structure of the [CHCl₃ \subset Cu₃L₂hfac₆] complex is shown in Figure S3 in the Supporting Information (SI).

Elemental analysis of the complex indicated that it was 2:3 L/ $Cu(hfac)_2$. Figure 2 shows the crystal structure of the $Cu(hfac)_2$ complex with L. The complex formed an M3L2A6-type cage containing two THF molecules in its cavity. Each Cu²⁺ ion is sixcoordinated by four O atoms of the hfac anions and two N atoms of the pyridines of **L**. The four O atoms and the Cu^{2+} ion lie in the same plane, and the N-Cu-N angles are in the range 174.7-178.5°. The mean N(pyridine)-Cu distance was found to be 2.000 Å and therefore comparable to that of a pyridine/ $Cu(hfac)_2$ complex.^{3k} The void volume of the $Cu_3L_2hfac_6$ cage was estimated to be 285 Å^{3,7} Lindoy et al.⁸ reported that the two di- β -diketone ligands 1,10-(1,4-phenylene)dihexane-1,3-dione (L¹) and 1,1'-biphenyl-4,4'-diylbis(4,4,4-trifluorobutane-1,3dione) (L^2) formed large neutral open-faced Fe₄L₆ tetrahedral cage complexes. The $\text{Fe}_4 \text{L}_6^1$ cage had a small cavity (void volume = 174 Å³), whereas the $\text{Fe}_4 \text{L}_6^2$ cage had a larger cavity (void volume = 844 Å³). The $\text{Fe}_4 \text{L}_6^1$ and $\text{Fe}_4 \text{L}_6^2$ cages contained one and four THF molecules, respectively. It is therefore important to note that the $Cu_3L_2hfac_6$ cage with a small void volume (285) $Å^3$) and small windows held on tightly to two molecules of THF (total 163.5 Å³)⁹ using its ligand framework and the six hfac anions (Figure 2b). Two atropisomers (Δ and Λ forms) existed for the conformation of L (Figure 2c), and the [2THFCCu₃L₂hfac₆] complex existed in its racemic form in the solid state.

The $[2THF \subset Cu_3L_2hfac_6]$ complex was subjected to thermogravimetric-differential thermal analysis (TG-DTA). As shown in Figure 3, the weight losses at 93 and 176 °C were 2.7 and 5.5%,

Received: October 16, 2013

Inorganic Chemistry



Figure 2. Side view (a), top view without the THF molecules (b), and atropisomers without the THF molecules and hfac anions (c) of the $[2THF \subset Cu_3L_2hfac_6]$ complex. The hfac anions are colored blue in part b.





example indicated that the ${\rm Cu}_3 L_2 h {\rm fac}_6$ cage was holding onto its THF molecules tightly.

Cold spray ionization mass spectrometry (CSI-MS) was used to determine whether L formed a cage structure with $Cu(hfac)_2$ in a solution at 298 K, using a mixture of THF and CH_3OH (1:1, v/v). As shown in Figure 4, a base peak corresponding to [L +



Figure 4. CSI-MS of the $Cu_3L_2hfac_6$ cage in THF/CH₃OH (1:1, v/v) at 298 K. The hfac anions are omitted in the cage models.

 $Cu^{2+} + hfac^{-}$ was observed at m/z 788. Fragment-ion peaks corresponding to $[2L + Cu^{2+} + hfac^{-}]^{+}$ and $[2L + 2Cu^{2+} + 3hfac^{-}]^{+}$ were also observed at m/z 1306 and 1785, respectively. Furthermore, a fragment-ion peak corresponding to $[2L + 3Cu^{2+} + 5hfac^{-}]^{+}$ was observed at m/z 2262. These CSI-MS data strongly supported the idea that the Cu(hfac)₂ complex with L formed a cage structure in solution. These fragment-ion peaks, however, did not include THF molecules because the lack of hfac anions effectively opened the cage windows and allowed the THF molecules to readily escape from the complex.

The binding constant between L and Cu(hfac)₂ was estimated by Cu(hfac)₂-induced titration experiments using UV–vis in CH₃OH (Figure 5 and the associated multimedia file in the SI). log K for the 2:3 $2/Cu(hfac)_2$ complex in CH₃OH was estimated to be 18.1 based on nonlinear least-squares curve fitting.¹¹ Walker and Li¹² reported that log β_2 for a 2:1 4-methylpyridine/ Cu(hfac)₂ complex was much less than that of the current complex at 2.7. The lower log β_2 value of the 4-methylpyridine/



Figure 5. Changes in the UV–vis spectra of L upon the addition of $Cu(hfac)_2$ in CH_3OH .

Inorganic Chemistry

 $Cu(hfac)_2$ system suggested that the $Cu_3L_2hfac_6$ cage would be stabilized by the synergistic coordination of six pyridines.

In conclusion, we have demonstrated that the new tripodal ligand, L, forms an $M_3L_2A_6$ cage with $Cu(hfac)_2$. This is the first reported example of a cage compound with small windows composed of hfac anions. TG-DTA indicated that the hfac anions played an important role in holding the guest molecules tightly. The application of this system to enhance the circular dichroism (CD) intensity of chiral molecules with small CD molecular ellipticity is now underway in our laboratory.

ASSOCIATED CONTENT

Supporting Information

Crystallographic data in CIF format, ¹H and ¹³C NMR spectra of compounds, conditions for the titration experiments, X-ray structure of the $[CHCl_3 \subset Cu_3L_2hfac_6]$ complex, crystal data, and UV-vis spectra and speciation analyses, which are also available as a multimedia file (habata.avi). This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*E-mail: habata@chem.sci.toho-u.ac.jp.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors thank Professor Masatoshi Hasegawa for the TG-DTA measurements. This research was supported by Grants-in-Aid 08026969 and 11011761, a High-Tech Research Center project (2005–2009), and the Supported Program for Strategic Research Foundation at Private Universities (2012–2016) from the Ministry of Education, Culture, Sports, Science and Technology of Japan for Y.H.

REFERENCES

(1) (a) Kusukawa, T.; Yoshizawa, M.; Fujita, M. Kagaku to Kogyo 2002, 55, 1237–1241. (b) Fujita, M.; Yoshizawa, M. New properties and reactions in self-assembled M_6L_4 coordination cages; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2008; pp 277–313. (c) Murase, T.; Fujita, M. Fain Kemikaru 2009, 38, 37–44. (d) Murase, T.; Fujita, M. Chem. Rec. 2010, 10, 342–347. (e) Yoshizawa, M.; Fujita, M. Bull. Chem. Soc. Jpn. 2010, 83, 609–618. (f) Fujita, M.; Murase, T. Noncovalent assemblies: design and synthesis; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2011; pp 7–30. (g) Stang, P. J.; Olenyuk, B. Transition-metal-mediated self-assembly of discrete nanoscopic species with well-defined shapes and geometries; Academic Press: New York, 2000; pp 167–224. (h) Thanasekaran, P.; Lee, C.-C.; Lu, K.-L. Acc. Chem. Res. 2012, 45, 1403–1418.

(2) (a) Bar, A. K.; Chakrabarty, R.; Mukherjee, P. S. Inorg. Chem. 2009, 48, 10880-10882. (b) Bringmann, G.; Breuning, M.; Pfeifer, R.-M.; Schreiber, P. Tetrahedron: Asymmetry 2003, 14, 2225-2228. (c) Cibin, F. R.; Di, B. N.; Doddi, G.; Fares, V.; Mencarelli, P.; Ullucci, E. Tetrahedron 2003, 59, 9971-9978. (d) de, l. R. I.; Mani, F.; Peruzzini, M.; Stoppioni, P. J. Organomet. Chem. 2004, 689, 164-169. (e) El, A. B.; Guenee, L.; Pal, P.; Hamacek, J. Inorg. Chem. 2011, 50, 8588-8597. (f) Fan, J.; Zhu, H.-F.; Okamura, T.-a.; Sun, W.-Y.; Tang, W.-X.; Ueyama, N. Chem.-Eur. J. 2003, 9, 4724-4731. (g) Li, X.-J.; Jiang, F.-L.; Wu, M.-Y.; Zhang, S.-Q.; Zhou, Y.-F.; Hong, M.-C. Inorg. Chem. 2012, 51, 4116-4122. (h) Liu, H.-K.; Sun, W.-Y.; Ma, D.-J.; Tang, W.-X.; Yu, K.-B. Chem. Commun. 2000, 591-592. (i) Liu, H.-K.; Tong, X. Chem. Commun. 2002, 1316-1317. (j) Mann, S.; Huttner, G.; Zsolnai, L.; Heinze, K. Angew. Chem., Int. Ed. Engl. 1997, 35, 2808-2809. (k) Meng, W.-L.; Fan, J.; Okamura, T.-a.; Kawaguchi, H.; Lv, Y.; Sun, W.-Y.; Ueyama, N. Z. Anorg. Allg. Chem. 2006, 632, 1890-1896.

(1) Nakajima, H.; Yasuda, M.; Baba, A. Dalton Trans. 2012, 41, 6602-6606. (m) Reger, D. L.; Semeniuc, R. F.; Smith, M. D. Inorg. Chem. 2003, 42, 8137-8139. (n) Saalfrank, R. W.; Maid, H.; Scheurer, A.; Heinemann, F. W.; Puchta, R.; Bauer, W.; Stern, D.; Stalke, D. Angew. Chem., Int. Ed. 2008, 47, 8941-8945. (o) Shahverdizadeh, G. H.; Hakimi, F.; Mirtamizdoust, B.; Soudi, A.; Hojati-Talemi, P. J. Inorg. Organomet. Polym. Mater. 2012, 22, 903-909. (p) Wang, Y.; Okamura, T.-a.; Sun, W.-Y.; Ueyama, N. Cryst. Growth Des. 2008, 8, 802-804. (q) Weberski, M. P., Jr.; McLauchlan, C. C. Synthesis and characterization of a tripodal vanadium(III) phosphonate cluster; American Chemical Society: Washington, DC, 2006; p GLRM-096. (r) Xu, X.; Nieuwenhuyzen, M.; Zhang, J.; James, S. L. J. Inorg. Organomet. Polym. Mater. 2006, 15, 431-437. (s) Xue, Z.-Z.; Sheng, T.-L.; Zhu, Q.-L.; Yuan, D.-Q.; Wang, Y.-L.; Tan, C.-H.; Hu, S.-M.; Wen, Y.-H.; Wang, Y.; Fu, R.-B.; Wu, X.-T. CrystEngComm 2013, 15, 8139-8145. (t) Zhang, L.-Y.; Liu, Y.; Li, K.; Pan, M.; Yan, C.; Wei, S.-C.; Chen, Y.-X.; Su, C.-Y. CrystEngComm 2013, 15, 7106-7112. (u) Fujita, M.; Nagao, S.; Ogura, K. J. Am. Chem. Soc. 1995, 117, 1649-1650. (v) Su, C.-Y.; Cai, Y.-P.; Chen, C.-L.; Lissner, F.; Kang, B.-S.; Kaim, W. Angew. Chem., Int. Ed. 2002, 41, 3371-3375.

(3) (a) Tabellion, F. M.; Seidel, S. R.; Arif, A. M.; Stang, P. J. Angew. Chem., Int. Ed. 2001, 40, 1529-1532. (b) Zaman, M. B.; Udachin, K. A.; Ripmeester, J. A. CrystEngComm 2002, 4, 613-617. (c) Shin, D. M.; Lee, I. S.; Chung, Y. K. Inorg. Chem. 2003, 42, 8838-8846. (d) Noveron, J. C.; Chatterjee, B.; Arif, A. M.; Stang, P. J. J. Phys. Org. Chem. 2003, 16, 420-425. (e) Sporer, C.; Wurst, K.; Ruiz-Molina, D.; Kopacka, H.; Veciana, J.; Jaitner, P. J. Organomet. Chem. 2003, 684, 44-49. (f) Granifo, J.; Garland, M. T.; Baggio, R. Inorg. Chem. Commun. 2005, 8, 568-573. (g) Horikoshi, R.; Mochida, T.; Kurihara, M.; Mikuriya, M. Cryst. Growth Des. 2005, 5, 243-249. (h) Nunokawa, K.; Onaka, S.; Mizuno, Y.; Okazaki, K.; Sunahara, T.; Ito, M.; Yaguchi, M.; Imai, H.; Inoue, K.; Ozeki, T.; Chiba, H.; Yosida, T. J. Organomet. Chem. 2005, 690, 48-56. (i) Zaman, M. B.; Udachin, K.; Ripmeester, J. A.; Smith, M. D.; Zur, L. H.-C. Inorg. Chem. 2005, 44, 5047-5059. (j) Winter, S.; Weber, E.; Eriksson, L.; Csoeregh, I. New J. Chem. 2006, 30, 1808-1819. (k) Aakeroy, C. B.; Schultheiss, N.; Desper, J.; Moore, C. CrystEngComm 2007, 9, 421-426. (1) German-Acacio, J. M.; Hernandez-Ortega, S.; Aakeroey, C. B.; Valdes-Martinez, J. Inorg. Chim. Acta 2009, 362, 4087-4090. (m) Romanenko, G. V.; Maryunina, K. Y.; Bogomyakov, A. S.; Sagdeev, R. Z.; Ovcharenko, V. I. Inorg. Chem. 2011, 50, 6597-6609. (n) Li, L.; Turnbull, M. M.; Ackers, J.; Chen, J.; Lin, H.; Pan, B.; Wang, H.; Foxman, B. M. Inorg. Chim. Acta 2009, 362, 3845-3852. (o) Zhang, Y.-J.; Wang, J.-J.; Chen, J. Z. Anorg. Allg. Chem. 2012, 638, 1849-1854. (4) Tabellion, F. M.; Seidel, S. R.; Arif, A. M.; Stang, P. J. J. Am. Chem. Soc. 2001, 123, 7740-7741.

(5) Birk, T.; Pedersen, K. S.; Thuesen, C. A.; Weyhermuller, T.; Schau-Magnussen, M.; Piligkos, S.; Weihe, H.; Mossin, S.; Evangelisti, M.; Bendix, J. *Inorg. Chem.* **2012**, *51*, 5435–5443.

(6) The volume of Cu(hfac)₂ was calculated by: *Spartan '10;* Wavefunction, Inc.: Irvine, CA, 2010.

(7) Spek, A. L. Acta Crystallogr., Sect. D 2009, 65, 148-155.

(8) (a) Clegg, J. K.; Lindoy, L. F.; Moubaraki, B.; Murray, K. S.; McMurtrie, J. C. *Dalton Trans.* **2004**, 2417–2423. (b) Clegg, J. K.; Li, F.; Jolliffe, K. A.; Meehan, G. V.; Lindoy, L. F. *Chem. Commun.* **2011**, 47, 6042–6044.

(9) The total volume of two THF molecules was calculated using the X-ray data.

(10) Ananchenko, G. S.; Udachin, K. A.; Pojarova, M.; Dubes, A.; Ripmeester, J. A.; Jebors, S.; Coleman, A. W. *Cryst. Growth Des.* **2006**, *6*, 2141–2148.

(11) Gans, P.; Sabatini, A.; Vacca, A. Talanta 1996, 43, 1739-1753.

(12) Walker, W. R.; Li, N. C. J. Inorg. Nucl. Chem. 1965, 27, 2255-2261.