

A NEW METHOD FOR THE SYNTHESIS OF 3-DEOXY SUGARS BY GRIGNARD REACTIONS WITH
METHYL 5,6-O-CYCLOHEXYLIDENE-3-O-MESYL- α - AND - β -D-ALLOFURANOSIDE¹

Masajiro KAWANA and Sakae EMOTO
The Institute of Physical and Chemical Research
Wako-shi, Saitama 351

The 3-deoxy sugars were easily prepared in good yield by the reaction of *t*-butylmagnesium bromide with the title compounds. The same reaction with 1-O-benzoyl- or -methyl-*cis*-cyclododecane-1,2-diol gave cyclododecanone.


In a recent communication,² we reported that the reactions of methyl 5,6-O-cyclohexylidene-3-O-mesyl- β -D-allofuranoside 1a and its α -anomer 1b with an excess of methylmagnesium iodide gave, by one step, branched-chain deoxy sugars, methyl 5,6-O-cyclohexylidene-3-deoxy-2-C-methyl- β -D-arabino- and - α -D-ribo-hexofuranoside, respectively. We have now extended the scope of these reactions to the synthesis of deoxy sugars.

A solid of 1a (2 mmol) was gradually added to a solution of *t*-butylmagnesium bromide (ca. 24 mmol) in a mixture of ether (20 ml) and benzene (10 ml) at room temperature over a period of 30 min, and the mixture was stirred for another 30 min. The usual workup exclusively afforded methyl 5,6-O-cyclohexylidene-3-deoxy- β -D-arabino-hexofuranoside 2a [syrup, $[\alpha]_D^{25}$ -86.0° (c 1, CHCl₃)]³ in 87% yield after column chromatography. Treatment of 2a with benzoyl chloride in pyridine gave a crystalline benzoate 2b [mp 86.0-86.5°, $[\alpha]_D^{25}$ -88.4° (c 1.2, CHCl₃)].

In a similar manner, the anomer 1b was reacted with *t*-butylmagnesium bromide at 60-65° (bath temperature) for 40 min to give methyl 5,6-O-cyclohexylidene-3-deoxy- α -D-ribo-hexofuranoside 3a [syrup, $[\alpha]_D^{25}$ +93.6° (c 1, CHCl₃)] in 75% yield. Its benzoate 3b [syrup, $[\alpha]_D^{25}$ +96.0° (c 1, CHCl₃)] was obtained by the conventional method.

Both structures of 2a and 3a were determined by means of their elemental analyses, spectroscopic data, and the following chemical conversions. A mixture of 2a (0.4 mmol) and cyclohexanone (0.3 ml) in benzene (1.5 ml) was stirred at room temperature in the presence of a small amount of sulfuric acid for 4 hr. After neutralization with calcium hydroxide, the excess cyclohexanone was reduced with sodium borohydride to give 1,2;5,6-di-O-cyclohexylidene-3-deoxy- β -D-arabino-hexofuranose 4 [syrup, $[\alpha]_D^{24}$ +6.1° (c 1.4, CHCl₃)]. Similarly, 3a gave 1,2;5,6-di-O-cyclohexylidene-

3-deoxy- α -D-ribo-hexofuranose 5 [syrup, $[\alpha]_D^{26} +1.4^\circ$ (c 1.2, CHCl_3)]. Although the furanose structures of 4 and 5 were tentatively assigned, the physical data of these compounds were identical with those of the corresponding samples derived from known methyl 4,6-O-benzylidene-3-deoxy- α -D-arabino-⁴ and -ribo-hexopyranoside,⁵ respectively.



The present reactions would involve the formation of 3-deoxy-2-keto sugars via the 1,2-hydride shift² from C-2 to C-3, followed by the stereoselective reductions of these ketones with *t*-butylmagnesium bromide. We have not succeeded in the

isolation of the keto-sugars, but 1,2;5,6-di-O-cyclohexylidene- α -D-ribo-hexofuranos-3-ulose⁶ was found to be reduced stereoselectively with *t*-butylmagnesium bromide at room temperature to give 1,2;5,6-di-O-cyclohexylidene- α -D-allofuranose⁶ in good yield. When a simple mesylate, 1-O-mesyl-cis-cyclododecane-1,2-diol 6a [mp 97-98°] was treated with a large excess of *t*-butylmagnesium bromide, cyclododecanone 7 was formed in 82% yield. A similar result could be obtained by the use of 1-O-benzoyl-cis-cyclododecane-1,2-diol 6b [mp 112.5-113.5°] in place of 6a.⁷

REFERENCES AND NOTES

1. A part of this work was presented at the 35th National Meeting of the Chemical Society of Japan, Sapporo, October 1976 and supported by a Research Grant for studies of Life Sciences by this Institute.
2. M. Kawana and S. Emoto, *Tetrahedron Lett.*, **1975**, 3395.
3. Satisfactory elemental analyses and spectral data were obtained for all new compounds.
4. D. A. Prins, *J. Am. Chem. Soc.*, **70**, 3955 (1948).
5. E. Vis and P. Karrer, *Helv. Chim. Acta*, **37**, 378 (1954).
6. K. James, A. R. Tatchell, and P. K. Ray, *J. Chem. Soc. (C)*, **1967**, 2681; M. Kawana, H. Ohru, and S. Emoto, *Bull. Chem. Soc. Jpn.*, **41**, 2199 (1968).
7. M. E. Jung, *Tetrahedron*, **32**, 3 (1976): He has quoted a private communication about a similar Grignard reaction which was found by G. Stork, A. A. Ponaras, and G. A. Garcia.

(Received April 9, 1977)