6 H). The arsonium salt 7 reversibly yields 1.1-dimethyl- λ^5 -arsabenzene 3 on treatment with dimsyl anion in Me₂SO. The proton NMR spectrum illustrated in Figure 1 shows signals for H₃, H₅ as a multiplet centered at δ 6.63, for H₄ as an 8 Hz triplet at 4.43 partially overlapping with 10 Hz doublet at 4.34 for H₂, H₆, while the methyl groups are at δ 1.5. ¹³C NMR (Me₂SO) δ 138.6 (for C₃, C₅), 91.8 (for C₄), 70.3 (for C₂, C₆) 22.4 (for CH₃). Again four ring protons of 7 at H₂, H₄, and H₆ are washed out with D₂O-Me₂SO and base.⁶

In the NMR spectra of 1b and 3 the H_2 , H_4 , and H_6 protons have chemical shift values well outside the aromatic region and are markedly upfield from the range of olefinic protons of their conjugate acids, 5 and 7. On the other hand the H_3 and H_5 protons of 1 and 3 are relatively highly deshielded. However, the signals for these protons occur approximately half a part per million upfield from the corresponding H₃ protons of conjugate acids 5 and 7. The absences of any shift to lower fields for H_3 and H_5 as well as the large upfield shift of H_2 , H_4 , and H₆ suggest that there are no appreciable ring current effects in λ^5 -phosphabenzenes and λ^5 -arsabenzenes.⁷ In fact the NMR spectra, together with proton exchange data, strongly suggest that C_2 , C_4 , and C_6 bear a high electron density. Thus we must view the ring carbon atoms as favoring a pentadienyl anion which is stabilized largely by electrostatic interaction with the positive heteroatom.

The C NMR spectral data on 1 and 3 further support these conclusions.⁷ Signals for C_2 and C_6 occur more than 60 ppm upfield from the signals of C_3 and C_5 .⁸ As in the proton NMR spectra the signal for C_4 is not shifted as far upfield. Very likely this is a consequence of polarization of the carbanionic pentadienyl system by the positive heteroatom.⁹ It is striking that the chemical shift values of ylides 1 and 3 are very nearly identical with those of anions 8 and 9, available from deprotonation of 1 and $3.^{4,10}$ It seems highly likely that the electron density at each carbon is very similar. In contrast to the near identity of chemical shift values of anions and ylides, the ³¹P-¹³C and ³¹P-¹H coupling constants are markedly divergent for 1 and 8,4,11 an expected consequence of the different hybridization at P in the two systems.

¹H and ¹³C NMR spectra of 1,1-dimethyl- λ^5 -phosphabenzene and 1,1-dimethyl- λ^5 -arsabenzene present strong presumptive evidence that these compounds are ylidic in character. In this respect they strongly resemble the thiabenzenes.^{9,12,13} Several bonding proposals have been put forward for λ^5 -phosphabenzenes¹³ and similar cyclic systems^{15,16,17} conjugated via d-p π -bonding to higher row elements. Our findings do not rule out the use of the ultimate arsenic and phosphorus d orbitals in bonding of the λ^5 -phosphabenzenes and the λ^5 -arsabenzenes, but they do strongly argue against any bonding proposal in which appreciable electron density is donated from carbon to the heteroatoms.

Acknowledgment. We wish to thank the National Science Foundation for partial support of this work under Grant No. MPS74-17594.

References and Notes

- (1) K. Dimroth, Fortschr. Chem. Forsch., 38, 1 (1973).
- G. Maerki, Lect. Heterocycl., 1, 69 (1972).
 G. Maerki, Angew. Chem., Int. Ed. Engl., 2, 153, 479 (1963); 4, 1023 (1965).
- A. J. Ashe, III, and T. W. Smith, Tetrahedron Lett., submitted.
- The NMR spectra were recorded using either a Varian T60 or JEOL JNM-(5) PFT/100 instrument. Me₄Si was used as an internal standard.
- The H NMR spectrum showed only singlets at δ 7.0 (1 H), 6.2 (1 H), and 2.2 (6)
- (6) The matrix spectrum showed only singlets at δ 4 for HOD. (7) For comparison see data for λ^3 -phosphabenzene and λ^3 -arsabenzene: A. J. Ashe, III, *J. Am. Chem. Soc.*, **93**, 3293 (1971); A. J. Ashe, III, R. R. Sharp, and J. W. Tolan, *ibid.*, **98**, 5451 (1976).
- Data for 2 are similar: T. Bundgaard, H. L. Jakobsen, K. Dimroth, and H. H. Pohl, *Tetrahedron Lett.*, 3179 (1974).
- (9) See A. G. Hortmann and R. L. Harris, J. Am. Chem. Soc., 93, 2471 (1971).

- (10) The ¹³C NMR spectrum of **8** consists of doublets at δ 71.8 (3 Hz) for C₂, 133.0 (8 Hz) for C₃, and 96.5 (9 Hz) for C₄, while signals for **9** occur at δ
- 103.0 (e1) for C_2 , and S_3 , and S_2 . (e) F_{L2} for C_4 , while signals for S occur at σ 72.9 for C_2 , 132.1 for C_3 , and 92.6 for C_4 . See ref 4. (11) For 8, $J_{PH}^2 = 53$ Hz while $J_{PH}^3 \sim 8$ Hz. (12) A. G. Hortmann, R. L. Harris, and J. A. Miles, *J. Am. Chem. Soc.*, 96, 6119 (1974); A. G. Hortmann and R. L. Harris, *ibid.*, 92, 1803 (1970).
- B. E. Maryanoff, J. Stackhouse, G. H. Senkler, Jr., and K. Mislow, J. Am. Chem. Soc., 97, 2718 (1975).
- (14) H. Oehling and A. Schweig, *Tetrahedron Lett.*, 4941 (1970).
 (15) D. P. Craig, M. L. Hoffman, R. Mason, and N. L. Paddock, *J. Chem. Soc.*, 1376 (1961).
- (16) C. A. Coulson, *Nature (London)*, **221**, 1106 (1969).
 (17) M. J. S. Dewar, "The Molecular Orbital Theory of Organic Chemistry", McGraw-Hill, New York, N.Y., 1969, pp 430–436.
- (18) A. P. Sloan Fellow 1972-1976.

Arthur J. Ashe, III,* 18 Timothy W. Smith

Department of Chemistry, The University of Michigan Ann Arbor, Michigan 48109 Received July 6, 1976

Ligand Effects and Product Distributions in **Molybdothiol Catalyst Systems**

Sir:

The reduction of acetylene to ethylene is one of the reactions catalyzed by nitrogenase and is commonly used to assay for the activity of this enzyme.¹ The complex, $Mo_2O_4(cys)_2^{2-}$, 1 (X = Y = H),² is reported to mimic the enzyme in this re-



spect,³⁻⁶ catalyzing the formation of ethylene (but with small amounts of ethane), the reducing power being supplied either by NaBH₄^{3,4} or electrolytically.⁵ Although complexes of molybdenum seem superior as catalysts to those of other transition metals,⁶ structural variations in the ligand have not been systematically pursued. We have therefore initiated a program of preparation and testing of Mo complexes containing cysteine-related ligands as potential catalysts. These studies reveal several rather striking results. Even with the cysteine complex, we find the major reduction product of acetylene in borate buffer to be 1,3-butadiene (C_4H_6), not ethylene. The production of C_4H_6 presumably went unnoticed in all previous investigations. Further, we observe that seemingly minor changes in ligand or buffer have profound effects on the product distribution and/or catalytic activity.

Typical reactions were run in 15-ml "Hypovials" (Pierce Chemical Co.) as follows: Septum-sealed vials containing 25 mg of complex and 3.5 ml of pH 9.6 buffer (0.1 M Na₂B₄O₇. $10H_2O$ or 0.2 M Na₂CO₃) were flushed with acetylene and kept at 30 °C for 4-6 h⁷ before initiating the reaction by injection of 0.10 ml of 2 M NaBH₄ in DMF.⁸ Reaction mixtures were then shaken⁹ (30 °C bath) and the gas phase analyzed¹⁰ by GC on Porapak N (temperature programming). The results are summarized in Table I.

Butadiene (C_4H_6) was found to be a major product for those complexes in Table I which were catalytically active in the reduction of acetylene. A preparative scale reaction using 1 (X = Y = H) yielded sufficient C_4H_6 for characterization by methods other than GC. The product was separated by trapping at -50 °C and confirmed to be butadiene by its NMR and mass spectra. Its formation constitutes an overall two-electron reductive dimerization of acetylene.

Our first attempts at ligand variation involved methyl sub-

Table I. Acetylene Reduction by Molybdocomplex Systems^a

	Gas phase (µmol)			
Complex (ligand)	$\overline{C_2H_4}$	C_2H_6	C_4H_6	C_2H_2
1 (X = Y = H) (cys)	52.5	0.6	142	362
$1 (X = Y = H) (cys)^{b}$	130	4.8	27	450
$1 (X = Y = CH_3) (pen)$	4.0	1.1	3.7	668
1, $X = CH_3$; $Y = H$ (<i>threo-β-Me</i>)	52.9	1.3	149	345
1, X = H; Y = CH ₃ (erythro- β -Me)	8.0	1.3	10.0	672
2 (edta)	4.9	0.9	0.6	672
3 (edcys)	0.6	0.3	0.1	668

^a 25-min reaction times and borate buffer (pH 9.6) were used unless noted otherwise. At the catalyst level employed, ca. 80 μ mol of reduction products correspond to one turn-over per Mo *atom*. That no other major product has escaped detection is clear from consideration of material balance. Control experiments lacking Mo or ligand yield less than 0.14 μ mol of reduced products. ^b 0.2 M carbonate buffer, pH 9.6. A similar effect is seen using the threo complex.

stitution of cysteine at the β position. The complexes, prepared by standard methods,¹¹ are extremely similar to Mo₂O₄-(cys)₂²⁻ in properties and are presumed to have structure **1**. Remarkably, the complex Mo₂O₄(pen)₂²⁻, **1** (X = Y = CH₃),¹² was found ineffective in catalyzing acetylene reduction. The presence of the two methyl groups in the β position has clearly compromised the reactivity of the complex.

To probe the nature of this effect, we prepared both the threo and erythro forms of β -methylcysteine¹³ and their corresponding $Mo_2O_4L_2^{2-}$ complexes. These isomeric complexes have identical electronic and very similar infrared spectra. Remarkably, the three complex (1) $(X = CH_3, Y = H)$ behaves quite like the cysteine complex in the reduction of acetylene (in both product yield and distribution) while the erythro complex (1) ($X = H, Y = CH_3$) resembles the penicillamine complex and has little activity. Because the two isomeric β -methylcysteine ligands are likely to have similar electronic influences in these complexes, a steric factor must be responsible for the observed phenomenon. Inspection of molecular models reveals a probable explanation, if dissociation of carboxylate is assumed necessary for activity. To produce an accessible metal site by carboxylate dissociation requires a twist of the C-C bond in the five-membered MoNCCS chelate ring. In the case of the inactive erythro complex, this twist is accompanied by eclipsing of the methyl and carboxylate groups, a 1-2 interaction which may be sufficiently unfavorable to hinder the dissociation process. In the active three isomer eclipsing of these groups does not occur and the dissociation of carboxylate should be energetically more favorable. Moreover, even if the carboxylate dissociation does occur, the β -methyl group of the erythro form can partially block the position trans to Mo-O_t, while in the threo form there is a clearly accessible metal site. In either event the open site in the threo form may be more accessible to nucleophilic attack, which could aid in the dissociation of the dimer to catalytically active monomers (see below). A specific buffer ion could be the nucleophile in question, for, as shown in Table I, the product distribution is profoundly affected by changing the buffer ion from borate to carbonate. The true nature of this effect awaits elucidation through additional experiments.

One of the hypotheses resulting from earlier ligand variation studies is that sulfur-donor ligands are required for activity.⁶ The lack of activity of $Mo_2O_4(edta)^{2-}$, **2**, is thus ascribed to absence of thiolate. Moreover, detailed electrochemical studies reveal that while the cysteine complex dissociates on reduction,

the dinuclear edta complex remains intact.^{14,15} It is possible, however, that the ethylenediamine (en) linkage bridging the two Mo atoms in the edta complex may be in part responsible for the lack of activity. To test this idea we prepared the N,N'-ethylenedicysteine¹⁶ complex, Mo₂O₄(edcys)²⁻, 3 (of



proposed structure¹² shown), wherein an ethylene bridge links the cysteine units on each Mo atom. We find that this complex resembles the edta complex in both electrochemistry and lack of activity in C_2H_2 reduction (Table I). It would appear that despite the presence of thiol ligands, the bridging en group does not allow dissociation into the discrete monomeric units apparently needed for activity. Furthermore, the little activity that is present leads to the formation of a substantial proportion of C_2H_6 , indicating the ability of the reduced dinuclear complex to effect the required four-electron reduction.

Finally, in view of our finding of extensive 1,3-butadiene formation in the "model" system, one must ask the question: does nitrogenase produce C_4H_6 from C_2H_2 ? It is notable that at high levels of acetylene, the reduction of C_2H_2 by nitrogenase shows substrate inhibition (at least insofar as production of ethylene is concerned).¹⁷ The possibility of nitrogenase catalyzing the production of C_4H_6 from C_2H_2 was therefore investigated. Using 1 atm of C_2H_2 and conditions where the Mo-Fe protein has a specific activity of 1600 nmol of C_2H_4 produced (mg of enzyme)⁻¹ min^{-1 18} there was no detectable butadiene (i.e., <0.3 nmol min⁻¹ (mg of enzyme)⁻¹). Therefore, it is clear that the molybdothiol model system does not mimic nitrogenase in this important respect.

The results presented in this communication illustrate both pitfalls and promise in the model approach. They emphasize the need to look at the chemistry of a system as a whole and not solely at those aspects which happen to mimic the enzyme. Further, the observation that subtle variations of the ligand in the molybdothiol complex have large effects on catalytic activity opens the possibility that the optimal activity for such systems has not yet been achieved.

Acknowledgment. The authors are grateful for the able technical assistance of Dale Work in ligand synthesis and Deloria Jacobs in the nitrogenase experiments.

References and Notes

- R. Schöllhorn and R. H. Burris, *Proc. Natl. Acad. Sci. U.S.A.*, **58**, 213 (1967);
 M. J. Dilworth, *Biochim. Biophys. Acta*, **127**, 285 (1966);
 R. W. F. Hardy,
 R. D. Holsten, E. K. Jackson, and R. C. Burns, *Plant Physiol.*, **43**, 1185 (1968).
- (2) A. Kay and P. C. H. Mitchell, *Nature (London)*, **219**, 267 (1968); L. R. Melby, *Inorg. Chem.*, **8**, 349 (1969); J. R. Knox and C. K. Prout, *Acta Crystallogr. Sect. B*, **25**, 1857 (1969).
- (3) G. N. Schrauzer and G. Schlesinger, J. Am. Chem. Soc., 92, 1808 (1970).
- (4) G. N. Schrauzer and P. Doemeny, J. Am. Chem. Soc., 93, 1608 (1971).
- D. A. Ledwith and F. A. Schultz, J. Am. Chem. Soc., 97, 6591 (1975).
 G. N. Schrauzer in Proceedings of the 1st International Symposium on Nitrogen Fixation, Vol. 1, W. E. Newton and C. J. Nyman, Ed., Washington State University Press, Pullman, Wash., 1976, p 79. Note Added in Proof. Personal communications from Professors F. A. Schultz and G. N. Schrauzer indicate that they too can observe butadiene formation (in varying amounts) from acetylene in their Mo-cysteine systems.
- (7) In the catalytic systems the "activity" increases substantially over 4–5 h. For those data in Table I, a 5-h "incubation" period was used. An effect of incubation time on activity has been previously noted and attributed to dissociation.⁴ Yields of products in the noncatalytic systems were relatively independent of this effect.

- (8) The use of DMF had no effect on the product yields.
- (9) Without shaking, the cysteine complex yielded C₂H₄ and C₄H₆ (borate buffer) in about equal amounts, probably due to depletion of acetylene in the liquid phase.
- (10) A 200-µl sample was taken using a valve-type syringe ("Pressure-Lok"). Thus an aliquot (0.0114) of the gas phase could be obtained regardless of pressure changes in the vial.
- (11) E. I. Stiefel, Prog. Inorg. Chem., in press.
- (12) Preparation of the new complexes will be described elsewhere. Satisfactory C, H, N analyses were obtained in all cases. Infrared, NMR, electronic spectra, and conductivity behavior in comparison with $Mo_2O_4(edta)^2^-$ and $Mo_2O_4(cys)_2^{2-}$ support the structures shown. Abbreviation: pen = peniciliamine = 3-mercaptovaline = $\beta_i\beta$ -dimethylcysteine.
- (13) H. E. Carter, C. M. Stevens, and L. F. Ney, J. Biol. Chem., 139, 247 (1941). The configurational assignments have been recently established: J. Hoogmartens, P. J. Claes, and H. Vanderhaeghe, J. Org. Chem., 39, 425 (1974).
- (14) V. R. Ott and F. A. Schultz, J. Electroanal. Chem., 59, 47 (1975).
- V. R. Ott and F. A. Schultz, J. Electroanal. Chem., 61, 81 (1975).
 P. Biondeau, C. Berse, and D. Gravel, Can. J. Chem., 45, 49 (1967). We are indebted to Dr. Harvey Schugar of Rutgers University for suggesting and initially supplying this ligand and Dr. C. T. C. Creedy for helpful discussions on this point.
- (17) J. C. Hwang and R. H. Burris, *Biochim. Biophys. Acta*, 283, 339 (1972).
 (18) The nitrogenase preparation used was from *Azotobacter vinelandii*, the specific activity being determined in the presence of excess Fe protein and 10% C₂H₂ in argon. The value reported is for the reaction mixture containing S₂O₄²⁻⁷, ATP, creatine phosphate, creatine phosphokinase, and Mg²⁺ as well as the enzyme. Details of enzyme preparation and purity will be reported elsewhere by E. I. Stiefel and D. Jacobs.

James L. Corbin,* Narayanankutty Pariyadath Edward I. Stiefel*

Contribution No. 579 Charles F. Kettering Research Laboratory Yellow Springs, Ohio 45387 Received July 13, 1976

Novel Thioxo- β -lactams from Thiosulfinates Derived from Penicillin Sulfoxides

Sir:

Recent publications¹ revealed the utilization of the trimethylsilyl (Me₃Si) group in the isolation and crystallization of β -lactam sulfenic acids II and III (R₂ = H, R₁ = CH₃, p-NO₂C₆H₄CH₂) generated thermally from penicillin sulfoxides. The Me₃Si group distinguished itself from other acid protecting groups in that it is hydrolyzed readily to permit the generation of the β -lactam sulfenic acids in situ and, therefore, provided a way of studying the chemistry of this reactive species.

The nucleophilic nature of the sulfur atom in the sulfenic acid is well known;² thus, sulfenic acid II ($R_2 = H$) obtained from hydrolysis of the Me₃Si ester (II, $R_2 = Me_3Si$) will add to the vinylic double bond under neutral conditions to regenerate the penicillin sulfoxides. However, due to the lack of reactivity of the conjugated double bond of the α,β -unsaturated ester (III) to the intramolecular 1,4 addition,³ a characteristic intermolecular condensation reaction of the sulfenic acid moiety is allowed to take place.⁴ Therefore, when sulfenic acid III ($R_1 = CH_3$, $R_2 = H$), generated from its Me₃Si ester (III, $R_1 = CH_3$, $R_2 = Me_3Si$) by the treatment of methanol, was allowed to stand at room temperature for several hours, this condensation reaction occurred readily and the resulting thiosulfinate crystallized from chloroform-ether to give a colorless crystalline solid IV ($R_1 = CH_3$): mp 156–157 °C; $[\alpha]^{27}_{D}$ +56° (CHCl₃); IR (CHCl₃) 1795, 1785, 1735, and 1115 cm⁻¹; NMR (CDCl₃) δ 2.08 (s, 3 H), 2.15 (s, 3 H), 2.21 (s, 3 H), 2.35 (s, 3 H), 3.76 (s, 3 H), 3.81 (s, 3 H), 5.62 (d, 2 H, J = 4.5 Hz), 5.70 (ABq, 2 H, J = 4.5 and 34 Hz) and 7.82 (s, 8 H). Similarly, treatment of III ($R_1 = CH_3$, p-NO₂C₆H₄- CH_2 , Me_3Si ; $R_2 = Me_3Si$) in chloroform with methanolic HCl resulted in the instantaneous precipitation of IV ($R_1 = CH_3$, p-NO₂C₆H₄CH₂, H, respectively) in high yield.

The thiosulfinate IV $(R_1 = CH_3)$, after brief heating or prolonged standing at room temperature in an inert solvent,



produced the novel thioxo- β -lactam V (R₁ = CH₃).⁵ This monocyclic thione lactam will not polymerize and can be crystallized from methanol as light yellow prisms, mp 126-127 °C, in 67% yield.^{6.7} The presence of an azetidine carbonyl group and a thiocarbonyl carbon in V $(R_1 = CH_3)$ was indicated respectively by an absorption in the IR (CHCl₃) at 1835 cm^{-1} and by a chemical shift at 200.5 ppm downfield from tetramethylsilane in the ¹³C NMR (CDCl₃).⁸ In addition, the 13 C NMR also indicated the presence of 2CH₃ (δ 22.0, 23.4), 1OCH₃ (δ 52.3), 1CH (δ 65.7), 2C=C (δ 117.1, 161.8), 1 NC=O (δ 165.9), 1 ester C (δ 167.2), and a phthaloyl group. A ¹H coupled ¹³C spectrum verified that the thiocarbonyl carbon is within three bonds of only one proton. The proton NMR (CDCl₃) of V had signals at δ 2.18 (s, 3 H), 2.42 (s, 3 H), 3.78 (s, 3 H), 5.92 (s, 1 H), and 7.84 (m, 4 H), and was also consistent with the structure.⁹ Using the same method, the thioxo- β -lactams (V, R₁ = H, p-NO₂C₆H₄CH₂) can be prepared from the respective thiosulfinates (IV, $R_1 = H$, p- $NO_2C_6H_4CH_2$).

We theorized that the corresponding thioxo- β -lactam (VI) in the β , γ -unsaturated ester series could be prepared if the sulfenic acid II could be intercepted by an external sulfur nucleophile. Treatment of I (R₁ = CH₃) with *n*-pentyl mercapton afforded a high yield of VII, NMR (CDCl₃) δ 5.74 (ABq, 2 H, J = 4.5 and 10 Hz), 5.10–5.35 (m, 3 H, characteristic of the vinylic CH₂ and the allylic methine H), 2.32–2.80 (m, 2 H, -SCH₂-) and 2.12 (s, 3 H).¹⁰ Subsequent oxidation of VII with *m*-chloroperbenzoic acid-CH₂Cl₂ gave selectively the thiosulfinate VIII, NMR (CDCl₃) δ 5.8–6.4 (m, 2 H), 2.8–3.2 (m, 2H, OSCH₂-), and 2.0 (s, 3 H).¹¹ Heating VIII in re-