

JOM 23926PC

Preliminary Communication



1,8-Bis(diphenylphosphino)naphthalene:
a rigid chelating, diphosphine analogue
of proton spongeRichard D. Jackson, Stuart James, A. Guy Orpen
and Paul G. PringleSchool of Chemistry, University of Bristol, Cantock's Close,
Bristol BS8 1TS (UK)

(Received April 2, 1993)

Abstract

The synthesis of the new diphosphine, 1,8-bis(diphenylphosphino)naphthalene (**1a**), and its X-ray crystal structure are reported. Protonation of **1a** gives a fluxional species to which a P–P bonded structure is assigned. Despite the strain apparent in the solid state due to the proximity of the diphenylphosphino groups, it appears that **1a** has a normal coordination chemistry with platinum(II) and palladium(II).

There is currently great interest in new diphosphine ligands with a C_3 -backbone because the six-membered palladium(II) chelates which they form have been demonstrated to be extremely active catalysts for CO/ethylene co-polymerization [1]. We report here the new C_3 -diphosphine ligand 1,8-bis(diphenylphosphino)naphthalene (**1a**), a phosphorus analogue of 1,8-bis(dimethylamino)naphthalene (**2**), which is also known as "proton sponge" because it is an effective non-nucleophilic base [2]. It might have been expected that diphosphine analogues of **2** would be confined to those having small substituents, and 1,8-bis(dimethylphosphino)naphthalene (**1b**) is known [3], but we have found that the bulky 1,8-bis(diphenylphosphino)naphthalene is surprisingly easily made and behaves as a normal chelating diphosphine towards palladium(II) and platinum(II).

**1a** R = Ph**1b** R = Me**1c** R = Cy**2**


Treatment of 1,8-dilithionaphthalene [3] with Ph_2PCl in diethyl ether at $0^\circ C$ gave an 82% yield of diphosphine **1a** as a yellow, air-stable powder which has been fully characterised [4*]. It is readily made in this way on a 2 g scale. Clear yellow crystals of **1a** suitable for X-ray crystallography were grown from CH_2Cl_2/Et_2O . The crystal structure analysis [5*] confirms the identity of **1a** (see Fig. 1) and provides information on the distortions present in the molecule. In common with other 1,8-substituted naphthalenes, **1a** exhibits both out-of-plane and in-plane bending of the PPh_2 substituents away from one another as well as buckling of the C_{10} unit (mean carbon atom deviation from least-squares C_{10} plane $0.006 \text{ \AA}$, phosphorus deviations: P(1) $0.50 \text{ \AA}$, P(2) $0.34 \text{ \AA}$). While these distortions lower the symmetry of the $C_{10}P_2$ nucleus from C_{2v} to approximately C_2 , the orientations of the PPh_2 groups are such as to reduce the molecular symmetry to C_1 in the solid state. The net effect of these distortions is to generate a P(1) $\cdots$ P(2) distance of $3.052 \text{ \AA}$, well within the sum of the van der Waals radii ($3.80 \text{ \AA}$). This distance is comparable with those in Pt^{II} or Pd^{II} complexes of conventional chelating diphosphines (e.g. P $\cdots$ P in $[PdCl_2(Ph_2PCH_2CH_2CH_2PPh)_2]$ is $3.193 \text{ \AA}$ [6]), and may explain the relatively orthodox coordinating properties of **1a** (see below).

The inequivalence of the two P atoms in **1a** in the solid state (see above) is confirmed by the MAS solid state $^{31}P\{^1H\}$ NMR spectrum which shows an AB pattern with $\delta(P_A) -18.0$ and $\delta(P_B) -10.2$. The $J(PP)$ value of 199 Hz is unusually large for a $^4J(PP)$ coupling and indicates that there is a significant through-space component. In CD_2Cl_2 solution the $^{31}P\{^1H\}$ NMR spectrum of **1a** is a sharp singlet at -14.7 ppm even at $-80^\circ C$ indicating that either the structure is symmetrical in solution or there is rapid fluxionality on the NMR timescale leading to equivalence of the P atoms (the solution chemical shift is close to the average of the solid state shifts).


Addition of MeI to **1a** gave the monophosphonium salt **3**, the $^{31}P\{^1H\}$ NMR spectrum of which at $+28^\circ C$ was an AB pattern with $^4J(PP)$ of 24 Hz . Addition of one equivalent of $HBF_4 \cdot OMe_2$ to diphosphine **1a** in CD_2Cl_2 gave rise to a new species which from variable

* Reference number with asterisk indicates a note in the list of references.

temperature ^{31}P NMR studies was found to be fluxional. At $+28^\circ\text{C}$, the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum is a broad singlet ($w_{1/2} = 35$ Hz) at -4.9 ppm. The proton-coupled ^{31}P NMR spectrum is a doublet with $J(\text{PH}) = 288$ Hz. At -80°C the spectrum is an AB pattern, $\delta(\text{P}_\text{A}) = 8.2$ ppm, $\delta(\text{P}_\text{B}) = -19.6$ ppm, $J(\text{P}_\text{A}\text{P}_\text{B}) = 110$ Hz. The proton-coupled ^{31}P NMR spectrum at -80°C shows that only P_A is coupled to the proton with $J(\text{P}_\text{A}\text{H}) = 576$ Hz. $J(\text{P}_\text{B}\text{H})$ must be less than the linewidth (20 Hz). The fluxionality of **4** is therefore proton exchange between the two phosphorus atoms and this is an intramolecular process since $J(\text{PH})$ at ambient temperatures is half the low temperature $J(\text{P}_\text{A}\text{H})$ value. The coalescence temperature is 275 K, corresponding to a $\Delta G^\ddagger$ of 49.5 kJ mol $^{-1}$ as calculated by the Gutowsky–Holm method [7]. The large $J(\text{PP})$ of 110 Hz for the protonated species contrast with that found for the phosphonium salt **3** and is consistent with the P–P bonded structure **4**, analogous to the recently reported [8] tricyclic compound **5**.



Diphosphine **1a** reacts readily with $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$ or $[\text{PdCl}_2(\text{NCPH}_2)_2]$ to give the chelates **6** and **7**, respectively, in high yields (80–90%). The spectroscopic properties of these complexes are apparently normal. For example the ^{31}P coordination chemical shift (Δ) of 16.1 for **6** when compared with Δ for the monophosphine complex **8** yields a ring contribution (Δ_R) of -7.9 , similar to the values for other six-membered



chelates [9]. Hence, contrary to expectation, ligand **1a** has normal diphosphine coordination properties, and is a member of a potentially large class of ligands since preliminary studies indicate that the even bulkier cyclohexyl analogue, **1c**, is available by a similar route.

Acknowledgements

We thank BP for supporting this work and Johnson-Matthey PLC for a generous loan of platinum metals.

References and notes

- (a) Shell, *Eur. Patent* 121965 (1984); (b) E. Drent, J.A.M. van Broekhoven and M.J. Doyle, *J. Organomet. Chem.*, **417** (1991) 235; (c) G.P.C.M. Dekker, C.J. Elsevier, K. Vrieze and P.W.N.M. van Leeuwen, *Organometallics*, **11** (1992) 1598; (d) G.P.C.M. Dekker, C.J. Elsevier, K. Vrieze, P.W.N.M. van Leeuwen and C.F. Roobeek, *J. Organomet. Chem.*, **430** (1992) 357.
- R.W. Alder, *J. Chem. Soc., Perkin Trans. 1*, (1981) 2840, and refs. therein.
- T. Costa and H. Schmidbaur, *Chem. Ber.*, **115** (1982) 1374.
- All isolated new compounds have been characterised by a combination of elemental analyses, IR, ^1H , ^{31}P and ^{13}C NMR spectroscopy. The ^{31}P NMR data for **3–9** (in CDCl_3) are: **1a**, $\delta(\text{P}) -14.7$; **1c**, $\delta(\text{P}) -16.1$; **3**, $\delta(\text{P}_\text{A}) +23.6$, $\delta(\text{P}_\text{B}) -13.0$, $^2J(\text{P}_\text{A}\text{P}_\text{B})$ 24; **4**, $\delta(\text{P}_\text{A}) +8.2$, $\delta(\text{P}_\text{B}) -19.6$, $^2J(\text{P}_\text{A}\text{P}_\text{B})$ 110; **6**, $\delta(\text{P}) +1.6$, $^1J(\text{PtP})$ 3318; **7**, $\delta(\text{P}) +20.8$; and **8**, $\delta(\text{P}) +9.8$, $^1J(\text{PtP})$ 3656.
- Crystal data for **3**: $\text{C}_{34}\text{H}_{26}\text{P}_2$, $M = 496.5$, triclinic, space group $\text{P}\bar{1}$ (No. 2), $a = 9.493(3)$, $b = 10.030(3)$, $c = 15.295(5)$ Å, $\alpha = 90.45(2)$, $\beta = 93.65(2)$, $\gamma = 117.70(2)^\circ$, $V = 1285.5(6)$ Å 3 , $Z = 2$, $D_\text{x} = 1.28$ g cm $^{-3}$, $\bar{\lambda} = 0.71069$ Å, $\mu(\text{Mo-K}\alpha) = 1.9$ cm $^{-1}$, $F(000) = 520$, $T = 295$ K. Data were collected on a Siemens P3m diffractometer for $4 < 2\theta < 45^\circ$. The structure was solved by direct methods and refined by least-squares to $R = 0.066$ for 2361 unique, observed ($I > \sigma(I)$) absorption corrected intensity data. Table of atom coordinates, bond lengths, bond angles, and thermal parameters have been deposited with the Cambridge Crystallographic Data Centre.
- W.L. Steffen and G.J. Palenik, *Inorg. Chem.*, **15** (1976) 2432.
- H. Gutowsky and C.H. Holm, *J. Chem. Phys.*, **25** (1956) 1228.
- R.W. Alder, C. Ganter, C.J. Harris and A.G. Orpen, *J. Chem. Soc., Chem. Commun.*, (1992) 1172.
- P.E. Garrou, *Chem. Rev.*, **81** (1981) 229.




Fig. 1. Molecular geometry of **1a**, important molecular parameters include: bond lengths, $\text{P}(1)\text{--C}(8)$ 1.844(6), $\text{P}(1)\text{--C}(29)$ 1.828(5), $\text{P}(1)\text{--C}(23)$ 1.837(5), $\text{P}(2)\text{--C}(1)$ 1.848(6), $\text{P}(2)\text{--C}(11)$ 1.839(5), $\text{P}(2)\text{--C}(17)$ 1.839(5) Å; bond angles, $\text{P}(1)\text{--C}(8)\text{--C}(9)$ $123.0(4)$, $\text{P}(2)\text{--C}(1)\text{--C}(9)$ $124.5(3)^\circ$; torsion angles, $\text{C}(1)\text{--C}(9)\text{--C}(8)\text{--P}(1)$ $-13.4(7)$, $\text{C}(8)\text{--C}(9)\text{--C}(1)\text{--P}(2)$ $-7.7(7)^\circ$.