

S0960-894X(96)00150-3

## Asymmetric Synthesis of *cis*-(-)-(2*R*4*S*)-4-(Phosphonomethyl)-2-Piperidinecarboxylic Acid, A Potent NMDA Receptor Antagonist


Jerry W. Skiles,<sup>\*,†</sup> Peter P. Giannousis,<sup>§</sup> and Kevin R. Fales<sup>§,¶</sup>

Ciba, Pharmaceuticals Division, 556 Morris Avenue, Summit, New Jersey 07901


**Abstract:** The asymmetric synthesis of *cis*-(-)-(2*R*4*S*)-4-(phosphonomethyl)-2-piperidinecarboxylic acid (**1**), a potent and specific NMDA receptor antagonist, via an olefin-iminium cyclization is described. Copyright © 1996 Elsevier Science Ltd

The excitatory amino acids,<sup>1</sup> aspartate and glutamate, mediate their actions via at least three classes of receptors which are generally represented by the prototypical antagonist *N*-methyl-D-aspartic acid (NMDA), quisqualic acid (QUIS), and kainic acid (KA). Of these the NMDA receptor has been the most studied.<sup>2</sup> Excess activity at this receptor has deleterious effects on CNS function. Antagonists of the NMDA receptor could thus have potential utility in a number of CNS disorders, most notably in the treatment of epilepsy and the neuronal damage resulting from cerebral ischemia.

One of the most potent selective competitive NMDA receptor antagonists<sup>3,4</sup> is **1**; *cis*-(±)-CGS 19755 (Selfotel) which is presently undergoing extensive clinical evaluation as an antiischemic agent in serious traumatic brain injury (TBI)<sup>5</sup> and stroke. The previously described synthesis<sup>3</sup> of **1** initiates with 4-pyridinecarbinol (**3**) to afford ultimately racemic CGS 19755 via a catalytic ring reduction of the 2-pyridine carboxamide **2** as is shown in the schematic below. In this manner *cis*-(±)-CGS 19755 is obtained as a mixture of enantiomers. The preparation and biological activities of the individual enantiomers of *cis*-(±)-CGS 19755 has not been previously described. Herein we describe for the first time the preparation of both stereoisomers of *cis*-(±)-CGS 19755 and report their biological activities in the NMDA receptor binding assay.




In 1986, Grieco and co-workers demonstrated that a variety of 4-hydroxypiperidines may be formed under aqueous conditions by an iminium ion cyclization of a homoallylic amine intermediate.<sup>6,7</sup> We envisioned that 4-substituted pipercolinic acids could be formed under aqueous conditions described by Grieco using an appropriately substituted homoallylic amine and glyoxylic acid as precursors.<sup>8,9</sup> In order to induce the formation of diastereomers, the homoallylic amines used for the cyclization were prepared from (*S*)-(-)- $\alpha$ -methylbenzylamine and (*R*)-(+)- $\alpha$ -methylbenzylamine.



According to the scheme given below the homoallylic amine **5** was prepared by refluxing a mixture of (*R*)-(+)- $\alpha$ -methylbenzylamine, 4-bromo-1-butene (**4**), and Et<sub>3</sub>N in CH<sub>3</sub>CN. The chiral homoallyl amine **5**, [ $\alpha$ ]<sub>D</sub> = +45.4° (*c* = 1 CHCl<sub>3</sub>), was dissolved in a mixture of CH<sub>3</sub>CN and H<sub>2</sub>O (1:1) and treated with glyoxylic acid at room temperature for several days to provide the diastereomeric lactams *cis*-(2*S*4*R*2'*R*)-**8a** and *cis*-(2*R*4*S*2'*R*)-**8b** in a diastereomeric ratio of 1:2 respectively. It was later found the reaction could be pushed to completion within 4 hours at room temperature if the reaction was conducted in CH<sub>3</sub>CN in the absence of H<sub>2</sub>O and in the presence of 4 Å molecular sieves. Of the several potential mechanisms for this conversion it is most likely that the incipient carbocation **7a** is trapped internally by the carboxylate and rarely, if ever, by H<sub>2</sub>O solvent to afford the lactones **8a** and **8b** directly. Supporting evidence for this mechanism, although not proven, is the fact that we observed no *trans*-

hydroxy acid in the NMR of the crude reaction mixture when run in the mixture of  $\text{CH}_3\text{CN}$  and  $\text{H}_2\text{O}$  (1:1). The diastereomeric lactones **8** were easily separated by a simple recrystallization from *n*-hexane. The lactone *cis*-(2*R*4*S*2'*R*)-**8b**, *vide infra*, was subjected to  $\text{CH}_3\text{OH}$  and anhydrous  $\text{NH}_3$  to afford the alcoholic methyl ester (2*R*4*S*2'*R*)-**9** which *via* a Swern oxidation provided the ketone (2*R*2'*R*)-**10**. The ketone (2*R*2'*R*)-**10** was easily transformed into a 2:1 mixture of *E* and *Z*-phosphonates (2*R*2'*R*)-**11** by treatment with  $[(\text{EtO})_2\text{P}(\text{O})]_2\text{CH}_2$ , DBU, and  $\text{LiCl}$  in  $\text{CH}_3\text{CN}$  or THF.<sup>10</sup> Catalytic reduction and hydrogenolysis (10 %  $\text{Pd/C}$ ,  $\text{H}_2$ ) of *cis*-(2*R*2'*R*)-**11** gave cleanly the triester *cis*-(2*R*4*S*)-**13** which was hydrolyzed with aqueous 6 *N*  $\text{HCl}$  to afford *cis*-(-)-(2*R*4*S*)-**1**, the minus isomer of ( $\pm$ )-CGS 19755. Alternatively, according to the scheme below, **11** could be converted in a stepwise manner to **13** by a reduction - hydrogenolysis procedure (10 %  $\text{Pt/C}$  followed by 10 %  $\text{Pd/C}$ ). In a manner similar to that described above but using the lactone *cis*-(2*S*4*R*2'*R*)-**8a** the *cis*-(+)-(2*S*4*R*)-**1** triacid was obtained which corresponds to the (+)-isomer of ( $\pm$ )-CGS 19755.<sup>11</sup>


Scheme I: Synthesis of *cis*-(-)-(2*R*4*S*)-CGS 19755



**Reagents:** All rotations taken in  $\text{CHCl}_3$  ( $c = 1$ ) with the exception of **1** (1 *N*  $\text{NaOH}$ ); a) (*R*)-(+)- $\alpha$ -methylbenzylamine,  $\text{Et}_3\text{N}$ ,  $\text{CH}_3\text{CN}$ ; b)  $\text{OHCCOOH}$ , 4 Å molecular sieves,  $\text{CH}_3\text{CN}$ ; c)  $\text{NH}_3$ ,  $\text{CH}_3\text{OH}$ ; d)  $(\text{ClCO})_2$ ,  $\text{Et}_3\text{N}$ ,  $\text{DMSO}$ ,  $\text{CH}_2\text{Cl}_2$ ; e)  $[(\text{EtO})_2\text{P}(\text{O})]_2\text{CH}_2$ , DBU,  $\text{LiCl}$ , THF; f) 5 %  $\text{Pt/C}$ ,  $\text{H}_2$  (45 psi),  $\text{EtOH}$ ; g) 10 %  $\text{Pd/C}$ ,  $\text{H}_2$  (45 psi),  $\text{EtOH}$ ; h) 6 *N*  $\text{HCl}$ ; i)  $\text{KHCO}_3$ ,  $\text{H}_2\text{O}$ ,  $\text{EtOH}$ .

Assignment of the absolute stereochemistry of the diastereomeric lactones **8** was done by correlation to the known natural product *cis*-(-)-(2*S*4*R*)-4-hydroxypipercolinic acid (**14b**) which was isolated from *Acacia oswaldii* leaves.<sup>12</sup> Using (*S*)-(-)-( $\alpha$ )-methylbenzylamine and (*R*)-(+)-( $\alpha$ )-methylbenzylamine all four possible isomers of **8** were prepared in an analogous manner to that described above and isolated via crystallization (*n*-hexane) and/or chromatography (silica gel; EtOAc/*n*-hexane, 1:3). Thus (2*R*4*S*2'*S*')-**8a**, *vide infra*, was subjected to hydrogenolysis conditions to afford *cis*-(+)-(2*R*4*S*)-4-hydroxypipercolinic acid (**14a**),  $[\alpha]_D = +20.4^\circ$  (*c* = 1.0 CH<sub>3</sub>OH). In a likewise manner (2*S*4*R*2'*R*')-**8b** was subjected to hydrogenolysis conditions to give *cis*-(-)-(2*S*4*R*)-4-hydroxypipercolinic acid (**14b**). The reported optical rotation for the (2*S*4*R*)-**14b** natural product is  $[\alpha]_D = -17^\circ$  (*c* = 1.0 CH<sub>3</sub>OH).<sup>12</sup> Thus in effect the conversion of the enantiomeric (2*R*4*S*2'*S*') and (2*S*4*R*2'*R*') lactones **8a** and **8b** to the known *cis*-4-hydroxypipercolinic acids **14a** and **14b** and comparison of optical rotations with those given in the literature has allowed assignment of the absolute configurations of all four lactones **8** and as well as the *cis*-acids **1**.<sup>13</sup>

**Scheme 2.** Cofirmation of absolute steeochemistry of **8a** and **8b** lactones.



**Reagents and conditions:** Optical rotations of **8** taken in CH<sub>2</sub>Cl<sub>2</sub>; **14** taken in CH<sub>3</sub>OH; a) 10 % Pd/C (45 psi), EtOAc, H<sub>2</sub>O.

The *cis*-(-)-(2*R*4*S*)-4-(phosphonomethyl)piperidine-2-carboxylic acid (**1**) has an IC<sub>50</sub> of 29 nM in a tritiated CPP receptor binding assay compared to 10,000 nM for *cis*-(+)-(2*S*4*R*)-**1**.<sup>14</sup> This activity is in agreement with *N*-methyl-D-aspartate having an (*R*)-configuration at what corresponds to the 2-position of *cis*-(-)-(2*R*4*S*)-**1**.

**Table 1.** NMDA receptor binding results for enantiomers of *cis*-( $\pm$ )-CGS 19755.

Compound	CGS Number	IC <sub>50</sub> (nM)
<i>cis</i> - ( $\pm$ ) - <b>1</b>	CGS 19755	69
<i>cis</i> - (-) - (2 <i>R</i> 4 <i>S</i> ) - <b>1</b>	CGS 20281	29
<i>cis</i> - (+) - (2 <i>S</i> 4 <i>R</i> ) - <b>1</b>	CGS 20282	10,000

**Acknowledgement:** We wish to thank Dr. M. Sills for evaluation of all reported compounds in the NMDA receptor binding assay.

\* To whom correspondence should be addressed.

† Research Department, Department of Medicinal Chemistry (OA/RA).


§ Process Research & Development.

¶ Current address: Wayne State University, Department of Chemistry, Detroit, Michigan 48202.

## References and Notes

- (a) Cotman, C. W.; Iversen, L. L. *Trends Neurosci.* **1987**, *10*, 263. (b) Watkins, J. C.; Olverman, H. J. *Trends Neurosci.* **1987**, *10*, 265.
- Ornstein, P. L.; Monn, J. A.; Schoepp, D. D. *DN&P* **1994**, *7*, 5.

3. Hutchison, A. J.; Williams, M.; Angst, C.; deJesus, R.; Blanchard, L.; Jackson, R. H.; Wilusz, E. J.; Murphy, D. E.; Bernard, P. S.; Schneider, J.; Campbell, T.; Guida, W.; Sills, M. A. *J. Med. Chem.* **1989**, *32*, 2171.
4. Lehmann, J.; Hutchison, A. J.; McPherson, S. E.; Mondadori, C.; Schmutz, M.; Sinton, C. N. M.; Tsai, C.; Murphy, D. E.; Steel, D. J.; Williams, M.; Cheney, D. L.; Wood, P. L. *J. Pharmacol. Exp. Ther.* **1987**, *240*, 778.
5. For a recent review concerning traumatic brain injury (TBI) see: Nair, S.; Lehmann, J.; Williams, J. M.; Weiss, B. *Drug News & Products* **1993**, *6*, 600.
6. Larsen, S. D.; Grieco, P. A.; Fobare, W. F. *J. Am. Chem. Soc.* **1986**, *108*, 3512.
7. Hays, S. J.; Malone, T. C.; Johnson, G. *J. Org. Chem.* **1991**, *56*, 4084.
8. Recently two excellent reviews have appeared on piperidine compounds: (a) Rubiralta, M.; Giralt, E.; Diez, A. "Piperidine: Structure, Preparation, Reactivity and Synthetic Applications of Piperidine and its Derivatives", Elsevier, New York, N.Y. **1991**. (b) Wang, C. -L.; Wuanola, M. A. *Org. Prepar. Proceed. Intern.* **1992**, *24*, 585.
9. (a) Agami, C.; Couty, F.; Poursoulis, M.; Vaissermann, J. *Tetrahedron*, **1992**, *48*, 431. (b) Teng, T. -F.; Lin, J. -H.; Yang, T. -K. *Heterocycles* **1990**, *31*, 1201. (c) Esch, P. M.; deBoer, R. F.; Hiemstra, H.; Boska, I. M.; Speckamp, W. N. *Tetrahedron* **1991**, *47*, 4063. (d) Evans, D. A.; Thomas, E. W.; Cherpeck, R. E. *J. Am. Chem. Soc.* **1982**, *104*, 3695. (e) Overman, L. E.; Sharp, M. J. *J. Am. Chem. Soc.* **1988**, *110*, 612. (f) Overman, L. E.; Sharp, M. J. *Tetrahedron Lett.* **1988**, *29*, 901. (g) Arnold, H.; Overman, L. E.; Sharp, M. J.; Witsschel, M. C. *Org. Syn.* **1991**, *70*, 111.
10. Conditions other than these (e.g., NaH, *n*-BuLi, LDA, etc. ) failed or gave low yields of inferior quality product. The conditions described by Masamune and Roush (DBU and LiCl in THF or CH<sub>3</sub>CN) consistently afforded **11** in good yield (82 - 90 %) as a mixture of *E* and *Z*-isomers: Blanchette, M. A.; Choy, W.; Davis, J. T.; Essenfield, A. P.; Masamune, S.; Roush, W. R.; Sakai, T. *Tetrahedron Lett.* **1984**, *25*, 2183.
11. All compounds exhibited satisfactory C, H, and N microanalysis and were within 0.4 % of theoretical values. All new compounds exhibited <sup>1</sup>H and <sup>13</sup>C NMR which are consistent with the assigned structures.
12. Clark-Lewis, J. W.; Mortimer, P. I. *J. Chem. Soc.* **1961**, 189.
13. As an additional proof of the absolute stereochemical assignments, the carboxamide diethyl ester of *cis*-(±)-**1** could also be resolved into the respective enantiomers *via* its di-*p*-toluoyl-L-tartrate salts (see **15** below). Hydrolysis of each enantiomer with 6 N HCl afforded both *cis*-(+)-**1** and *cis*-(-)-**1**. Optical rotations as well as NMR, MS, and IR of these resolved materials were identical to those prepared by synthetic means. In addition to correlation with the known natural product (2*S*,4*R*)-**14b** the single crystal X-ray structure of the *cis*-(+)-carboxamide diethyl ester di-*p*-toluoyl-L-tartaric acid salt of **1**, i.e. **15** below, confirmed our absolute stereochemical assignments. Complete listings of atomic coordinates, bond lengths, bond angles, thermal parameters and structure factors for compound **15** have been deposited at the Cambridge Crystallographic Data Centre, University Chemical Laboratory, Lensfield Road, Cambridge, CB2 1EW England.



14. For details of the NMDA receptor binding assay see references 3 and 4.