in the presence and absence of oxygen, (2) uncertainties regarding the nature of axial bases (if any) supplied by the graphite surface, and (3) the competitive inhibition by water shown in the present work. However, the surprisingly high dioxygen affinity of 1 can give new insight into the factors that influence dioxygen binding to metal centers.

Acknowledgment. The support of this research by the CNRS (Centre National de la Recherche Scientifique), France, the National Science Foundation (Grant CHE83-18512), the National Institutes of Health (Grant NIHGM17880-14,15), and NATO (North Atlantic Treaty Organization) is gratefully acknowledged. The Bio-organic, Biomedical Mass Spectrometry Resource (A. L. Burlingame, Director), supported by NIH Division of Research Resources Grant RR 01614, is also gratefully acknowledged.

Registry No. 1, 89906-36-5; 1.O2, 89906-37-6; 1.(N-MeIm), 99665-74-4; 1·(H<sub>2</sub>O), 99665-75-5; O<sub>2</sub>, 7782-44-7.

## Photochemical Carbonylation of Benzene by Iridium(I) and Rhodium(I) Square-Planar Complexes

Amanda J. Kunin and Richard Eisenberg\*

Department of Chemistry, University of Rochester Rochester, New York 14627 Received August 19, 1985

The photochemical behavior of d<sup>8</sup> square-planar complexes represents an area of current interest as evidenced by Ford's recent flash photolysis studies on RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> and its well-known Ir analogue.<sup>1-3</sup> In these studies all intermediates and observed transients are thought to form via initial photodissociation of CO. In the absence of added substrate, the principal chemical reaction following the photolysis is recombination of MCl(PPh<sub>3</sub>)<sub>2</sub> and CO. although for M = Rh, dimerization to  $[MCl(PPh_3)_2]_2$  followed by slower carbonylation to starting complex is also noted. Flash photolysis of the Rh complex with either  $H_2$  or  $C_2H_4$  present gives spectral changes indicative of initial reaction of RhCl(PPh<sub>3</sub>)<sub>2</sub> with substrate, but subsequent and rapid carbonylation leads to no net chemical change for the system. In this paper we report that photolysis of these complexes under long-term continuous irradiation leads to significant and new overall reaction chemistry. This chemistry, which is only observed upon photolysis, involves arene C-H functionalization and, in particular, the carbonylation of benzene to benzaldehyde. This observed C-H bond functionalization contrasts with the known oxidative carbonylation of arenes using  $Pd^{2+}$  in that no prior activation of the benzene ring is needed.<sup>4</sup> We are also able to show that the system is limited by the overall thermodynamics of the arene carbonylation reaction and not by the reactivity of the d<sup>8</sup> photocatalyst system.

The starting point for this work derives from a mechanistic proposal made previously in a report on benzene carbonylation using either  $IrH_3(CO)(dppe)$  (1) or  $IrH(CO)_2(dppe)$  (2) as the photocatalyst complex.<sup>5</sup> These coordinatively saturated complexes undergo exchange reaction (1) thermally or photochemically, presumably through the intermediacy of the four-coordinate species IrH(CO)(dppe) (3). Since benzene carbonylation was only observed upon photolysis, it was proposed that 3 was the photochemically active species for the system. A test of this proposal was performed based on the assumption that 3, as a four-coordinate, 16-e<sup>-</sup> species, would be yellow or orange and therefore absorb light at longer wavelengths than the two precursor complexes 1 and 2. Two samples containing 6 mM solutions of



IrH(CO)<sub>2</sub>(dppe) (2) in benzene under 600 torr of CO were photolyzed through a 366-nm cut-off filter. Under these conditions, 2 is the only observable complex in solution, but it exhibits no significant absorptions at  $\lambda$  greater than 320 nm. The first sample, which was maintained at room temperature, showed no benzaldehyde formation after 48 h of photolysis. The second sample was heated during photolysis to 75 °C at which temperature dissociation of CO from 2 is known to occur.<sup>4</sup> After 6 h of photolysis the second sample was analyzed by gas chromatography and found to contain  $1.1 \times 10^{-3}$  mmol (1.8 mM) of benzaldehyde. This result provides strong support that 3 is the photochemically active species in the previously reported benzene carbonylation system.5

The formation of benzaldehyde is also seen when benzene solutions of the chelated di(phosphine) complexes IrH<sub>3</sub>(CO)(dppp) (4) (dppp = 1,3-bis(diphenylphosphino)propane) and IrBr-(CO)(dppe) (5) are irradiated under 600 torr of CO. Complex 4 is synthesized analogously to the procedure used for 1<sup>5</sup> and, like 1, undergoes thermal and photochemical exchange with CO and H<sub>2</sub>, presumably through the  $16-e^{-}$  species IrH(CO)(dppp). The four-coordinate complex 5 exists under CO in equilibrium with its five-coordinate carbonyl adduct  $IrBr(CO)_2(dppe)$  and  $\nu_{CO}$  for both species are observed in solution.<sup>6</sup>

The observation of benzene carbonylation promoted photochemically by four-coordinate species of general formula IrX-(CO)(P P) prompted us to examine the photolysis of benzene solutions of Ir(I) and Rh(I) monodentate phosphine complexes under CO. We find that for  $IrCl(CO)(PPh_3)_2$ ,  $RhCl(CO)(PPh_3)_2$ , and RhH(CO)(PPh<sub>3</sub>)<sub>3</sub>, PhCHO forms under long-tern continuous The most extensively studied of these systems is irradiation. RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub>. In a typical experiment, a 7.2 mM solution of RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> [3.0 mg of complex in 0.6 mL of benzene] was placed under 100 mm of CO and the tube was flame-sealed. The sample was irradiated at ambient temperature through Pyrex. Benzaldehyde was observed by <sup>1</sup>H NMR in a concentration of 0.15 mM after 2 h of photolysis.<sup>7</sup> The concentration continued to increase until it remained essentially constant at 15 mM after 40 h of photolysis. This final aldehyde concentration was further confirmed by GC analysis.<sup>8</sup>

In all runs the amounts of PhCHO formed are small, never exceeding 3 turnovers. This is consistent with the unfavorable thermodynamics for the reaction  $C_6H_6 + CO = C_6H_5CHO$ , for

Wink, D.; Ford, P. C. J. Am. Chem. Soc. 1985, 107, 1794.
 (a) Ford, P. C. Biennial Inorganic Chemical Symposium, York University, Toronto, June 6-9, 1985, Session VI.
 (b) Ford, P. C., personal communication.
 (c) Wink, D. A.; Ford, P. C. J. Am. Chem. Soc. 1985, 107, 5566.

<sup>(3)</sup> Other photochemical studies of mononuclear d<sup>8</sup> complexes have focused primarily on the luminescence spectra and the nature of the excited state: Geoffroy, G. L.; Wrighton, M. S.; Hammond, G. S.; Gray, H. B. J. Am. Chem. Soc. 1974, 96. 3105. Brady, R.; Miller, W. V.; Vaska, L. J. Chem. Soc., Chem. Commun. 1974, 393. Johnson, C. E.; Eisenberg, R.; Evans, T. R.; Burberry, M. S. J. Am. Chem. Soc. 1983, 105, 1795. Fordyce, W. A.; Crosby, G. A. Inorg. Chem. 1982, 21, 1023. Recent work on binuclear d<sup>8</sup> complexes has included photochemical as well as photophysical studies. See, for example: Roundhill, D. M. J. Am. Chem. Soc. 1985, 107, 4354. Marshall, J. L.; Stobart, S. R.; Gray, H. B. J. Am. Chem. Soc. 1984, 106, 3029. (4) Parshall, G. W. "Homogeneous Catalysis" Wiley: New York, 1980; p 127. Collman, J. P.; Hegedus, L. S. "Principles and Applications of Organotransition Metal Chemistry"; University Science Books: Mill Valley, CA, 1980; p 585. (3) Other photochemical studies of mononuclear d<sup>8</sup> complexes have focused

<sup>1980;</sup> p 585.

<sup>(5)</sup> Fisher, B. J.; Eisenberg, R. Organometallics 1983, 2, 764. (6) (a) Infrared data in benzene solution for IrBr(CO)(dppe),  $\nu_{CO}$ , 1980 cm<sup>-1</sup>; IrBr(CO)<sub>2</sub>(dppe),  $\nu_{CO}$  2040 and 1940 cm<sup>-1</sup>. (b) Johnson, C. E.; Ei-senberg, R. J. Am. Chem. Soc. 1985, 107, 3148. (7) (a) All proton spectra were recorded on a Bruker WH-400 400-MHz spectrometer. (b) Benzaldehyde (PhCHO) was observed by <sup>1</sup>H NMR by using a solvent peak suppression program. The reaction solvent was 20% C<sub>6</sub>D<sub>6</sub> in C. H. and the installated foreware user 2777 67 H.

this is a solvent peak suppression program. The reaction solvent was  $20\% C_6 D_6$ in  $C_6 H_6$ , and the irradiated frequency was 7277.67 Hz. (8) Benzaldehyde was detected on 10 ft of Apiezon L on Chromosorb W on a Hewlett-Packard 5730A GLC (FID) equipped with an HP 3380A au-tomatic integrator. GLC conditions: injector port, 200 °C; oven, 150 °C; detector, 200 °C.



Figure 1. Aldehyde proton resonance (400 MHz) for RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> +  ${}^{13}$ CO + PhCHO in C<sub>6</sub>D<sub>6</sub> showing successive incorporation of  ${}^{13}$ C isotope into benzaldehyde. The resonance is at  $\delta$  9.63 ppm with  $J({}^{13}$ C-H) of 173 Hz. Irradiation times: (a) t = 0; (b) 1 day; (c) 5 days; (d) 10 davs.

which we calculate  $\Delta G^{\circ}_{298} = +1.7$  kcal.<sup>9</sup> The reverse reaction. PhCHO decarbonylation, has been studed in detail by others.<sup>10</sup> As a catalytic reaction using RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub>, it is found to require temperatures greater than 170 °C. In contrast, we find that  $RhCl(CO)(PPh_3)_2$  is a photochemical decarbonylation catalyst at ambient temperature, as shown by photolysis of a 7.2 mM solution of RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> (4  $\mu$ mol) containing 20  $\mu$ mol of PhCHO. After 24 h of photolysis, the amount of PhCHO had decreased to 12  $\mu$ mol as determined by GC analysis. In an earlier report, Geoffroy et al. observed that treatment of photolyzed solutions of RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> with PhCHO led to benzene formation, but they did not see direct photochemical decarbonylation using heptanal as substrate.<sup>11</sup>

In order to confirm that the low yields of PhCHO produced in the photolysis of benzene solutions of  $RhCl(CO)(PPh_3)_2$  were indeed the result of thermodynamic limitations and not kinetic inactivity of the system, an experiment was performed using <sup>13</sup>CO. A benzene- $d_6$  solution containing 3.0 mg of RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> (4.3  $\mu$ mol) and 2.0  $\mu$ L of PhCHO (19.6  $\mu$ mol) was placed under 1.2 atm of <sup>13</sup>CO and irradiated at 22 °C through a Pyrex filter. Examination of the aldehyde resonance by <sup>1</sup>H NMR ( $\delta$  9.63,  $J_{C-H}$ = 173 Hz) revealed significant incorporation of <sup>13</sup>CO into benzaldehyde, confirming the occurrence of carbonylation/decarbonylation sequences. As shown in Figure 1, the <sup>13</sup>C satellites of the aldehyde proton showed a steady increase in intensity over 10 days at which time the extent of <sup>13</sup>CO incorporation into PhCHO amounted to 38%. During this period, the PhCHO concentration decreased from 36 mM (5 equiv) to 21 mM (2.9 equiv) as determined by GC analysis.<sup>8</sup>

While this demonstrates the reversibility of photochemical carbonylation, it raises the question of whether there is a common intermediate for both the forward and reverse reactions or a photostationary state resulting from two independent photoreactions. To test this, three identical samples were prepared (7.5 mM complex, 12 mM PhCHO, and 1.0 atm of CO) and irradiated through a Pyrex filter for 3 days. One tube was opened and GC analysis showed that the PhCHO concentration had decreased to 10.5 mM. The other two samples were placed at distances 4 and 16 cm from the light source and irradiated for another day. Analysis of these two samples showed each to be 10.1 mM in PhCHO, indicating that carbonylation and decarbonylation proceed through a common intermediate and result eventually in an equilibrium mixture of PhCHO, CO, and benzene.

To account for the incorporation of <sup>13</sup>CO into PhCHO under photolysis, equivalence of <sup>12</sup>CO and <sup>13</sup>CO is required at some point in the catalysis. This may be achieved via free CO in solution or a dicarbonyl intermediate. The pathways leading to photochemical carbonylation and decarbonylation involve the same initial step which we believe is loss of a ligand to generate a 14-e intermediate. While Ford and Wink have demonstrated efficient

photochemical dissociation of CO from MCl(CO)(PPh<sub>3</sub>)<sub>2</sub>, they see no net reaction chemistry under the conditions of their experiment.<sup>1,2</sup> The formation of benzaldehyde in the present study may result from either a relatively inefficient reaction of MCl- $(PPh_3)_2$  with C<sub>6</sub>H<sub>6</sub> leading to arene activation and carbonylation, or from a second photochemical process which occurs much less efficiently than CO photodissociation. This process could be phosphine dissociation to generate MCl(CO)(PPh<sub>3</sub>).

Support for the notion of ligand dissociation as the photochemical step is obtained from photolyses carried out under different CO pressures and in the presence of added phosphine. Two 7.5 mM benzene solutions of RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> under 36 and 615 mm of CO were prepared. The samples were irradiated and monitored by <sup>1</sup>H NMR spectroscopy<sup>7b</sup> for 20 h. Analysis by GC showed the samples to be 4.1 and 1.8 mM, respectively, in benzaldehyde. Thus the efficiency of benzene carbonylation is reduced as CO pressure is increased. The addition of  $PPh_3$  to a benzene solution of RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub> under CO causes rapid formation of  $RhCl(PPh_3)_3$  upon irradiation in accord with the results of Ford<sup>1</sup> and inhibition of benzaldehyde formation.

While further mechanistic details remain to be elucidated, the key point of the present study is that Rh(I) and Ir(I) phosphine complexes do exhibit photochemical reactivity which leads to significant, new reaction chemistry. If the arene carbonylation reaction seen here can be driven to overcome its unfavorable thermodynamics such as by subsequent hydrogenation, this photochemical functionalization may prove useful as well as interesting.

Acknowledgment. We thank the National Science Foundation (CHE 83-08064) for support of this work, and the Johnson Matthey Co., Inc., for a generous loan of rhodium salts. We also thank Prof. Jack Kampmeier and Tong-Zhou Liu for helpful discussions and Dr. Romana Lashewycz-Rubycz for experimental assistance.

## Sustained Epoxidation of Olefins by Oxygen Donors Catalyzed by Transition Metal Substituted Polyoxometalates, Oxidatively Resistant Inorganic Analogues of Metalloporphyrins

Craig L. Hill\* and Robert B. Brown, Jr.

Department of Chemistry, Emory University Atlanta, Georgia 30322

Received July 1, 1985

The homogeneous catalytic oxygenation of saturated and unsaturated hydrocarbons has been one of the most active areas of chemical research in the last 5 years.<sup>1-12</sup> Numerous metallo-

<sup>(9)</sup> Calculated with thermodynamic constants from "Handbook of Chem-

 <sup>(3)</sup> Calculated with the individuality constants from Francock of Chemistry and Physics", 62nd ed.; CRC Press: Boca Raton, FL, 1981. Benson,
 S. W. "Thermochemical Kinetics"; Wiley-Interscience: New York, 1976.
 (10) Doughty, D. H.; Pignolet, L. H. In "Homogeneous Catalysis with Metal Phosphine Complexes"; Pignolet, L. H., Ed.; Plenum Press: New York,

<sup>1983;</sup> Chapter 11.
(11) Geoffroy, G. L.; Denton, D. A.; Keeney, M. E.; Bucks, R. R. Inorg. Chem. 1976, 15, 2382.

<sup>(1)</sup> Sheldon, R. A.; Kochi, J. K. "Metal-Catalyzed Oxidations of Organic Compounds"; Academic Press: New York, 1981.

 <sup>(2) (</sup>a) Smegal, J. A.; Schardt, B. C.; Hill, C. L. J. Am. Chem. Soc. 1983, 105, 3510. (b) Smegal, J. A.; Hill, C. L. Ibid. 1983, 105, 3515. (c) Hill, C. L.; Smegal, J. A.; Henly, T. J. J. Org. Chem. 1983, 48, 3277. (d) Smegal, J. A.; Hill, C. L. J. Am. Chem. Soc. 1983, 105, 2920 and references cited therein.

<sup>(3) (</sup>a) Groves, J. T.; Quinn, R. J. Am. Chem. Soc. 1985, 107, 5790. (b)
(c) Groves, J. T.; Subramanian, D. V. Ibid. 1984, 106, 2177. (c) Groves, J. T.;
Nemo, T. E. Ibid. 1983, 105, 5786. (d) Groves, J. T.; Takahashi, T. Ibid.
1983, 105, 2073. (e) Groves, J. T.; Wantanabe, Y.; McMurry, T. J. Ibid.
1983, 105, 4489 and references cited in each.
(4) (a) Mashiko, T.; Dolphin, D.; Nakano, T.; Traylor, T. G. J. Am. Chem.
Soc. 1985, 107, 3735. (b) Traylor, P. S.; Dolphin, D. H.; Traylor, T. G. J.

Chem. Soc., Chem. Commun. 1984, 279.

<sup>(5) (</sup>a) Powell, M. F.; Pai, E. F.; Bruice, T. C. J. Am. Chem. Soc. 1984, 106, 3277. (b) Nee, M. W.; Bruice, T. C. Ibid. 1982, 104, 6123 and references cited therein.