

96. 2'-Deoxy- β -D-ribofuranosides of N^6 -Methylated 7-Deazaadenine and 8-Aza-7-deazaadenine: Solid-Phase Synthesis of Oligodeoxyribonucleotides and Properties of Self-Complementary Duplexes

by Frank Seela*, Klaus Kaiser, and Uwe Bindig

Laboratorium für Organische und Bioorganische Chemie, Fachbereich Biologie/Chemie, Universität Osnabrück, Barbarastr. 7, D-4500 Osnabrück

(17.II.89)


The syntheses of 7-deaza- N^6 -methyladenine N^9 -(2'-deoxy- β -D-ribofuranoside) (**2**) as well as of 8-aza-7-deaza- N^6 -methyladenine N^8 - and N^9 -(2'-deoxyribofuranosides) (**3** and **4**, resp.) are described. A 4,4'-dimethoxytritylation followed by phosphorylation yielded the methyl phosphoramidites **12–14**. They were employed together with the phosphoramidite of 2'-deoxy- N^6 -methyladenosine (**15**) in automated solid-phase oligonucleotide synthesis. Alternating or palindromic oligonucleotides derived from d(A-T)₆ or d(A-T-G-C-A-G-A*-T-C-T-G-C-A) but containing one methylated pyrrolo[2,3-*d*]pyrimidine or pyrazolo[3,4-*d*]pyrimidine moiety in place of a N^6 -methylaminopurine (A*) were synthesized. Melting experiments showed that duplex destabilization induced by a N^6 -Me group of 2'-deoxy- N^6 -methyladenosine is reversed by incorporation of 8-aza-7-deaza-2'-deoxy- N^6 -methyladenosine, whereas 7-deaza-2'-deoxy- N^6 -methyladenosine decreased the T_m value further. Regiospecific phosphodiester hydrolysis of d(A-T-G-C-A-G-m⁶A-T-C-T-G-C-A) by the endodeoxyribonuclease Dpn I, yielding d(A-T-G-C-A-G-m⁶A) and d(pT-C-T-G-C-A), was prevented when the residue c⁷m⁶A_d (**2**), c⁷m⁶z⁸A_d (**3**), or c⁷m⁶z⁸A_d' (**4**) replaced m⁶A_d (**1**) indicating that N(7) of N^6 -methyladenine is a proton-acceptor site for the endodeoxyribonuclease.

1. Introduction. – The methylation of the adenine 6-NH₂ group and of the 5-position of cytosine is the most common naturally occurring modification of the DNA molecule. With those methylated bases, gene expression is controlled [1] and enzymatic phosphodiester hydrolysis of DNA by endodeoxyribonucleases can be prevented [2]. As a consequence, it has been suggested that the Me group attached to 2'-deoxyadenosine alters DNA's tertiary structure [3].






Up to now, 2'-deoxy- N^6 -methyladenosine (m⁶A_d, **1**) has been incorporated into DNA fragments employing chemical [4] [5] or enzymatic methods [3]. As the T_m value of those oligonucleotides is strongly decreased and only little information exists about the interaction of endodeoxyribonucleases recognizing such methylated purines or pyrimidines [6], we became interested in N^6 -methylated 2'-deoxyadenosine derivatives. They should lack potential enzyme binding positions (N(7) of purine) and form more stable duplexes by increased base stacking or stronger base pairing. In the following, we report on the syntheses of 7-deaza-2'-deoxy- N^6 -methyladenosine¹⁾ (c⁷m⁶A_d, **2**) as well as of 8-aza-7-deaza- N^6 -methyl-2'-deoxyadenosine (**3**) and its N^8 -(2'-deoxy- β -D-ribofuranoside) **4** and on their conversion into the corresponding phosphoramidites **12–14**. These monomeric building blocks are then used together with regular phosphoramidites in automated oligonucleotide synthesis. The resulting oligomers are studied with respect to duplex stability or regiospecific phosphodiester hydrolysis by the endodeoxyribonuclease Dpn I.

¹⁾ Purine nomenclature is used throughout *Chapt. 1–4*.




2. Synthesis of 7-Deaza-*N*⁶-methyladenine *N*⁹-(2'-Deoxy- β -D-ribofuranoside) (2) as well as of 8-Aza-7-deaza-*N*⁶-methyladenine *N*⁸- and *N*⁹-(2'-Deoxy- β -D-ribofuranosides) (3 and 4, respectively). – For the preparation of 2, three methods were considered: *i*) methylation of 9e at N(1) followed by *Dimroth* rearrangement, as reported for the corresponding ribonucleoside [7], *ii*) nucleophilic displacement of a reactive 6-substituent by MeNH₂, and *iii*) glycosylation of 7-deaza-*N*⁶-methyladenine (5a) with 2-deoxy-3,5-di-*O*-(*p*-toluoyl)- α -D-*erythro*-pentofuranosyl chloride (6). As our laboratory has developed a method of regio- and stereoselective glycosylation of purine analogues with the




1 (purine numbering)

2 (systematic numbering)¹⁾3 (systematic numbering)¹⁾4 (systematic numbering)¹⁾


KOH, TDA-1







9a R¹ = CH₃NH, R² = H, R³ = Tol
 b R¹ = CH₃NH, R² = Tol, R³ = H
 c R¹ = CH₃NH, R² = (MeO)₂Tr, R³ = Tol
 d R¹ = CH₃SO₂, R² = R³ = Tol
 e R¹ = NH₂, R² = R³ = H



10a R¹ = CH₃O, R² = R³ = Tol
 b R¹ = CH₃NH, R² = (MeO)₂Tr, R³ = H
 c R¹ = NH₂, R² = R³ = H



11a R¹ = CH₃O, R² = R³ = Tol
 b R¹ = CH₃NH, R² = (MeO)₂Tr, R³ = H
 c R¹ = NH₂, R² = R³ = H



halogenose **6** [8] [9], we have employed solid-liquid phase-transfer glycosylation on compound **5a**. The reaction was performed in MeCN in the presence of powdered KOH and the cryptand *TDA-1* at room temperature. Within 15 min, **7a** was formed in 66% yield together with a minor amount of partially deprotected **9a** and **9b**. In contrast to glycosylations performed on other pyrrolo[2,3-*d*]pyrimidines, this reaction was particularly sensitive towards sugar deprotection. It is not clear whether this is due to a catalytic action of the MeNH group. However, detoluoylation of **7a**, **9a**, or **9b** in conc. aq. NH₃/MeOH yielded **2** which was isolated after chromatographic purification.

The glycosylation site as well as the anomeric configuration of **2** were determined by ¹H-NMR NOE difference spectroscopy [10]. Table 1 shows NOE's for H-C(4') and H_α-C(2') upon irradiation of H-C(1') which indicate spatial proximity of those protons and confirm the β-D-configuration. The NOE of H-C(6) confirms the glycosylation position. The structure of the partially deprotected compounds **9a** and **9b** was assigned by the ¹³C-NMR data of the sugar moiety (Table 2). Deprotection at the C(3') position of **7a** (→**9b**) shifts C(3') upfield and C(2') as well as C(1') downfield (**9b**). Deprotection of the 5'-position (→**9a**) results in an upfield shift of C(5') and smaller downfield shift of the other C-atoms [11].

Table 1. NOE Data [%] of **2**, **8a**, and **11c** upon Irradiation of H-C(1') and of **4** upon Irradiation of H-C(3) ((D₆)Me₂SO, 23°)

	H-C(1')	H _α -C(2')	H-C(4')	H-C(5')	H-C(6) ^{a)}	H-C(3) ^{b)}
2	–	6.1	2.3	–	4.8	–
8a	–	10.0	3.5	–	9.5	–
4	9.5	1.3	1.6	2.7	–	–
11c	–	4.8	2.0	–	–	8.3

^{a)} Pyrrolo[2,3-*d*]pyrimidine numbering.

^{b)} Pyrazolo[3,4-*d*]pyrimidine numbering.

Alternatively, compound **2** is accessible from the sugar-protected 6-chloro compound **7c** [11] by halogen displacement with MeNH₂ under simultaneous sugar deprotection in 65% yield. The reaction conditions required elevated temperature and a steel bomb, thus making this procedure more laborious. The reaction proceeded much faster and occurred already at 60° in a pressure bottle, when the sugar-protected sulfone **9d** was used, yielding **2** in 83%. For the synthesis of the sulfone **9d**, 4-(methylthio)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**5b**) was glycosylated under the same conditions as already described for compound **5a**. The glycosylation product **7b** (80% yield) was deprotected to give **8a** which was oxidized with 3-chloroperbenzoic acid to **9d** as the only product. Under the same conditions, the oxidation of the deprotected 6-ethylthio compound **8c** prepared from **8b** with ethanethiol gave a mixture of the sulfone **8e** and the diastereoisomeric sulfoxides **8d**/**8d'**. The ¹³C-NMR chemical shifts of **5b**, **7b**, **8a**, and **9d** were assigned on the basis of gated-decoupled ¹³C-NMR spectra.

Preparation of **2** by methylation of **9e** was not considered as this route afforded too many steps. Comparing the total yields of the different synthetic routes (43% of **2** via **5c**, **5a**, **7a**; 53% of **2** via **5c**, **7c**; 54% of **2** via **5b**, **7b**, **9d**) shows that introduction of the MeNH group on the nucleoside level gave the best results. Although an additional step was necessary for the route via the sulfone, a reaction in a steel bomb was not required.

Table 2. ^{13}C -NMR Chemical Shifts ($(\text{D}_6)\text{Me}_2\text{SO}$)^{a)} of Purine, Pyrrolo[2,3-*d*]pyrimidine, and Pyrazolo[3,4-*d*]pyrimidine 2'-Deoxyribofuranosides^{b)}

	C(2) (C(2)) [C(6)]	C(6) (C(4)) [C(4)]	C(5) (C(4a)) [C(3a)]	– (C(5)) [C(3)]	C(8) (C(6)) –	C(4) (C(7a)) [C(7a)]	CH ₂	CH ₃
1	152.5	155.1	119.9	–	139.4	147.9	–	27.1
2	151.6	156.7	103.2	99.3	121.6	149.0	–	27.2
3	156.1	156.8	101.1	132.7	–	153.2	–	27.0
4	156.6	158.2	101.9	123.6	–	159.2	–	27.0
5a	151.5	156.7	102.6	98.6	120.8	150.5	–	27.2
5b	150.3	159.8	115.1	98.1	125.4	148.8	–	10.9
7a	151.8	156.2	103.4	100.1	121.1	149.4	–	27.2
7b	150.5	160.9	116.1	99.8	125.9	148.1	–	11.4
8a	150.5	160.6	115.9	99.3	125.9	147.8	–	11.4
8c	150.4	159.8	116.0	99.2	125.6	147.9	22.6	15.0
8d	151.52 ^{c)}	163.87 ^{c)}	117.38	100.01	129.08	149.97 ^{c)}	46.71	5.47
8d'	151.50 ^{c)}	163.87 ^{c)}	117.38	100.01	129.04	149.97 ^{c)}	46.68	5.41
8e	152.9 ^{c)}	153.9 ^{c)}	115.3	100.3	131.4	150.3 ^{c)}	46.1	6.8
9a	151.6	156.8	103.5	99.6	121.8	149.0	–	27.2
9b	151.9	156.7	103.2	99.6	121.0	149.3	–	27.2
9c	151.8	156.7	103.2	99.6	120.9	149.3	–	27.2/55.1
9d	153.2	155.6	114.3	100.6	131.4	150.4	–	–
10b	156.1	156.8	101.1	132.5	–	153.3	–	26.9/55.0
11b	156.5	158.3	102.2	124.1	–	159.3	–	27.0/55.0

	C(1')	C(2')	C(3')	C(4')	C(5')
1	84.1	– ^{d)}	71.1	88.1	62.0
2	83.4	– ^{d)}	71.3	87.4	62.2
3	84.1	38.1	71.2	87.7	62.6
4	90.6	39.5	70.8	88.5	62.2
7a	81.1	36.2	75.2	83.2	64.4
7b	81.3	36.1	75.1	83.6	64.3
8a	83.1	– ^{d)}	71.1	87.5	62.0
8c	83.0	40.1	71.0	87.5	61.9
8d	83.00	– ^{d)}	71.00	87.67	61.86
8d'	82.94	– ^{d)}	71.00	87.67	61.86
8e	83.3	– ^{d)}	70.9	87.8	61.8
9a	83.9	37.0	76.2	84.7	62.0
9b	82.6	– ^{d)}	70.8	83.5	64.6
9c	82.5	38.0	70.9	85.5	64.3
9d	81.6	36.2	74.9	83.9	64.2
10b	83.5	38.3	70.9	85.2	64.5
11b	89.9	39.5	70.7	85.4	64.2

a) δ Values given in ppm relative to Me_4Si as internal standard.
b) Pyrrolo[2,3-*d*]pyrimidine numbering in parentheses, pyrazolo[3,4-*d*]pyrimidine numbering in brackets.
c) Tentative.
d) Superimposed by Me_2SO .

For the synthesis of 8-aza-7-deaza- N^6 -methyladenine N^8 - and N^9 -(2'-deoxy- β -D-ribofuranosides) (**3** and **4**, resp.), the already described 4-methoxy-1*H*-pyrazolo[3,4-*d*]pyrimidine 2'-deoxy- β -D-ribofuranosides **10a** and **11a** [12a,b] were used as starting materials. As their MeO groups are much more reactive than those of pyrrolo[2,3-*d*]-

pyrimidines and even of purines, displacement by MeNH_2 occurred at 60° in a pressure bottle affording compounds **3** and **4**, respectively, in 95% yield. Their structures were confirmed by ^1H - and ^{13}C -NMR spectroscopy (see *Exper. Part* and *Table 2*). The strong NOE of $\text{H}-\text{C}(1')$ of **4** upon irradiation of $\text{H}-\text{C}(3)$ ($\text{H}-\text{C}(7)$ in purines numbering) provided the unequivocal assignment of $\text{H}-\text{C}(6)$ and $\text{H}-\text{C}(3)$ and showed that $\text{H}-\text{C}(3)$ resonates downfield from $\text{H}-\text{C}(6)$.

This conclusion should be transferable to compound **11c** and requires a correction of an earlier ^1H -NMR assignment [12a]. This correction was verified by an NOE-difference spectrum of **11c** with saturation of $\text{H}-\text{C}(1')$ (*Table 1*). In connection with these NMR experiments, ^1H , ^{13}C correlation spectra of **1** and **dA** have been measured. They showed that $\text{H}-\text{C}(2)$ signals are located upfield as compared to $\text{H}-\text{C}(8)$, which is opposite to an assignment reported earlier [13].

3. Solid-Phase Synthesis of Oligonucleotides Containing 1–4. – Solid-phase synthesis [14] of the oligonucleotides **19–27** was performed using P(III) chemistry [15] which




Fig. 1. HPLC profiles after phosphodiester hydrolysis of **19b–22b** with snake-venom phosphodiesterase followed by alkaline phosphatase. Gradient IV; conditions, see *Exper.* Part. a) From **19b**; b) from **20b**; c) from **21b**; d) from **22b**.

required the phosphoramidites **12–15** as monomeric building blocks. The synthesis of **15** has already been reported [4] [5]. For the synthesis of **12–14**, the 5'-OH group of **2–4** was protected with 4,4'-dimethoxytrityl chloride ((MeO)₂TrCl) to give **9c**, **10b**, and **11b**, respectively. The amorphous compounds were characterized by NMR spectroscopy and elemental analyses. The position of dimethoxytritylation was determined from the characteristic ¹³C-NMR upfield shift of the C(4') signal and the downfield shift of C(5') (Table 2).

It has been shown earlier for 2'-deoxy-*N*⁶-methyladenosine and related nucleosides that protection of the 6-methylamino group was not required during oligonucleotide synthesis [6]. Nucleobase protection was also not necessary in the case of **2–4**. The AcNCH₃ group, which can be formed upon the capping procedure of the oligonucleotide synthesis, will be cleaved together with the other base-protecting groups by NH₃ treatment. The phosphoramidites **12–14** were obtained from **9c**, **10b**, and **11b** by reaction with chloro(diisopropylamino)methoxyphosphine. They were purified by flash chromatography and characterized by ³¹P-NMR spectroscopy.

The synthesis of the oligonucleotides **19–27** was carried out on solid support in an automated DNA synthesizer using the regular phosphoramidites of dA, dT, dG, and dC

d(A-T) ₆ [13] 16	d(c ⁷ z ⁸ A-T) ₆ [13] 17	d(c ⁷ z ⁸ A'-T) ₆ [13] 18
d([(MeO) ₂ Tr]A-T-A-T-A-T-m ⁶ A-T-A-T-A-T) 19a		d(A-T-A-T-A-T-m ⁶ A-T-A-T-A-T) 19b
d([(MeO) ₂ Tr]A-T-A-T-A-T-c ⁷ m ⁶ z ⁸ A-T-A-T-A-T) 20a		d(A-T-A-T-A-T-c ⁷ m ⁶ z ⁸ A-T-A-T-A-T) 20b
d([(MeO) ₂ Tr]A-T-A-T-A-T-c ⁷ m ⁶ z ⁸ A'-T-A-T-A-T) 21a		d(A-T-A-T-A-T-c ⁷ m ⁶ z ⁸ A'-T-A-T-A-T) 21b
d([(MeO) ₂ Tr]A-T-A-T-A-T-c ⁷ m ⁶ A-T-A-T-A-T) 22a		d(A-T-A-T-A-T-c ⁷ m ⁶ A-T-A-T-A-T) 22b
d([(MeO) ₂ Tr]A-T-G-C-A-G-A-T-C-T-G-C-A) 23a		d(A-T-G-C-A-G-A-T-C-T-G-C-A) 23b
d([(MeO) ₂ Tr]A-T-G-C-A-G-m ⁶ A-T-C-T-G-C-A) 24a		d(A-T-G-C-A-G-m ⁶ A-T-C-T-G-C-A) 24b
d([(MeO) ₂ Tr]A-T-G-C-A-G-c ⁷ m ⁶ z ⁸ A-T-C-T-G-C-A) 25a		d(A-T-G-C-A-G-c ⁷ m ⁶ z ⁸ A-T-C-T-G-C-A) 25b
d([(MeO) ₂ Tr]A-T-G-C-A-G-c ⁷ m ⁶ z ⁸ A'-T-C-T-G-C-A) 26a		d(A-T-G-C-A-G-c ⁷ m ⁶ z ⁸ A'-T-C-T-G-C-A) 26b
d([(MeO) ₂ Tr]A-T-G-C-A-G-c ⁷ m ⁶ A-T-C-T-G-C-A) 27a		d(A-T-G-C-A-G-c ⁷ m ⁶ A-T-C-T-G-C-A) 27b

c⁷z⁸A_d = 8-aza-7-deaza-2'-deoxyadenosine; m⁶A_d = 2'-deoxy-*N*⁶-methyladenosine; c⁷m⁶A_d = 7-deaza-2'-deoxy-*N*⁶-methyladenosine; c⁷m⁶z⁸A_d = 8-aza-7-deaza-2'-deoxy-*N*⁶-methyladenosine. A' is the symbol used for the 8-aza-7-deazaadenosine moiety with the unusual N⁸- instead of the usual N⁹-glycosylic linkage (purine numbering).

together with the modified compounds **12–15**. The syntheses followed a protocol of detritylation, coupling, capping, and oxidation according to [16]. The protected oligomers were demethylated with thiophenol and cleaved from the solid support by treatment with NH_3 . After removal of the base-protecting groups by ammonolysis, purification of the $(\text{MeO})_2\text{Tr}$ -protected oligomers **19a–27a** by HPLC, and removal of the $(\text{MeO})_2\text{Tr}$ residue with aq. AcOH , the oligomers **19b–27b** were recovered and again purified by HPLC to give colorless solids.

The nucleoside content of the oligonucleotides (see *Fig. 1*) was determined after tandem hydrolysis with snake-venom phosphodiesterase followed by alkaline phosphatase. UV Quantification of the HPLC pattern confirmed the molar amount of monomers.

4. Physical Properties of the Oligomers 16–27b and Influence of 1–4 on the Phosphodiester Hydrolysis by Endodeoxyribonuclease Dpn I. – It has been reported that N^6 -methylation of dA within oligonucleotides decreases the T_m value strongly [3] [4]. This is demonstrated by the T_m values of $\text{d}(\text{A-T})_6$ (**16**) and the modified dodecamer **19b**, showing a ΔT_m of 6° (*Table 3*). The destabilization by N^6 -methylation has been attributed to a restricted rotation of the $N^6\text{-Me}$ group, preferentially located in *cis* orientation with respect to $\text{N}(1)$ [17]. This should interfere with *Watson-Crick* base pairing towards dT.

Table 3. Melting Temperatures (T_m Values) of the Oligonucleotides **16–18^{a)}**, **19b–22b^{a)}**, and **23b–27b^{b)}**

Oligomer		T_m [°]
$\text{d}(\text{A-T})_6$ [12b]	(16)	32
$\text{d}(\text{c}^7\text{z}^8\text{A-T})_6$ [12b]	(17)	36
$\text{d}(\text{c}^7\text{z}^8\text{A'-T})_6$ [12b]	(18)	48
$\text{d}(\text{A-T-A-T-A-T-m}^6\text{A-T-A-T-A-T})$	(19b)	26
$\text{d}(\text{A-T-A-T-A-T-c}^7\text{m}^6\text{z}^8\text{A-T-A-T-A-T})$	(20b)	28
$\text{d}(\text{A-T-A-T-A-T-c}^7\text{m}^6\text{z}^8\text{A'-T-A-T-A-T})$	(21b)	31
$\text{d}(\text{A-T-A-T-A-T-c}^7\text{m}^6\text{A-T-A-T-A-T})$	(22b)	24
$\text{d}(\text{A-T-G-C-A-G-A-T-C-T-G-C-A})$	(23b)	59
$\text{d}(\text{A-T-G-C-A-G-A-T-C-T-G-C-A})$	(23b)	59
$\text{d}(\text{A-T-G-C-A-G-m}^6\text{A-T-C-T-G-C-A})$	(24b)	53
$\text{d}(\text{A-T-G-C-A-G-c}^7\text{m}^6\text{z}^8\text{A-T-C-T-G-C-A})$	(25b)	54
$\text{d}(\text{A-T-G-C-A-G-c}^7\text{m}^6\text{z}^8\text{A'-T-C-T-G-C-A})$	(26b)	55
$\text{d}(\text{A-T-G-C-A-G-c}^7\text{m}^6\text{A-T-C-T-G-C-A})$	(27b)	51

^{a)} In 1M NaCl, 0.1M MgCl_2 , and 60 mM cacodylic acid, pH 7.0 at $4\ \mu\text{M}$ single-strand concentration.

^{b)} In 150 mM NaCl, 6 mM MgCl_2 , 7 mM 2-mercaptoethanol, and bovine serum albumine (100 $\mu\text{g/ml}$) at $4\ \mu\text{M}$ single-strand concentration.

However, the situation seems to be much more complex. From the $+I$ effect of the Me group, the MeNH group of **1** is expected to be a weaker proton donor than an NH_2 group, thus destabilizing the H-bond to the carbonyl group of dT for electronic as well as for sterical reasons. Furthermore, the bathochromic shift (10 nm) in the UV of **1** as compared to dA indicates that N^6 -methylation affects the electronic state of the nucleobase and thus stacking interactions. Finally, the H_2O spine located within the major groove of an oligonucleotide duplex [18] may be affected by introducing a $N^6\text{-Me}$ group. As the H_2O -accepting $\text{N}(7)$ may not be accessible in case of **1** and is not present in compound **2**, a destabilization of the duplex may occur. The residues **3** or **4**, however, having an N-atom within the major groove ($\text{N}(8)$) may again allow the formation of a hydrate.

We have already reported on increased T_m values of $d(A-T)_6$ (**16**) which contains 8-aza-7-deaza-2'-deoxyadenosine instead of dA residues [12b] and have explained this finding by a better base overlap. A stabilization is also observed in the case of the oligomer **20b** as compared to **19b**. However, the T_m increase is smaller due to the fact that in this case, only one m^6A_d is replaced by **3**. In contrast, the pyrrolo[2,3-*d*]pyrimidine nucleoside **2** destabilizes the duplex structure as demonstrated for the oligomer **22b** (Table 3).

A strong stabilization of $d(A-T)_6$ (**16**) occurs also by replacing every dA residue with the 8-aza-7-deazaadenine N^8 -(2'-deoxy- β -D-ribofuranoside) ($= c^7z^8A'_d$; **11c**; see T_m of **16** and **18** in Table 3) [12b]. The same phenomenon is observed in the case of **21b** when compared to **19b**. Because of the strong T_m increase for the oligomer **18** as compared to **16**, a special tertiary structure of this duplex is assumed. As DNA tertiary structures can be visualized by the CD spectrum [19], we have measured the CD spectra of **16–18** and




Fig. 2. CD Spectra of the dodecamers a) **16** (—), **17** (---), and **18** (----) and b) **19b** (—), **20b** (---), **21b** (-----), and **22b** (—). In 1M NaCl, 0.1M MgCl₂, and 60 mM sodium cacodylate at 10 μ M single-strand concentration.

19b–22b (Fig. 2). The CD of **17** and **19b–22b** are very similar to that of $d(A-T)_6$ (**16**) indicating that those oligomers form a right-handed B-DNA-resembling structure as found for $\text{poly}[d(A-T)] \cdot \text{poly}[d(A-T)]$ at low salt concentration [20]. However, the CD of **18** is different (Fig. 2a). The existence of a negative Cotton effect at 300 nm and a positive band at 270 nm points to a left-handed helix as found for Z-DNA [21]. This left-handed DNA is characteristic for $\text{poly}[d(G-C)] \cdot \text{poly}[d(G-C)]$ at high salt concentration [22] and has also been observed for $\text{poly}[d(A-T)] \cdot \text{poly}[d(A-T)]$ at 5M NaCl in the presence of 95 mM NiCl_2 [20]. Characteristic features of Z-DNA are repeating elements of a $C(3')\text{-endo/syn}$ and $C(2')\text{-endo/anti}$ sugar conformation with the purine bases in *syn* orientation [23].

In the case of the **11c** residue a reversed *Watson-Crick* base pair between the modified adenine and thymine should be preferred. Computer modelling of a $c^2z^8A_d'-T_d$ base pair indicates that a left-handed Z-DNA can be formed if a reversed *Watson-Crick* base pair is formed. As the special CD spectrum is only found for **18**, (Fig. 2a) in which every dA residue is replaced by **11c**, but not for **21b** one can conclude that one m^6A_d replacement by **4** is not sufficient to alter the structure of the whole duplex.

Apart from the alternating oligomers, we have determined the T_m values of the 13-mers **23b–27b**. Compound **24b** contains the recognition sequence $d(G-m^6A-T-C)$ of the endodeoxyribonuclease Dpn I (*Diplococcus Pneumonia*) [24–26]. This restriction enzyme is one of the rare examples where N^6 -methylation of the adenine moiety within the recognition site is required for regiospecific phosphodiester hydrolysis [27]. In most other cases, N^6 -methylation of adenine protects DNA from phosphodiester hydrolysis by restriction enzymes [2]. As it is unknown if N(7), located in the major groove of DNA, is necessary as a proton acceptor for enzym binding, we have incorporated compounds **2–4** instead of m^6A_d into the 13-mer $d(A-T-G-C-A-G-m^6A-T-C-T-G-C-A)$ (**24b**). The oligomers **23b–27b** were then subjected to hydrolysis by the endodeoxyribonuclease Dpn I. Enzymatic hydrolysis was followed by HPLC. As Fig. 3 shows, the 13-mer **24b** is cleaved into the heptamer $d(A-T-G-C-A-G-m^6A)$ and the hexamer $d(pT-C-T-G-C-A)$ within 20 h at 22°. Complete hydrolysis of the oligomer by snake-venom phosphodiesterase followed by alkaline phosphatase resulted in a HPLC pattern (Fig. 4) from which the nucleoside




Fig. 3. Time course of phosphodiester hydrolysis of the oligomer $d(A-T-G-C-A-G-m^6A-T-C-T-G-C-A)$ (**24b**) by treatment with the endodeoxyribonuclease Dpn I. Conditions, see Exper. Part.




Fig. 4. HPLC profiles of the products a) of phosphodiester hydrolysis of *d*(*A-T-G-C-A-G-m*⁶*A-T-C-T-G-C-A*) (**24b**) by the endodeoxyribonuclease Dpn I (conditions, see *Exper. Part*; gradient II) and b), c) after tandem hydrolysis of *d*(*pT-C-T-G-C-A*) (peak I of a) and *d*(*A-T-G-C-A-G-m*⁶*A*) (peak II of a), respectively, with snake-venom phosphodiesterase followed by alkaline phosphatase (gradient III). ☆: buffer peaks.

content was calculated on the basis of the extinction coefficients. As expected, the non-methylated **23b** was not hydrolysed at all by the endodeoxyribonuclease Dpn I, a finding which confirms results on the oligonucleotide level already observed on native DNA [24]. None of the oligomers **25b–27b** containing the *m*⁶*A*_d derivatives **2–4** were hydrolyzed by the endodeoxyribonuclease Dpn I which shows that N(7) is required as a proton-acceptor position for regiospecific phosphodiester hydrolysis.

We thank Dr. H. Rosemeyer for the discussion of NMR spectra and Mrs. M. Beste for technical assistance. Financial support from the *Deutsche Forschungsgemeinschaft* and the *Stiftung Volkswagenwerk* is gratefully acknowledged.

Experimental Part

General. The phosphoramidites were prepared from the 5'-tritylated and appropriately base-protected nucleosides [(MeO)₂Tr]bz⁶A_d, [(MeO)₂Tr]ib²G_d, [(MeO)₂Tr]bz⁴C_d, and [(MeO)₂Tr]T_d (Biosyntech, Hamburg, FRG). *Fractosil 500* polymeric support was purchased from Biosyntech (Hamburg, FRG). Snake-venom phosphodiesterase (EC 3.1.4.1., *Crotallus durissus*), alkaline phosphatase (EC 3.1.3.1., *E. coli*), and endodeoxyribonuclease DpnI were purchased from Boehringer Mannheim (FRG). Oligonucleotide synthesis was carried out on an automated DNA synthesizer, model 380B, of Applied Biosystems (Weiterstadt, FRG) on a 1-μmol scale. TLC: silica gel *SIL G-25 UV*₂₅₄ plates (Macherey-Nagel, FRG). Flash chromatography: silica gel 60H (Merck, FRG), at 0.9 bar; Solvent systems: CH₂Cl₂/MeOH 9:1 (A), CH₂Cl₂/MeOH 8:2 (B), CH₂Cl₂/acetone 7:3 (C), CH₂Cl₂/MeOH 95:5 (D), CH₂Cl₂/AcOEt/Et₃N 45:45:10 (E), CHCl₃ (F), CHCl₃/MeOH 95:5 (G), CHCl₃/MeOH 9:1 (H), CHCl₃/MeOH 8:2 (I), cyclohexane/AcOEt 6:4 (K). UV spectra: Hitachi-150-20 spectrophotometer; (Hitachi, Japan); λ_{max} (ε) in nm. CD-spectra: Jasco-J-600 spectropolarimeter. NMR spectra: Bruker AC 250 spectrometer; δ in ppm relative to tetramethylsilane (= 0 ppm) as internal standard (¹H and ¹³C) or to external 85% phosphoric acid (³¹P; = 0 ppm).

Melting Experiments. The melting experiments were carried out in a thermostatically controlled cell holder with a Shimadzu-210-A UV spectrophotometer connected with a Kipp and Zonen BD 90 recorder. The increase of absorbance at the appropriate wavelength as a function of time was recorded, while the temp. of the soln. was increased linearly with time at a rate of 20°/h using a Lauda PM-350 programmer and a Lauda RCS 6 bath equipped with a R 22 unit (MWG Lauda, FRG). The actual temp. was measured in the sample cell with a Pt resistor.

HPLC Separation. HPLC was carried out on a 4 × 250 and 4 × 25 mm (10 μm) RP-18 LiChrosorb column (Merck) using a Merck-Hitachi HPLC apparatus with one pump (model 655A-12) connected with a proportion-

ing valve, a variable wavelength monitor (model 655A), and a controller (model L-5000), connected with an integrator (model D-2000). The solvent gradients consisting of 0.1M Et₃NHOAc (pH 7) MeCN 95:5 (A), and MeCN (B) were used in the following order: gradient I, 15 min 20–35% B in A; II, 10 min 5–20% B in A; III, 12 min 100% A, 13 min 0–10% B in A; IV, 8 min 100% B, 4 min 0–16% B in A.

Phase-Transfer Glycosylation of 4-(Methylamino)-7H-pyrrolo[2,3-d]pyrimidine (5a) with 2-Deoxy-3,5-di-O-(p-toluoyl)- α -D-erythro-pentofuranosyl Chloride (6). Powdered KOH (700 mg, 12.5 mmol), **5a** [28] (700 mg, 4.72 mmol), and tris[2-(2-methoxyethoxy)ethyl]amine (TDA-1; 100 μ l, 0.31 mmol) were stirred at r.t. in anh. MeCN (50 ml). After 5 min, **6** [29] (1.85 g, 4.77 mmol) was added. Stirring was continued for 10 min, insoluble material removed by filtration and washed with MeCN, the filtrate evaporated, and the residue chromatographed on silica gel 60 (10 $\times$ 5 cm column): 3 zones were separated, containing **7a**, **9a**, and **9b**, respectively.

7-[2'-Deoxy-3',5'-di-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-7H-pyrrolo[2,3-d]pyrimidine (7a). From the fast migrating zone (eluant G), **7a** was isolated as colorless foam (1.56 g, 66%). TLC (silica gel, H): R_f 0.5. UV (MeOH): 240 (35500), 271 (18000). ¹H-NMR ((D₆)DMSO): 2.38, 2.40 (2s, 2 CH₃); 2.65 (m, H β -C(2')); 2.96 (d, J = 4.6, CH₃N); 3.07 (m, H α -C(2')); 4.58 (m, H-C(4')); 5.72 (m, H-C(3')); 6.61 (d, J = 3.6, H-C(5)); 6.67 (m, H-C(1')); 7.32, 7.93 (m, 8 arom. H, H-C(6)); 7.55 (q, J = 4.6, NH); 8.17 (s, H-C(2)). Anal. calc. for C₂₈H₂₈N₄O₅: C 67.19, H 5.63, N 11.19; found: C 67.00, H 5.67, N 11.01.

7-[2'-Deoxy-3'-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-7H-pyrrolo[2,3-d]pyrimidine (9a). From the 2nd zone (eluant H), a colorless foam was obtained (110 mg, 6%). TLC (silica gel, H): R_f 0.4. UV (MeOH): 239 (19100), 273 (15200). ¹H-NMR ((D₆)DMSO): 2.41 (s, CH₃); 2.89 (m, H α -C(2')); H β -C(2') superimposed by DMSO; 2.96 (d, J = 4.6, CH₃N); 3.68 (m, 2 H-C(5')); 4.19 (m, H-C(4')); 5.51 (t, J = 5.7, OH-C(5')); 5.58 (m, H-C(3')); 6.60 (m, H-C(1'), H-C(1'), H-C(5)); 7.37, 7.95 (m, 4 arom. H); 7.44 (d, J = 3.7, H-C(6)); 7.58 (q, J = 4.6, NH); 8.16 (s, H-C(2)). Anal. calc. for C₂₀H₂₂N₄O₄: C 62.81, H 5.80, N 14.65; found: C 62.54, H 5.77, N 14.28.

7-[2'-Deoxy-5'-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-7H-pyrrolo[2,3-d]pyrimidine (9b). The slow migrating zone (eluant H) afforded a colorless foam (100 mg, 6%). TLC (silica gel, I): R_f 0.8. UV (MeOH): 236 (17900), 274 (15100). ¹H-NMR ((D₆)DMSO): 2.29 (m, H β -C(2')); 2.38 (s, CH₃); 2.64 (m, H α -C(2')); 2.95 (d, J = 3.5, CH₃N); 4.10 (m, H-C(4')); 4.35, 4.49 (m, 2 H-C(5'), H-C(3')); 5.53 (d, J = 4.3, OH-C(3')); 6.56 (m, H-C(1')); 7.27 (d, J = 3.7, H-C(6)); 7.33, 7.86 (m, 4 arom. H); 7.52 (q, J = 3.7, NH); 8.16 (s, H-C(2)). Anal. calc. for C₂₀H₂₂N₄O₄: C 62.81, H 5.80, N 14.64; found: C 62.54, H 5.70, N 14.40.

7-(2'-Deoxy- β -D-erythro-pentofuranosyl)-4-(methylamino)-7H-pyrrolo[2,3-d]pyrimidine (2). From **7a**: A soln. of **7a** (100 mg, 0.2 mmol) in MeOH (25 ml, saturated with NH₃ at 0°) was stirred for 6 h at r.t. and evaporated. The residue was adsorbed on silica gel 60H (2.0 g) and chromatographed on a silica-gel 60H column (10 $\times$ 2 cm, H): **2** (40 mg, 76%), colorless foam.

From 7c: A soln. of 4-chloro-7-[2'-deoxy-3',5'-di-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-7H-pyrrolo[2,3-d]pyrimidine [11] (**7c**; 1.0 g, 1.98 mmol) MeOH (100 ml) and 40% aq. MeNH₂ (10 ml) was treated in a steel bomb for 6 h at 150°. The cold mixture was evaporated, the residue dissolved in MeOH, adsorbed on silica gel (2.0 g), and chromatographed on a silica gel column (5 $\times$ 2 cm, H): the main zone yielded **2** as colorless foam (340 mg, 65%).

From 9d: A soln. of **9d** (see below; 150 mg, 0.27 mmol) in 40% aq. MeNH₂/dioxane 1:1 (40 ml) was stored at 60° for 4 h in a pressure bottle. After evaporation of the solvent, the residue was chromatographed on silica gel (column: 4 $\times$ 3 cm, G). From the main zone, colorless amorphous **2** (60 mg, 83%) was isolated. TLC (silica gel, I): R_f 0.4. UV (MeOH): 210 (30300), 274 (15600). ¹H-NMR ((D₆)DMSO): 2.15 (m, H β -C(2')); 2.95 (d, J = 4.6, CH₃N); 3.54 (m, 2 H-C(5')); 3.83 (m, H-C(4')); 4.35 (m, H-C(3')); 5.18 (m, OH-C(5')); 5.27 (m, OH-C(3')); 6.49 (m, H-C(1')); 6.57 (d, J = 3.6, H-C(5)); 7.34 (d, J = 3.6, H-C(6)); 7.52 (q, J = 4.6, NH); 8.14 (s, H-C(2)). Anal. calc. for C₁₂H₁₆N₄O₃: C 54.54, H 6.10, N 21.20; found: C 54.61, H 6.11, N 20.95.

7-(2'-Deoxy- β -D-erythro-pentofuranosyl)-4-(ethylthio)-7H-pyrrolo[2,3-d]pyrimidine (8c). A mixture of **8b** (400 mg, 1.48 mmol) [11] [30], MeOH (5 ml), ethanethiol (6 ml), and 20% aq. NaOH soln. (2 ml) was stirred for 1 h at r.t. Then the soln. was neutralized with AcOH and evaporated. The residue was suspended in CHCl₃/MeOH 9:1, adsorbed on silica gel (2.0 g), and chromatographed on a silica-gel column (20 $\times$ 2 cm, G). The main zone gave a colorless foam upon evaporation which was crystallized from acetone: **8c** as colorless needles (410 mg, 94%). M.p. 140°. TLC (silica gel, H): R_f 0.5. UV (MeOH): 222 (24000), 250 (7000), 293 (14000). ¹H-NMR ((D₆)DMSO): 1.35 (t, J = 7.3, CH₃CH₂S); 2.23 (m, H β -C(2')); 2.52 (m, H α -C(2')); 3.32 (q, J = 7.3, CH₃CH₂S); 3.55 (m, 2 H-C(5')); 3.85 (m, H-C(4')); 4.38 (m, H-C(3')); 5.00 (t, J = 5.3, OH-C(5')); 5.32 (d, J = 3.8, OH-C(3')); 6.55 (d, J = 3.7, H-C(5)); 6.62 (m, H-C(1')); 7.75 (d, J = 3.7, H-C(6)); 8.62 (s, H-C(2)). Anal. calc. for C₁₃H₁₇N₃O₃S: C 52.87, H 5.80, N 14.23, S 10.86; found: C 52.80, H 5.83, N 14.12, S 11.08.

Oxidation of 8c with 3-Chloroperbenzoic Acid. To a soln. of **8c** (400 mg, 1.36 mmol) in EtOH (40 ml), 3-chloroperbenzoic acid (300 mg; technical grade, 80%) was added. After stirring for 30 min at r.t., the soln. was

evaporated and the residue chromatographed on silica gel 60H (column: 10 × 5 cm, G). The diastereoisomeric mixture **8d/8d'** was isolated from the slower migrating main zone as a colorless foam (250 mg, 59%). TLC (silica gel, H): R_f 0.3. The faster migrating zone yielded **8e** as a colorless foam (100 mg, 22%). TLC (silica gel, H): R_f 0.3.

7-(2'-Deoxy- β -D-erythro-pentofuranosyl)-4-(ethylsulfinyl)-7H-pyrrolo[2,3-d]pyrimidine (**8d/8d'**). UV (MeOH): 228 (21000), 286 (4000), 305 (4200). $^1\text{H-NMR}$ ((D_6) DMSO): 1.10 (t, $J = 7.3$, CH_3CH_2); 2.29 (m, $\text{H}_\beta\text{-C}(2')$); 2.56 (m, $\text{H}_\alpha\text{-C}(2')$); 3.11, 3.31 (m, CH_3CH_2); 3.56 (m, 2 H-C(5')); 3.87 (m, H-C(4')); 4.40 (m, H-C(3')), 4.97 (t, $J = 5.4$, OH-C(5')); 5.36 (d, $J = 4.1$, OH-C(3')); 6.73 (t, $J = 6.8$, H-C(1')); 7.12 (d, $J = 3.7$, H-C(5)); 8.01 (d, $J = 3.7$, H-C(6)); 8.86 (s, H-C(2))².

7-(2'-Deoxy- β -D-erythro-pentofuranosyl)-4-(ethylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidine (**8e**). UV (MeOH): 228 (26900), 285 (4200). $^1\text{H-NMR}$ ((D_6) DMSO): 1.19 (t, $J = 7.4$, CH_3CH_2); 2.32 (m, $\text{H}_\beta\text{-C}(2')$); 2.58 (m, $\text{H}_\alpha\text{-C}(2')$); 3.57 (m, CH_3CH_2 , 2 H-C(5')); 3.89 (m, H-C(4')); 4.42 (m, H-C(3')); 4.99 (t, $J = 6.5$, OH-C(5')); 5.37 (d, $J = 4.2$, OH-C(3')); 6.75 (t, $J = 7.5$, H-C(1')); 7.00 (d, $J = 3.7$, H-C(5)); 8.20 (d, $J = 3.7$, H-C(6)); 9.04 (s, H-C(2))².

7-[2'-Deoxy-3'-5'-di-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-4-(methylthio)-7H-pyrrolo[2,3-d]pyrimidine (**7b**). Glycosylation was carried out as described for **7a** by using powdered KOH (650 mg, 11.6 mmol), TDA-1 (100 μ l, 0.31 mmol), 4-(methylthio)-7H-pyrrolo[2,3-d]pyrimidine [**31**] (**5b**; 500 mg, 3.03 mmol), and **6** (1200 mg, 3.10 mmol). After chromatographic purification (F), colorless crystals (1.26 g, 80%) were obtained from EtOH. M.p. 118°. TLC (silica gel, K): R_f 0.7. UV (MeOH): 223 (38300), 240 (39200), 291 (14300). $^1\text{H-NMR}$ ((D_6) DMSO): 2.37, 2.40 (2s, CH_3); 2.65 (s, CH_3S); 2.73 (m, $\text{H}_\beta\text{-C}(2')$); 3.15 (m, $\text{H}_\alpha\text{-C}(2')$); 4.44 (m, H-C(4')), 2 H-C(5')); 5.75 (m, H-C(3')); 6.61 (d, $J = 3.7$, H-C(5)); 6.75 (m, H-C(1')); 7.34, 7.91 (m, 8 arom. H); 7.74 (d, $J = 3.7$, H-C(6)); 8.63 (s, H-C(2)). Anal. calc. for $\text{C}_{28}\text{H}_{27}\text{N}_3\text{O}_5\text{S}$: C 64.98, H 5.25, N 8.12, S 6.19; found: C 65.03, H 5.28, N 7.98, S 6.12.

7-[2'-Deoxy-3'-5'-di-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-4-(methylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidine (**9d**). By warming, **7b** (1000 mg, 1.93 mmol) was dissolved in MeOH (50 ml) and 3-chloroperbenzoic acid (1.6 g; technical grade, 80%) was added and stirred for 1 h at r.t. Upon cooling, a precipitate was formed which was recrystallized from MeOH. Colorless crystals of **9d** (860 mg, 81%). M.p. 143°. TLC (silica gel, K): R_f 0.5. UV (MeOH): 232 (49500). $^1\text{H-NMR}$ ((D_6) DMSO): 2.38, 2.41 (2s, 2 CH_3); 2.80 (m, $\text{H}_\beta\text{-C}(2')$); 3.20 (m, $\text{H}_\alpha\text{-C}(2')$); 3.42 (s, CH_3SO_2); 4.61 (m, H-C(4')), 2 H-C(5')); 5.78 (m, H-C(3')); 6.87 (m, H-C(1')); 7.01 (d, $J = 3.8$, H-C(5)); 7.35, 7.92 (m, 8 arom. H); 8.19 (d, $J = 3.8$, H-C(6)); 9.02 (s, H-C(2)). Anal. calc. for $\text{C}_{28}\text{H}_{27}\text{N}_3\text{O}_7\text{S}$: C 61.19, H 4.95, N 7.65, S 5.83; found: C 61.25, H 5.00, N 7.58, S 5.82.

7-(2'-Deoxy- β -D-erythro-pentofuranosyl)-4-(methylthio)-7H-pyrrolo[2,3-d]pyrimidine (**8a**). As described for **7a** ($\rightarrow$ 2), **7b** (500 mg, 0.97 mmol) was deprotected. The soln. was evaporated, the residue adsorbed on silica gel 60 (2.0 g) and chromatographed on a silica-gel 60H column (10 × 3 cm, G). The residue of the main zone was crystallized from acetone. Colorless crystals (250 mg, 92%). M.p. 148°. TLC (silica gel, H): R_f 0.3. UV (MeOH): 222 (28000), 249 (8300), 291 (16000). $^1\text{H-NMR}$ ((D_6) DMSO): 2.23 (m, $\text{H}_\beta\text{-C}(2')$); 2.54 (m, $\text{H}_\alpha\text{-C}(2')$); 2.65 (s, CH_3S); 3.56 (m, 2 H-C(5')); 3.85 (m, H-C(4')); 4.38 (m, H-C(3')); 5.00 (t, $J = 5.5$, OH-C(5')); 5.32 (d, $J = 4.1$, OH-C(3')); 6.58 (d, $J = 3.8$, H-C(5)); 6.62 (m, H-C(1')); 7.76 (d, $J = 3.8$, H-C(6)); 8.64 (s, H-C(2)). Anal. calc. for $\text{C}_{12}\text{H}_{15}\text{N}_3\text{O}_5\text{S}$: C 51.23, H 5.37, N 14.94, S 11.40; found: C 51.20, H 5.43, N 14.73, S 11.36.

1-[2'-Deoxy- β -D-erythro-pentofuranosyl]-4-(methylamino)-1H-pyrazolo[3,4-d]pyrimidine (**3**). To a soln. of 1-[2'-deoxy-3,5-di-O-(p-toluoyl)- β -D-erythro-pentofuranosyl]-4-methoxy-1H-pyrazolo[3,4-d]pyrimidine [**12a**, b] (**10a**; 500 mg, 1 mmol) in dry MeOH (60 ml), 33% $\text{MeNH}_2/\text{EtOH}$ (6 ml) was added and the mixture stored at 60° for 12 h. After evaporation, the residue was separated on a 15 × 4 cm column (silica gel 60H, A) by flash chromatography. Isolation of the material of the main zone yielded colorless, amorphous **3** (250 mg, 95%). TLC (silica gel, A): R_f 0.3. UV (MeOH): 281 (12800), 264 (sh) (9050). $^1\text{H-NMR}$ ((D_6) DMSO): 8.33 (q, $J = 4.5$, NH); 8.29 (s, H-C(6)); 8.14 (s, H-C(3)); 6.55 (t, $J = 6.4$, H-C(1')); 5.27 (d, $J = 4.5$, OH-C(3')); 4.83 (t, $J = 5.8$, OH-C(5')); 4.44 (m, H-C(3')); 3.81 (m, H-C(4')); 3.42 (m, 2 H-C(5')); 2.98 (d, $J = 4.4$, CH_3N); 2.80 (m, $\text{H}_\alpha\text{-C}(2')$); 2.24 (m, $\text{H}_\beta\text{-C}(2')$). Anal. calc. for $\text{C}_{11}\text{H}_{15}\text{N}_5\text{O}_3$: C 49.81, H 5.70, N 26.40; found: C 49.69, H 5.69, N 26.49.

2-(2'-Deoxy- β -D-erythro-pentofuranosyl)-4-(methylamino)-2H-pyrazolo[3,4-d]pyrimidine (**4**). As for **3**, using **11a** (1000 mg, 2 mmol), 33% $\text{MeNH}_2/\text{EtOH}$ (10 ml), and dry MeOH (80 ml): colorless amorphous **4** (510 mg, 95%). TLC (silica gel, B): R_f 0.4. UV (MeOH): 296 (13200). $^1\text{H-NMR}$ ((D_6) DMSO): 8.48 (s, H-C(3)); 8.27 (q, $J = 4.8$, NH); 8.23 (s, H-C(6)); 6.31 (t, $J = 5.9$, H-C(1')); 5.33 (d, $J = 4.3$, OH-C(3')); 4.90 (t, $J = 5.6$, OH-C(5')); 4.39 (m, H-C(3')); 3.90 (m, H-C(4')); 3.52 (m, 2 H-C(5')); 2.96 (d, $J = 4.6$, CH_3N); 2.61 (m, $\text{H}_\alpha\text{-C}(2')$); 2.36 (m, $\text{H}_\beta\text{-C}(2')$). Anal. calc. for $\text{C}_{11}\text{H}_{15}\text{N}_5\text{O}_3$: C 49.80, H 5.70, N 26.40; found: C 49.56, H 5.72, N 26.37.

²) No sufficient elemental analyses data were obtained due to decomposition of the materials on drying.

1-[2'-Deoxy-5'-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-1H-pyrazolo[3,4-d]-pyrimidine (**10b**). By co-evaporation with anhyd. pyridine (10 ml), **3** (400 mg, 1.5 mmol) was dried. Then, 4-(dimethylamino)pyridine (90 mg, 0.7 mmol) and 4,4'-dimethoxytrityl chloride (900 mg, 2.7 mmol) were added and stirred for 1 h under N₂ at r.t. (TLC control; silica gel, C). To the soln., 5% aq. NaHCO₃ soln. (50 ml) was added and the mixture extracted with CH₂Cl₂ (3 $\times$ 100 ml). The combined org. layers were dried (Na₂SO₄), filtered, and evaporated. The residue was flash chromatographed (9 $\times$ 3 cm column, silica gel, C). Isolation of the material of the main zone yielded colorless, amorphous **10b** (700 mg, 82%). TLC (silica gel, A): R_f 0.66. UV (MeOH): 281 (13950), 231 (23650). ¹H-NMR ((D₆)DMSO): 8.32 (s, H-C(6)); 8.29 (m, NH); 8.08 (s, H-C(3)); 6.59 (m, H-C(1')); 5.31 (d, J = 5, OH-C(3')); 4.57 (m, H-C(3')); 3.94 (m, H-C(4')); 3.69, 3.70 (2s, 2 CH₃O); 3.05 (m, 2 H-C(5')); 2.98 (d, J = 4.3, CH₃N); 2.79 (m, H_x-C(2')); 2.31 (m, H_β-C(2')). Anal. calc. for C₃₂H₃₃N₅O₅: C 67.71, H 5.86, N 12.33; found: C 67.60, H 5.98, N 12.16.

2-[2'-Deoxy-5'-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-1H-pyrazolo[3,4-d]-pyrimidine (**11b**). As for **10b**, using **4** (500 mg, 1.9 mmol), 4,4'-dimethoxytrityl chloride (1.1 g, 3.3 mmol), and 4-(dimethylamino)pyridine (110 mg, 0.85 mmol): colorless amorphous **11b** (750 mg, 70%). TLC (silica gel, D): R_f 0.3. UV (MeOH): 297 (12650), 234 (24000). ¹H-NMR ((D₆)DMSO): 8.50 (s, H-C(3)); 8.27 (m, NH); 8.24 (s, H-C(6)); 6.38 (m, H-C(1')); 5.38 (d, J = 4.9, OH-C(3')); 4.48 (m, H-C(3')); 3.98 (m, H-C(4')); 3.70, 3.67 (2s, 2 CH₃O); 3.09 (m, 2 H-C(5')); 2.98 (d, J = 4.5, CH₃N); 2.71 (m, H_x-C(2')); 2.36 (m, H_β-C(2')). Anal. calc. for C₃₂H₃₃N₅O₅: C 67.71, H 5.86, N 12.33; found: C 67.60, H 6.05, N 12.32.

7-[2'-Deoxy-5'-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-1H-pyrrolo[2,3-d]-pyrimidine (**9c**). As for **10b**, using **2** (310 mg, 1.2 mmol), 4,4'-dimethoxytrityl chloride (680 mg, 2 mmol), 4-(dimethylamino)pyridine (75 mg, 0.6 mmol), and anhyd. pyridine: **9c** (550 mg, 81%), colorless, amorphous material. TLC (silica gel, A): R_f 0.6. UV (MeOH): 275 (17000). ¹H-NMR ((D₆)DMSO): 8.14 (s, H-C(2)); 7.48 (m, NH); 6.53 (m, H-C(1')); 5.33 (d, J = 4.3, OH-C(3')); 4.35 (m, H-C(3')); 5.91 (m, H-C(4')); 3.72 (2s, 2 CH₃O); 3.14 (m, 2 H-C(5')); 2.95 (d, J = 4.6, CH₃N); 2.6 (m, H_x-C(2')); 2.23 (m, H_β-C(2')). Anal. calc. for C₃₃H₃₄N₄O₆: C 69.95, H 6.05, N 9.88; found: C 70.05, H 6.12, N 9.90.

General Procedure for the Preparation of the Phosphoramidites 12–14. The 5'-tritylated nucleoside was dissolved in anhyd. MeCN (5 ml) in a flask preflushed with Ar. Chloro(diisopropylamino)methoxyphosphine and (i-Pr)₂EtN were added by syringe to the soln. kept under Ar at r.t. After 30 min, the soln. was added to 5% aq. NaHCO₃ (30 ml) and extracted 3 times with CH₂Cl₂. The org. layer was dried (Na₂SO₄) and evaporated and the foam purified by flash chromatography on silica gel 60H (column 12 $\times$ 2.5 cm, E) affording the colorless, amorphous phosphoramidite.

1-[2'-Deoxy-5'-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-1H-pyrazolo[3,4-d]-pyrimidine 3'-[Methyl N,N-Diisopropylphosphoramidite] (**12**). From **10b** (400 mg, 0.7 mmol), chloro-(diisopropylamino)methoxyphosphine (0.14 ml, 0.7 mmol), and (i-Pr)₂EtN (0.36 ml, 2.1 mmol): colorless, amorphous **1** (350 mg, 69%). TLC (silica gel, E): R_f 0.65. ³¹P-NMR (CDCl₃): 147.08.

2-[2'-Deoxy-5'-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-1H-pyrazolo[3,4-d]-pyrimidine 3'-[Methyl N,N-Diisopropylphosphoramidite] (**13**). As for **12**: **13** (370 mg, 71%) as colorless foam. TLC (silica gel, E): R_f 0.35. ³¹P-NMR (CDCl₃): 148.4.

7-[2'-Deoxy-5'-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4-(methylamino)-7H-pyrrolo[2,3-d]-pyrimidine 3'-[Methyl N,N-Diisopropylphosphoramidite] (**14**). As described for **12**: colorless, amorphous **14** (310 mg, 61%). TLC (silica gel, E): R_f 0.7. ³¹P-NMR (CDCl₃): 147.0, 147.2.

Solid-Phase Synthesis of the Oligomers 19–27. The synthesis of the oligonucleotides was accomplished on a 1- μ mol scale using the methyl phosphoramidites of [(MeO)₂Tr]j**bz**⁶A_d, [(MeO)₂Tr]j**bz**²G_d, [(MeO)₂Tr]j**bz**⁴C_d, and [(MeO)₂Tr]j**d as well as the phosphoramidites **12–15**. The synthesis of **19–27** followed the regular protocol of the DNA synthesizer for methyl phosphoramidites [16]. Deprotection of NH₂ groups was carried out with 25% aq. NH₃ at 60° for 48 h. The 4,4'-dimethoxytrityl residues of **19a–27a** were removed by treatment with 80% aq. CH₃CO₂H, for 5 min at r.t. Purification was accomplished by HPLC on RP-18 columns using solvent system I for the (MeO)₂Tr derivatives **19a–27a**, and II for the detritylated oligonucleotides **19a–27b**. The oligomers were desalted on a 25 $\times$ 4 HPLC cartridge (RP-18 silica gel). Inorg. material was eluted with H₂O (10 ml), while the oligomer was eluted with MeOH/H₂O 3:2 (5 ml). The oligomers were lyophilized on a Speed-Vac evaporator to yield a colorless foam which was dissolved in H₂O (1 ml) and stored frozen at –20°.**

Enzymatic Hydrolysis of the Oligomers 19b–27b. The oligomer (0.4 A₂₆₀ units) was dissolved in 0.1M Tris-HCl buffer (pH 8.5; 200 μ l) and treated with snake-venom phosphodiesterase (3 μ g) at 25° for 60 min and alkaline phosphatase (2 μ g) for 15 min at 25°. The mixture was analyzed on reverse-phase HPLC (RP-18, solvent system III or IV). Quantification of the material was made at 260 nm on the basis of the peak areas which were divided by the

extinction coefficients of the nucleoside constituents (ϵ_{260} : A_d, 15 400; C_d, 7300; G_d, 11 700; T_d, 8800; m⁶A_d, 11 600; m⁶c⁷A_d, 9700; c⁷m⁶z⁸A_d, 8400; c⁷m⁶z⁸A'_d, 5300).

Phosphodiester Hydrolysis of the Oligomers 23b–27b with the Endodeoxyribonuclease Dpn I. The oligomers **23b–27b** (16 μ M) were dissolved in 6 mM Tris-HCl buffer (pH 7.7; 140 μ l) containing 150 mM NaCl, 6 mM MgCl₂, 7 mM 2-mercaptoethanol, and bovine serum albumine (100 μ g/ml) and treated with the endodeoxyribonuclease Dpn I (10 units) at 25°. Samples of 20 μ l were taken at different intervals of time and analyzed on reverse-phase HPLC (RP-18, gradient II). The hydrolysis products were collected, lyophilized, and analyzed after hydrolysis with snake-venom phosphodiesterase and alkaline phosphatase, as described in the previous section.

REFERENCES

- [1] W. Doerfler, *Angew. Chem.* **1984**, 96, 917.
- [2] P. Modrich, R. J. Roberts, in 'Nucleases', Eds. S. M. Linn and R. J. Roberts, Cold Spring Harbor Laboratory, New York, 1982.
- [3] J. D. Engel, P. H. von Hippel, *J. Biol. Chem.* **1978**, 253, 927.
- [4] A. M. Delort, A. Guy, D. Molko, R. Teoule, *Nucleos. Nucleot.* **1985**, 4, 201.
- [5] T. Tanaka, S. Tamatsukuri, M. Ikehara, *Chem. Pharm. Bull.* **1986**, 34, 2044.
- [6] A. Ono, T. Ueda, *Nucleic Acids Res.* **1987**, 15, 219.
- [7] J. F. Gerster, B. Carpenter, R. K. Robins, L. B. Townsend, *J. Am. Chem. Soc.* **1967**, 10, 326.
- [8] H. D. Winkler, F. Seela, *J. Org. Chem.* **1983**, 26, 3119.
- [9] F. Seela, B. Westermann, U. Bindig, *J. Chem. Soc., Perkin Trans. I* **1988**, 697.
- [10] H. Rosemeyer, G. Toth, F. Seela, *Nucleos. Nucleot.* **1989**, in press.
- [11] F. Seela, H.-P. Muth, U. Bindig, *Synthesis* **1988**, 9, 670.
- [12] a) F. Seela, H. Steker, *Helv. Chim. Acta* **1985**, 68, 563; b) *ibid.* **1988**, 71, 1813.
- [13] F. Seela, W. Herdering, A. Kehne, *Helv. Chim. Acta* **1987**, 70, 1649.
- [14] M. D. Matteucci, M. H. Caruthers, *J. Am. Chem. Soc.* **1981**, 103, 3185.
- [15] L. J. McBride, M. H. Caruthers, *Tetrahedron Lett.* **1983**, 24, 245.
- [16] *Applied Biosystems*, users manual of the DNA synthesizer 380B.
- [17] J. D. Engel, P. H. von Hippel, *Biochemistry* **1974**, 13, 4143.
- [18] R. E. Dickerson, H. R. Drew, B. N. Conner, R. M. Wing, A. V. Fratini, M. L. Kopka, *Science* **1982**, 216, 475.
- [19] I. Tinoco, C. R. Cantor, *Meth. Biochem. Anal.* **1970**, 18, 81.
- [20] B. Bourtayre, J. Liquier, L. Pizzorni, E. Taillandier, *J. Biol. Struct. Dyn.* **1987**, 5, 97.
- [21] A. H.-J. Wang, G. J. Quigley, F. J. Kolpak, J. L. Crawford, J. H. van Boom, G. van der Marel, A. Rich, *Nature (London)* **1979**, 282, 680.
- [22] F. M. Pohl, T. M. Jovin, *J. Mol. Biol.* **1972**, 67, 375.
- [23] W. Saenger, in 'Principles of Nucleic Acid Structures', Ed. C. R. Cantor, Springer Verlag, New York, 1984.
- [24] S. Lacks, B. Greenberg, *J. Biol. Chem.* **1975**, 250, 4060.
- [25] G. E. Geier, P. Modrich, *J. Biol. Chem.* **1979**, 254, 1408.
- [26] S. Lacks, B. Greenberg, *J. Mol. Biol.* **1977**, 114, 153.
- [27] M. McClelland, L. G. Kessler, M. Bittner, *Proc. Natl. Acad. Sci. USA* **1984**, 81, 983.
- [28] G. H. Hitchings, K. W. Ledig, R. A. West, to *Burroughs Wellcome & Co.*, 1962, U.S. Patent 3037980.
- [29] M. Hoffer, *Chem. Ber.* **1960**, 93, 2777.
- [30] Z. Kazimierzczuk, H. B. Cottam, G. R. Revankar, R. K. Robins, *J. Am. Chem. Soc.* **1984**, 106, 6379.
- [31] F. Seela, H. Steker, *Liebigs Ann. Chem.* **1983**, 1576.