

Formation and structure of $[\text{Co}\{\text{Ph}_2\text{PN}(\text{iBu})\text{PPh}_2\text{-}P,P'\}_2(\text{CO})][\text{Co}(\text{CO})_4]$ – a cage molecule-pair or an ion-pair complex?

Zheng-Zhi Zhang, Ao Yu and Hua-Ping Xi

Elemento-Organic Chemistry Laboratory, Nankai University, 300071 Tianjin (China)

Ru-Ji Wang and Hong-Gen Wang

Central Laboratory, Nankai University, 300071 Tianjin (China)

(Received April 28, 1993; in revised form September 24, 1993)

Abstract

The strong π -acid ligand $\text{Ph}_2\text{PN}(\text{iBu})\text{PPh}_2$ reacts with $\text{Co}_2(\text{CO})_8$ (1:1) to give $\text{Co}_2[\mu\text{-Ph}_2\text{PN}(\text{iBu})\text{PPh}_2](\mu\text{-CO})_2(\text{CO})_4$ (1); however, when the ratio is 2:1 a novel species $[\text{Co}\{\text{Ph}_2\text{PN}(\text{iBu})\text{PPh}_2\text{-}P,P'\}_2(\text{CO})][\text{Co}(\text{CO})_4]$ (2) has been obtained. Crystal data for 2: $M_r = 1140.83$; triclinic, space group $P\bar{1}$, $a = 12.330(2)$, $b = 13.340(2)$, $c = 18.122(3)$ Å, $\alpha = 86.63(1)$, $\beta = 80.75(1)$, $\gamma = 84.24(1)^\circ$, $V = 2924$ Å³, $Z = 2$; $R = 0.060$ for 3711 reflections having $I > 3\sigma(I)$. The results of X-ray diffraction, ESR, variable-temperature magnetic susceptibility, conductivity, and XPS analysis support that the species 2 is a d^9 - d^9 cage molecule-pair. The mechanism for the formation of the species 2 has been investigated initially by ^{31}P NMR.

Key words: Cobalt; Cage compound; Phosphazane; Carbonyl

1. Introduction

A variety of organometallic complexes containing diphosphino methane ($\text{X}_2\text{PCH}_2\text{PX}_2$) has been investigated extensively for more than two decades because of their potential as catalysts and their novel structural and reactive features [1–3]. Interest in the analogous diphosphino amine ligands $[\text{X}_2\text{PN}(\text{R})\text{PX}_2]$ has grown rapidly in recent years [4–11]. It has been found that small structural changes in these ligands produce large effects on aspects of the synthesis of metal complexes.

In the present work, $\text{Co}_2(\text{CO})_8$ is found to react with $\text{Ph}_2\text{PN}(\text{iBu})\text{PPh}_2$ in different ratios to give different products. When the ratio is 1:1, a predicted product, $\text{Co}_2[\mu\text{-Ph}_2\text{PN}(\text{iBu})\text{PPh}_2](\mu\text{-CO})_2(\text{CO})_4$ (1), is obtained. However, with the ratio 1:2, a novel product, $[\text{Co}\{\text{Ph}_2\text{PN}(\text{iBu})\text{PPh}_2\text{-}P,P'\}_2(\text{CO})][\text{Co}(\text{CO})_4]$ (2), has been obtained and there is evidence for it being a cage molecule-pair. The mechanism for the formation of 2

has been investigated initially by ^{31}P NMR, and the crystal structure of 2 has been studied by X-ray diffraction.

2. Experimental details

2.1. Reagents and solvents

$\text{Co}_2(\text{CO})_8$ (Strem) was used without further purification. All solvents were reagent grade and were purified by general procedures.

2.2. Physical measurements

Microanalyses for C, H and N were carried out with a Perkin-Elmer analyzer model 240. Infrared spectra (KBr disc) were recorded with a WFD-14 spectrophotometer. An AC-P200 NMR spectrometer was used to record ^{31}P spectra at 81.03 Hz (CDCl_3 , external standard: H_3PO_4). ESR spectrum was measured with a JES-FEIXG apparatus using the X-band. Conductivity was measured in acetonitrile with a DDS-11A conductometer. Variable-temperature magnetic susceptibility was carried out with a Vibrating-Sample Magnetic

Correspondence to: Dr. Z.-Z. Zhang.

meter model CF. Diamagnetic corrections were made with Pascal's constants. The magnetic moments were calculated by the equation $\mu_{\text{eff}} = 2.828(\chi T)^{1/2}$. The XPS spectra (Mg K α) were recorded on the PHI 5300 ESCA system (Perkin-Elmer) at 250 VA (12.5 kV $\times$ 20 mA) calibrated by assuming the binding energy of the adventitious carbon to be 284.6 eV, with experimental error ± 0.2 eV.

2.3. Synthesis of the compounds

All preparations were carried out under N₂ using Schlenk-type glassware.

2.3.1. Ph₂PN(ⁱBu)PPh₂

A solution of Ph₂PCl (15 g, 0.067 mmol) in benzene (25 ml) was added to a stirred solution of isobutylamine (12.4 g, 0.17 mmol) in 25 ml of benzene at -5°C , and stirring was continued for 1 h. The amine hydrochloride was filtered off and washed with benzene. The filtrate was passed through a silica column (100–140 mesh), then a solution of triethylamine (6.8 g, 0.067 mol) in 25 ml of benzene was added at 0°C and this was followed by dropwise addition of Ph₂PCl (15 g, 0.067 mol) in benzene (25 ml). Stirring was continued for 10 h and the precipitate was removed by filtration. The filtrate was passed through a silica column, then the volume of the solvent was reduced and ethanol was added to give the product (16 g, 54%); M.p. 108–109°C. Anal. Calcd for C₂₈H₂₉NP₂: C, 76.20; H, 6.62; N, 3.17. Found: C, 76.18; H, 6.60; N, 3.02%. ³¹P NMR (CDCl₃) δ 62.73 ppm. FD-MS (m/z): 441 (M⁺, 100%).

2.3.2. Co₂[μ -Ph₂PN(ⁱBu)PPh₂](μ -CO)₂(CO)₄ (1)

A solution of Co₂(CO)₈ (0.35 g, 1.02 mmol) and Ph₂PN(ⁱBu)PPh₂ (0.45 g, 1.02 mmol) in dichloromethane (20 ml) was stirred for 24 h at room temperature. The volume of the solution was reduced to about 5 ml, then ether (15 ml) was added to give an orange precipitate. The crude product was recrystallized from CH₂Cl₂/hexane. Yield: 0.4 g (52%), M.p. 115°C (dec). Anal. Calcd for C₃₄H₂₉Co₂NO₆P₂: C, 56.12; H, 3.99; N, 1.92. Found: C, 56.54; H, 3.53; N, 1.84%. IR: $\nu(\text{CO})$, 2040s, 1995s, 1986s, 1976s, 1814w, 1790s cm⁻¹. ³¹P NMR (CDCl₃) δ 79.15 ppm.

2.3.3. [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)][Co(CO)₄] (2)

A solution of Co₂(CO)₈ (0.33 g, 0.95 mmol) in THF (30 ml) was added dropwise to a solution of Ph₂PN(ⁱBu)PPh₂ (0.88 g, 2.00 mmol) in THF (25 ml) at 0°C . Immediately CO gas was evolved, then the solution changed from pale yellow to dark red. After completion of the addition, stirring was continued for 1

TABLE 1. Experimental details of the X-ray diffraction study of [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)][Co(CO)₄] (2)

Formula	C ₆₁ H ₅₈ Co ₂ N ₂ O ₅ P ₄
Molecular weight	1140.83
Colour	red
Crystal system	triclinic
Space group	$P\bar{1}$
<i>a</i> , Å	12.330(2)
<i>b</i> , Å	13.340(2)
<i>c</i> , Å	18.122(3)
α , °	86.63(1)
β , °	80.75(1)
γ , °	84.24(1)
<i>V</i> , Å ³	2924.3
<i>Z</i>	2
<i>D</i> _{calcd} , g cm ⁻³	1.30
<i>D</i> _{exp} (floatation in ClCH ₂ CH ₂ Cl/CHCl ₃), g cm ⁻³	1.29
Radiation	Mo K α (0.71073 Å)
Temperature, °C	23
μ , cm ⁻¹	7.20
Scan mode	ω -2 θ
Data collected	$4^\circ \leq 2\theta \leq 40^\circ$ $\pm h \pm k \pm l$
Unique data measured	5470
Obs. data [$I > 3\sigma(I)$]	3711
Weighting scheme	unit weights
No. of variables	667
<i>R</i> factor	0.060
<i>R</i> _w factor	0.068
<i>F</i> (000)	1184
Residual extreme in final difference map, e Å ⁻³	0.73

h. The solvent was removed by vacuum. The residue was dissolved in CH₂Cl₂ and diethyl ether. The solution was left overnight at room temperature and red crystals were produced. Yield: 0.46 g (43%), 157–162°C. Anal. Calcd for C₆₁H₅₈Co₂N₂O₅P₄: C, 64.22; H, 5.12; N, 2.46. Found: C, 64.18; H, 5.01; N, 2.32%. IR: $\nu(\text{CO})$, 1960s, 1870s, br cm⁻¹. ³¹P NMR (CDCl₃) δ 83.72 ppm. ESR (solid, 77 K): $g_{\text{iso}} = 2.132$ and 2.357. Conductivity (CH₃CN): 62.8 Ω^{-1} cm² mol⁻¹. XPS (Co, 2p_{3/2}): 778.9 (FWHM = 1.71) and 780.0 (FWHM = 1.92) eV.

2.4. X-ray crystallography

A crystal of **2** was mounted on a glass fibre in a random orientation. Preliminary examination and data collection were performed with Mo K α radiation ($\lambda = 0.71073$ Å) on an Enraf-Nonius CAD4 diffractometer equipped with a graphite crystal, incident beam monochromator. The corrections for LP factors and for empirical absorptions were applied to the intensity data. The experimental details of X-ray diffraction study of **2** are listed in Table 1.

The crystal structure of **2** was solved by the direct phase determination method (MULTAN 82). Two independent Co atoms were located on an E-map, and the coordinates of the other non-hydrogen atoms were found by successive difference Fourier syntheses. The hydrogen atoms were not included in the refinements and the calculations of structure factors. In the final refinement by full-matrix least-squares, anisotropic thermal parameters were refined for all non-hydrogen atoms, where all carbon atoms were refined isotropically in order to reduce the number of variables. The final reliability indices are given in Table 1. The fractional coordinates and thermal parameters of non-hydrogen atoms are listed in Table 2.

All calculations were performed on a PDP11/44 computer using SDP-PLUS programs.

3. Results and discussion

Dicobalt complexes of the type Co₂(CO)_{8-2n}(L-L)_n (*n* = 1–3) are normally obtained from reaction of bidentate ligands R₂PXPR₂ (X = CH₂ or NR') with Co₂(CO)₈ [12–15]. The ligand Ph₂PN(ⁱBu)PPh₂ thus reacts with Co₂(CO)₈ (1:1) to give a complex of the type Co₂(CO)₆(L-L), Co₂(CO)₆[μ-Ph₂PN(ⁱBu)PPh₂]₂ (1). However, reaction of Ph₂PN(ⁱBu)PPh₂/Co₂(CO)₈ in a 2:1 mole ratio does not give Co₂[μ-Ph₂PN(ⁱBu)PPh₂]₂(CO)₄, but instead a novel species, [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)][Co(CO)₄] (**2**), has been obtained. The perspective drawing of complex **2** based on X-ray diffraction analysis is shown in Fig. 1. Selected bond distances and angles for **2** are given in Table 3.

The structure of **2** suggests formulation as an ion-pair complex, since phosphines or phosphites are well known to cause disproportionation of Co₂(CO)₈ to form ion-pair complexes, [Co(PR₃)_n(CO)_{5-n}][Co(CO)₄] [16–18]. However, X-ray diffraction, ESR, variable-temperature magnetic susceptibility, conductivity and XPS measurement appear to show formation of **2** as a d⁹-d⁹ cage molecule-pair instead of as an ion-pair complex. The structure determination by X-ray diffraction analysis has shown distinct differences between the structural data of the neutral species [Co(CO)₄] and those reported for anionic [Co(CO)₄]⁻ [19,20]. The average Co-C and C-O distances are 1.81 and 1.09 Å, respectively, in the neutral Co(CO)₄ unit, but 1.77 and 1.14 Å in the Co(CO)₄⁻ anion, which show that the cobalt atom exhibits a weaker d_π-p_π^{*} property in the former than in the latter. The latter shows the ν(CO) stretching vibration at 1870 vs cm⁻¹, and the former has three absorption peaks at 2015 w, 1931 s and 1895 vs cm⁻¹.

TABLE 2. Fractional coordinates and equivalent isotropic thermal parameters of non-hydrogen atoms for **2**

	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{eq} (Å ²)
Co(1)	0.1118(1)	0.26129(9)	0.28761(7)	2.73(3)
P(1)	-0.0251(2)	0.3507(2)	0.2458(2)	3.05(6)
P(2)	0.1601(2)	0.4127(2)	0.2590(2)	3.06(6)
P(3)	0.2694(2)	0.1710(2)	0.2872(2)	3.07(6)
P(4)	0.0931(2)	0.1107(2)	0.2548(2)	3.07(6)
O	0.0267(6)	0.2581(6)	0.4485(4)	5.8(2)
N(1)	0.0289(6)	0.4617(5)	0.2498(4)	3.1(2)
N(2)	0.2149(6)	0.0600(5)	0.2828(4)	3.3(2)
C	0.0605(8)	0.2585(7)	0.3861(6)	4.0(2)
C(5)	-0.0077(8)	0.5639(6)	0.2190(6)	3.9(2)
C(6)	-0.0208(9)	0.6429(7)	0.2772(7)	5.0(3)
C(7)	-0.123(1)	0.631(1)	0.3394(8)	7.2(4)
C(8)	-0.033(1)	0.7494(8)	0.2351(9)	8.1(4)
C(9)	0.2726(9)	-0.0428(7)	0.2666(6)	4.4(3)
C(10)	0.2240(9)	-0.1231(7)	0.3243(6)	4.8(3)
C(11)	0.258(1)	-0.114(1)	0.4011(7)	7.3(4)
C(12)	0.266(1)	-0.2294(8)	0.2914(9)	7.1(4)
C(21)	-0.0514(8)	0.3411(7)	0.1509(6)	3.7(2)
C(22)	0.0121(8)	0.3887(7)	0.0907(6)	4.3(3)
C(23)	-0.004(1)	0.3729(8)	0.0169(6)	5.5(3)
C(24)	-0.082(1)	0.3097(9)	0.0047(7)	6.7(3)
C(25)	-0.146(1)	0.2605(9)	0.0660(7)	5.9(3)
C(26)	-0.1317(8)	0.2761(7)	0.1383(6)	4.9(3)
C(31)	-0.1656(7)	0.3560(7)	0.2965(5)	3.6(2)
C(32)	-0.2475(8)	0.4249(8)	0.2741(7)	4.9(3)
C(33)	-0.3560(9)	0.4236(8)	0.3103(7)	5.7(3)
C(34)	-0.3837(9)	0.3526(8)	0.3663(7)	5.7(3)
C(35)	-0.3021(9)	0.2821(8)	0.3891(7)	5.2(3)
C(36)	-0.1928(7)	0.2819(7)	0.3538(6)	3.9(2)
C(41)	0.2483(7)	0.4535(7)	0.1740(5)	3.3(2)
C(42)	0.2713(8)	0.5571(8)	0.1641(7)	5.0(3)
C(43)	0.327(1)	0.5901(9)	0.0946(8)	6.7(3)
C(44)	0.356(1)	0.524(1)	0.0359(8)	6.9(4)
C(45)	0.3341(9)	0.4216(9)	0.0480(7)	6.0(3)
C(46)	0.2791(8)	0.3885(8)	0.1175(6)	4.0(2)
C(51)	0.2043(8)	0.4764(7)	0.3334(5)	3.7(2)
C(52)	0.3177(8)	0.4898(7)	0.3307(6)	4.5(3)
C(53)	0.3531(9)	0.5288(8)	0.3936(7)	6.1(3)
C(54)	0.215(1)	0.5512(9)	0.4575(7)	7.0(3)
C(55)	0.165(1)	0.5383(9)	0.4589(7)	5.9(3)
C(56)	0.1290(9)	0.4975(7)	0.3982(6)	4.7(3)
C(61)	0.3423(7)	0.1650(7)	0.3668(5)	3.4(2)
C(62)	0.4366(8)	0.0966(7)	0.3692(6)	4.7(3)
C(63)	0.4936(8)	0.0949(8)	0.4293(7)	5.3(3)
C(64)	0.4611(8)	0.1602(9)	0.4868(6)	5.4(3)
C(65)	0.3688(9)	0.2298(9)	0.4843(6)	5.6(3)
C(66)	0.3080(8)	0.2329(8)	0.4237(6)	4.4(2)
C(71)	0.3821(7)	0.1792(6)	0.2102(5)	3.4(2)
C(72)	0.3853(9)	0.1250(8)	0.1458(6)	4.6(3)
C(73)	0.469(1)	0.1381(9)	0.0839(7)	6.2(3)
C(74)	0.548(1)	0.208(1)	0.0880(8)	7.0(4)
C(75)	0.5455(9)	0.2617(9)	0.1516(7)	5.8(3)
C(76)	0.4618(8)	0.2474(8)	0.2146(7)	5.0(3)
C(81)	-0.0194(8)	0.0467(6)	0.3086(6)	3.7(2)
C(82)	-0.0131(9)	0.0155(7)	0.3829(6)	4.3(3)
C(83)	-0.103(1)	-0.0281(9)	0.4265(7)	6.2(3)
C(84)	-0.195(1)	-0.0414(9)	0.3954(8)	6.7(3)
C(85)	-0.204(1)	-0.0078(9)	0.3223(8)	6.7(3)
C(86)	-0.1134(8)	0.0357(8)	0.2756(7)	5.1(3)



TABLE 2 (continued)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{eq} Å ²
C(91)	0.0997(8)	0.0719(7)	0.1614(6)	3.9(2)
C(92)	0.1332(8)	0.1399(7)	0.1001(6)	4.1(2)
C(93)	0.151(1)	0.1090(9)	0.0281(7)	6.4(3)
C(94)	0.138(2)	0.009(1)	0.0149(8)	11.0(5)
C(95)	0.101(2)	-0.059(1)	0.0744(9)	11.5(6)
C(96)	0.085(1)	-0.0284(8)	0.1489(7)	6.8(3)
Co(2)	0.6132(2)	-0.1963(2)	0.1799(1)	9.65(7)
O(1)	0.4217(8)	-0.1635(8)	0.1065(6)	9.4(3)
O(2)	0.564(1)	-0.242(1)	0.3365(7)	17.6(6)
O(3)	0.717(1)	-0.007(1)	0.1561(8)	15.9(5)
O(4)	0.756(1)	-0.364(1)	0.1224(8)	14.6(4)
C(1)	0.496(1)	-0.178(1)	0.1355(8)	7.8(4)
C(2)	0.583(1)	-0.223(2)	0.2729(7)	13.4(7)
C(3)	0.679(2)	-0.081(1)	0.165(1)	15.8(6)
C(4)	0.699(2)	-0.296(1)	0.1449(9)	12.4(5)

$$B_{eq} = (4/3) \cdot [a^2 \cdot B_{1,1} + b^2 \cdot B_{2,2} + c^2 \cdot B_{3,3} + ab(\cos \gamma) \cdot B_{1,2} + ac(\cos \beta) \cdot B_{1,3} + bc(\cos \alpha) \cdot B_{2,3}]$$

In contrast with the reported trigonal bipyramidal configurations of [Co(dppm)₂(CO)]⁺ [21] and [Co(dppe)₂(CO)]⁺ [22,23] the species [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)] exhibits a square pyramidal geometry and its structure is similar to a five-coordinate cobalt(0) complex with stronger π -acid diphosphine ligand, namely [2,3-bis(diphenylphosphino) maleic anhydride tricarbonyl cobalt(0)] [24]. Four P atoms lie in the same plane and the Co atom is located over the plane about 0.4 Å towards the carbonyl in this species. The average Co–P distance (2.17 Å) is shorter than

those in [Co(dppm)₂(CO)]⁺ (2.217 Å) and [Co(dppe)₂(CO)]⁺ (2.231 Å). Orpen [25] has recently described that the σ^* orbitals of P–R bonds have been recognized as playing the role of acceptor in PR₃. The order of increasing π -acid character is:



Therefore, the ligand Ph₂PN(ⁱBu)PPh₂ is expected to be a stronger π -acid than dppm and dppe, and a stronger back-donating d_{π} – σ^* bond could be formed with the Co atom, which may be an important reason for the formation of the neutral species 2.

The ESR spectrum of the powdered sample of the species 2 at 77 K shows two broad signals centred at $g_{\text{iso}} = 2.132$ and 2.357. The former is close to $g_{\perp} = 2.128$ of species Co(CO)₄ in the matrix ESR spectrum at 6 K [26]. The magnetic moment μ_{eff} is calculated as 3.75 BM based on the determination of variable-temperature magnetic susceptibility (1.4–295 K) (Fig. 2), which suggests that there are two unpaired electrons in 2. The conductivity of 2 in CH₃CN is 62.8 Ω⁻¹ cm² mol⁻¹ at room temperature, which falls within the range for a non-electrolyte [27]. In contrast [Co(dppm)₂(CO)]⁺ complexes are diamagnetic solids which give conducting solutions [28]. The photoelectron spectrum of 2 shows that the Co 2p_{3/2} lines locate at 778.9 eV (FWHM = 1.71 eV) and 780.0 eV (FWHM = 1.92 eV). The latter is close to that (780.2 eV) of Co₂(CO)₈ determined at the same conditions. The




Fig. 1. A perspective view of 2 showing the atomic labelling scheme.

TABLE 3. Selected bond distances (Å) and angles (°) for 2

Co(1)–P(1)	2.179(2)	Co(1)–P(4)	2.173(2)
Co(1)–P(2)	2.174(2)	Co(1)–C	1.796(9)
Co(1)–P(3)	2.180(2)	O–C	1.140(7)
P(1)–Co(1)–P(2)	71.94(6)	P(2)–Co(1)–P(4)	150.29(7)
P(1)–Co(1)–P(3)	159.74(7)	P(2)–Co(1)–C	105.9(2)
P(1)–Co(1)–P(4)	102.24(6)	P(3)–Co(1)–P(4)	71.94(6)
P(1)–Co(1)–C	100.0(2)	P(3)–Co(1)–C	100.3(2)
P(2)–Co(1)–P(3)	103.17(6)	P(4)–Co(1)–C	103.8(2)
Co(1)–C–O	178.8(6)		
Co(2)–C(1)	1.75(2)	Co(2)–C(3)	1.80(1)
Co(2)–C(2)	1.70(1)	Co(2)–C(4)	1.70(1)
O(1)–C(1)	1.128(9)	O(3)–C(3)	1.13(1)
O(2)–C(2)	1.16(1)	O(4)–C(4)	1.14(1)
C(1)–Co(2)–C(2)	113.3(5)	C(2)–Co(2)–C(3)	108.7(7)
C(1)–Co(2)–C(3)	106.9(6)	C(2)–Co(2)–C(4)	105.6(6)
C(1)–Co(2)–C(4)	110.7(5)	C(3)–Co(2)–C(4)	111.7(8)
Co(2)–C(1)–O(1)	177.9(9)	Co(2)–C(3)–O(3)	177(2)
Co(2)–C(2)–O(2)	179(1)	Co(2)–C(4)–O(4)	180(1)
P(1)–N(1)	1.694(4)	P(3)–N(2)	1.698(4)
P(1)–C(21)	1.814(7)	P(3)–C(61)	1.812(7)
P(1)–C(31)	1.823(6)	P(3)–C(71)	1.811(7)
P(2)–N(1)	1.712(5)	P(4)–N(2)	1.724(5)
P(2)–C(41)	1.826(6)	P(4)–C(81)	1.819(7)
P(2)–C(51)	1.816(7)	P(4)–C(91)	1.785(6)
N(1)–C(5)	1.498(7)	N(2)–C(9)	1.504(7)
Co(1)–P(1)–N(1)	93.9(2)	Co(1)–P(3)–N(2)	93.9(2)
Co(1)–P(1)–C(21)	121.6(2)	Co(1)–P(3)–C(61)	121.3(2)
Co(1)–P(1)–C(31)	122.2(2)	Co(1)–P(3)–C(71)	122.0(2)
N(1)–P(1)–C(21)	107.6(3)	N(2)–P(3)–C(61)	109.4(2)
N(1)–P(1)–C(31)	110.4(2)	N(2)–P(3)–C(71)	107.9(3)
C(21)–P(1)–C(31)	100.3(3)	C(61)–P(3)–C(71)	101.2(3)
Co(1)–P(2)–N(1)	93.7(2)	Co(1)–P(4)–N(2)	93.4(2)
Co(1)–P(2)–C(41)	126.5(2)	Co(1)–P(4)–C(31)	116.5(2)
Co(1)–P(2)–C(51)	115.8(2)	Co(1)–P(4)–C(91)	126.4(2)
N(1)–P(2)–C(41)	105.9(2)	N(2)–P(4)–C(31)	108.4(3)
N(1)–P(2)–C(51)	108.3(3)	N(2)–P(4)–C(91)	105.5(3)
C(41)–P(2)–C(51)	104.5(3)	C(81)–P(4)–C(91)	104.5(3)
P(1)–N(1)–P(2)	97.3(2)	P(3)–N(2)–P(4)	96.8(2)
P(1)–N(1)–C(5)	128.9(4)	P(3)–N(2)–C(9)	129.3(4)
P(2)–N(1)–C(5)	128.2(4)	P(4)–N(2)–C(9)	126.7(4)

Numbers in parentheses are estimated standard deviations in the least significant digits.

crystal structural analysis indicated that the species Co(CO)₄ is enclosed in the cage consisted of species [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)] (Fig. 3) and the Co(1)–Co(2) distance is 8.320 Å.

The mechanism for the formation of 2 has initially been investigated by ³¹P NMR. The ³¹P NMR species shows that an intermediate is present during the reaction, in which there might be three different environmental phosphorus atoms and the coupling interaction is complicated. It has been established [18] that Co₂(CO)₈ reacts with dppm rapidly (a few minutes) to give a salt [Co(dppm)(CO)₃][Co(CO)₄], which loses CO to form Co₂(μ-dppm)(μ-CO)₂(CO)₄. Following the




Fig. 2. Plot of molar susceptibility, χ_M , and effective moment, μ_{eff} for 2.

above considerations and in view of the ³¹P NMR spectrum we consider the intermediate probably has the following structure:





Fig. 3. A view of the packing arrangement in a cell of crystal 2.





Fig. 4. $^{13}P\{^1H\}$ NMR monitoring of the reaction between ligand $Ph_2PN(^iBu)PPh_2$ and $Co_2(CO)_8$ (solvent: $CDCl_3$, room temperature). (a) $t = 5$ – 35 min; (b) $t = 60$ min; (c) peak height of chemical shift as a function of reaction time. +, ligand $Ph_2PN(^iBu)PPh_2$, δ : 62.73 ppm. *, species 2, δ : 83.52 ppm. x, species 3, δ : 63.81 ppm.

We have isolated a minor orange species (**3**) by fractional crystallization, which gives three CO absorption bands at 2000s, 1962s and 1888s, cm^{-1} , and this seems to support our hypothesis. The relation of the amount of phosphazane, intermediate **3** and product **2**, *vs.* the reaction time detected by ³¹P NMR is given in Fig. 4.

Here the ³¹P signal of **1** ($\delta = 79.13$ ppm) is not observed. It is possible that **1** reacts rapidly with Ph₂PN(ⁱBu)PPh₂ to form the stable intermediate **3**, and the latter changes slowly to **2** by homolytic Co–Co bond cleavage and heterolytic P–Co bond cleavage. After reacting for 35 min the peak height for the ligand, Ph₂PN(ⁱBu)PPh₂, disappeared (Fig. 4(a)), however the peak height for species **2** continued to increase (Fig. 4(b) and (c)) which shows that the intermediate **3** continues to transfer to product **2**. Therefore, the total process of the reaction of Co₂(CO)₈ with Ph₂PN(ⁱBu)PPh₂ (1:2) is probably as follows:



2 probably has different structures in the solid state and in solution, and a strong interaction between [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)] and [Co(CO)₄] might be present when in solution [29].

Acknowledgment

We are grateful to the National Science Foundation and Elemento-Organic Chemistry Laboratory, Nankai University, for financial support.

References

- 1 J. Puddephatt, *Chem. Soc. Rev.*, 12 (1983) 99.
- 2 P. Chaudret, B. Delavaux and R. Poilblanc, *Coord. Chem. Rev.*, 86 (1988) 191.
- 3 Z.Z. Zhang, X.K. Wang and Z. Xi, *Huaxue Tongbao*, 8 (1988) 1.
- 4 T.R. Johnson and J.F. Nixon, *J. Chem. Soc. A*, (1969) 2518.
- 5 R.B. King and T.W. Lee, *Inorg. Chem.*, 21 (1982) 319.
- 6 G. de Leeuw, T.S. Field, R.T. Haines, B. McCulloch, E. Meitjies, C. Moberg, G.M. Olivier, P. Ramdial, C.N. Sampson, B. Sigwarth, N.D. Steen and K.G. Moodley, *J. Organomet. Chem.*, 275 (1984) 99.
- 7 A. Blagg, S.W. Carr, G.R. Cooper, I.D. Dobson, J.B. Gill, D.C. Goodall, B.L. Shaw, N. Taylor and T. Boddington, *J. Chem. Soc., Dalton Trans.*, (1985) 1213.
- 8 J. Ellermann, G. Szucsanyi and E. Wilhelm, *Chem. Ber.* 118 (1985) 1588.
- 9 A. Tarassoli, H.J. Chen, M.L. Thompson, V.S. Allured, R.C. Haltiwanger and A.D. Norman, *Inorg. Chem.*, 25 (1986) 4052.
- 10 I. Uson, J. Forries, R. Navarro and I.J. Cebollada, *J. Organomet. Chem.*, 304 (1986) 381.
- 11 M.S. Balakrishna, T.K. Prakasha, S.S. Krishnamurthy, U. Siriwardane and N.S. Hosmane, *J. Organomet. Chem.*, 390 (1990) 203.
- 12 C. Moreno, Ma.J. Macazaga and S. Delgado, *Inorg. Chim. Acta*, 182 (1991) 55.
- 13 G. de Leeuw, J.S. Field and R.J. Haines, *J. Organomet. Chem.*, 359 (1989) 245.
- 14 C.M. Brown, J.E. Finholt, R.B. King and J.W. Bibber, *Inorg. Chem.*, 21 (1982) 2139.
- 15 J. Ellermann, N. Geheeb, G. Zoubek and G. Thiele, *Z. Naturforsch.*, 32b (1977) 1271.
- 16 M.S. Arah, A. Maisonat, S. Attali and R. Polilblanc, *J. Organomet. Chem.*, 67 (1974) 109 and references cited therein.
- 17 M. Absi-Halabi, J.D. Atwood, N.P. Forbus and T.L. Brown, *J. Am. Chem. Soc.*, 102 (1980) 6245.
- 18 E.C. Linic and R.B. Hanson, *Inorg. Chem.*, 25 (1986) 812.
- 19 N.C. Baenziger, R.M. Flynn and N.L. Holy, *Acta Crystallogr. B*, 35 (1979) 741.
- 20 F. Calderazzo, G. Fachinetti and F. Marchetti, *J. Chem. Soc., Chem. Commun.*, (1981) 181.
- 21 D.J. Elliot, D.G. Holah, A.N. Hughes, V.R. Magnuson, I.M. Moser, R.J. Puddephatt and W. Xu, *Organometallics*, 10 (1991) 3933.
- 22 Z.Z. Zhang, X.K. Wang, Z. Xi, R.J. Wang and H.G. Wang, *J. Organomet. Chem.*, 352 (1988) C22.
- 23 X.K. Yao, H.G. Wang, Z.Z. Zhang, X.K. Wang and Z. Xi, *Acta Chim. Sinica*, 48 (1990) 770.
- 24 D. Fenske, *Chem. Ber.*, 112 (1979) 363.
- 25 A.G. Orpen and N.G. Connelly, *J. Chem. Soc., Chem. Commun.*, (1985) 1310.
- 26 L.A. Hanlan, P.E. Kundig, B.R. McGarvey and G.A. Ozin, *J. Am. Chem. Soc.*, 97 (1975) 7054.
- 27 W. Geary, *Coord. Chem. Rev.*, 7 (1971) 81.
- 28 C. Carriedo, P. Gomer-Sal, P. Royo, S. Martinier-Carrera and S. Garcia-Blanco, *J. Organomet. Chem.*, 301 (1986) 79.
- 29 G. Fachinetti, G. Fochi and T. Funaioli, *J. Organomet. Chem.*, 301 (1986) 91.