

Reductive Elimination of a Carbon-Carbon Bond from Bis(trialkylphosphine)-3,3-dimethylplatinacyclobutanes Produces Bis(trialkylphosphine)platinum(0) and 1,1-Dimethylcyclopropane¹


Robert DiCosimo² and George M. Whitesides*

Contribution from the Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139. Received October 26, 1981


Abstract: The thermal decompositions of several platinacyclobutanes—bis(triethylphosphine)- (1), bis(triisopropylphosphine)- (2), and bis(tricyclohexylphosphine)-3,3-dimethylplatinacyclobutane (3)—have been examined in cyclohexane solution. Decomposition of 1 proceeds heterogeneously: it produces platinum metal, neopentane, 1,1-dimethylcyclopropane, and ethylene and seems to be autocatalytic. Decompositions conducted using reaction mixtures containing mercury(0) proceed homogeneously and produce predominately 1,1-dimethylcyclopropane; any platinum metal produced is apparently amalgamated by the Hg(0) and rendered catalytically inactive. Thermal decompositions of 2 and 3 proceed homogeneously, even in the absence of Hg(0), and yield 1,1-dimethylcyclopropane, bis(trialkylphosphine)platinum(0), and a small quantity of neopentane. The rates of decomposition are decreased by addition of trialkylphosphine, and a study of this dependence of rate on concentration of added phosphine for 2 indicates that one phosphine dissociates to produce a three-coordinate platinum(II) intermediate prior to reductive elimination of 1,1-dimethylcyclopropane. The Arrhenius activation parameters for decomposition of 2 and 3 are respectively $E_a = 46 \pm 3 \text{ kcal mol}^{-1}$, $\log A = 23 \pm 2$ and $E_a = 42 \pm 2 \text{ kcal mol}^{-1}$, $\log A = 21 \pm 1$. Substitution of deuterium for hydrogen in the triisopropylphosphine groups of 2 produces no change in the rate of reaction. We cannot distinguish between mechanisms in which reductive elimination of 1,1-dimethylcyclopropane occurs directly from the three-coordinate intermediate and those in which oxidative addition of a C-H bond to platinum by cyclometalation of the phosphine produces a Pt(IV) intermediate in a step preceding rate-limiting reductive elimination of 1,1-dimethylcyclopropane.

Heterogeneous reactions of hydrocarbons with transition metals (cracking, reforming, hydrogenation) involve cleaving and reforming C-H and C-C bonds, and are believed to proceed through metallacyclic and surface-alkyl intermediates.³ Although studies of the mechanisms of these heterogeneous reactions have been difficult, the studies of related reactions in homogeneous transition-metal systems have provided examples of C-H and C-C bond cleavage and formation by metathesis,⁴ β -hydride elimination and insertion of olefins into metal-hydrogen bonds,^{5,6} oxidative addition and reductive elimination of C-H^{7,8} and C-C bonds,⁹⁻¹³ and cyclometalation.¹⁴⁻¹⁶ This paper is concerned with the

Scheme I. Reaction of Triisopropylphosphine with (1,5-Cyclooctadiene)dineopentylplatinum(II) (L = *i*-Pr₃P; Yields Based on (1,5-Cyclooctadiene)dineopentylplatinum(II))



Scheme II. Reaction of Tricyclohexylphosphine with (1,5-Cyclooctadiene)dineopentylplatinum(II) (L = *n*-C₆H₁₃P; Yields Based on (1,5-Cyclooctadiene)dineopentylplatinum(II))



mechanism of carbon-carbon bond formation by reductive elimination from platinum. A previous study of the thermal decom-

- (1) Supported by the National Science Foundation, Grant CHE 8012722.
- (2) Chevron Fellow, 1980-1982.
- (3) Clarke, J. K. A.; Rooney, J. J. *Adv. Catal.* **1976**, *25*, 125-183. Sinfelt, J. H. *Science (Washington, D.C.)* **1977**, *195*, 641-646.
- (4) Grubbs, R. H.; Hoppin, C. R. *J. Am. Chem. Soc.* **1979**, *101*, 1499-1508. Leconte, M.; Basset, J. M. *Ann. N.Y. Acad. Sci.* **1980**, *333*, 165-187. Rappe, A. K.; Goddard, W. A.; III *J. Am. Chem. Soc.* **1980**, *102*, 5114-5115.
- (5) Ozawa, F.; Ito, T.; Yamamoto, A. *J. Am. Chem. Soc.* **1980**, *102*, 6457-6463. Komiya, S.; Yamamoto, A.; Yamamoto, T. *Chem. Lett.* **1978**, 1273-1276.
- (6) McCarthy, T. J.; Nuzzo, R. G.; Whitesides, G. M. *J. Am. Chem. Soc.* **1981**, *103*, 3396-3403. Whitesides, G. M.; Gaasch, J. F.; Stedronsky, E. R. *Ibid.* **1972**, *94*, 5258-5270.
- (7) Abis, L.; Sen, A.; Halpern, J. *J. Am. Chem. Soc.* **1978**, *100*, 2915-2916.
- (8) Crocker, C.; Errington, R. J.; Markham, R.; Molton, C. J.; Odell, K. J.; Shaw, B. L. *J. Am. Chem. Soc.* **1980**, *102*, 4373-4379. Simpson, S. J.; Turner, H. W.; Andersen, R. A. *Ibid.* **1979**, *101*, 7728-7729.
- (9) Ruddick, J. D.; Shaw, B. L. *J. Chem. Soc. A* **1969**, 2969-2970. Clark, H. C.; Manzer, L. E. *Inorg. Chem.* **1973**, *12*, 362-368. Brown, M. P.; Puddenphatt, R. J.; Upton, C. E. E. *J. Chem. Soc., Dalton Trans.* **1974**, 2457-2465. Al-Essa, R. J.; Puddephatt, R. J.; Thompson, P. J.; Tipper, C. F. H. *J. Am. Chem. Soc.* **1980**, *102*, 7546-7553.
- (10) Tamaki, A.; Magennis, S. A.; Kochi, J. K. *J. Am. Chem. Soc.* **1974**, *96*, 6140-6148. Komiya, A.; Albright, T. A.; Hoffmann, R.; Kochi, J. K. *Ibid.* **1976**, *98*, 7255-7265.
- (11) Gillie, A.; Stille, J. K. *J. Am. Chem. Soc.* **1980**, *102*, 4933-4941. Moravskiy, A.; Stille, J. K. *Ibid.* **1981**, *103*, 4182-4186.
- (12) Ito, T.; Tsuchiya, H.; Yamamoto, A. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 1319-1327.
- (13) Smith, G.; Kochi, J. K. *J. Organomet. Chem.* **1980**, *198*, 199-214.
- (14) Foley, P.; DiCosimo, R.; Whitesides, G. M. *J. Am. Chem. Soc.* **1980**, *102*, 6713-6725. Moore, S. S.; DiCosimo, R.; Sowinski, A. F.; Whitesides, G. M. *Ibid.*, **1981**, *103*, 948-949.

- (15) Andersen, R. A.; Jones, R. A.; Wilkinson, G. *J. Chem. Soc., Dalton Trans.* **1978**, 466-453. Andersen, R. A.; Jones, R. A.; Wilkinson, G.; Hursthouse, M. B.; Malik, K. M. A. *J. Chem. Soc., Chem. Commun.* **1977**, 283-285.

position of platinum(II) metallacycles¹⁷ indicated that reductive elimination of a C–C bond took place only after the oxidative addition of a halogenated solvent molecule (methylene chloride) to platinum(II) had occurred and generated a platinum(IV) intermediate. In nonhalogenated solvents, only β -hydride elimination of alkene was observed. Here we describe a study of the mechanism of thermal decomposition of bis(trialkylphosphine)-3,3-dimethylplatinacyclobutanes to bis(trialkylphosphine)platinum(0) and 1,1-dimethylcyclopropane in hydrocarbon solvents. The reaction proceeds by an initial dissociation of one of the trialkylphosphine ligands and generation of a three-coordinate platinum(II) intermediate. The large Arrhenius preexponential frequency factors that are observed suggest that the reductive elimination of 1,1-dimethylcyclopropane is rate limiting. The possibility that oxidative addition of a C–H bond of the coordinated phosphine to Pt(II) to form a Pt(IV) intermediate precedes reductive elimination of dimethylcyclopropane can neither be proved nor disproved.

Results

Preparation of Bis(triisopropylphosphine)-3,3-dimethylplatinacyclobutane (2) and Bis(tricyclohexylphosphine)-3,3-dimethylplatinacyclobutane (3). Platinacyclobutanes **2** and **3** were isolated in 19% and 17% yields, respectively, as air-stable, white crystalline solids from the reaction of 2 equiv of the corresponding trialkylphosphine with dineopentyl(1,5-cyclooctadiene)platinum(II) in cyclohexane at 50 °C and were characterized by a combination of ¹H and ³¹P{¹H} NMR spectroscopy, elemental analysis, and chemical reactions. Addition of DCl to a solution of **2** in cyclohexane generated 1 equiv of neopentane which was determined by GC/MS analysis to be $\geq 95\%$ (CH₃)₂C(CH₂D)₂; repeating the procedure with **3** produced 1 equiv of neopentane which was $\geq 94\%$ (CH₃)₂C(CH₂D)₂.

Unlike the reaction of 2 equiv of triethylphosphine with (1,5-cyclooctadiene)dineopentylplatinum(II), which cleanly produces dineopentylbis(triethylphosphine)platinum(II),¹⁴ the corresponding reaction of (1,5-COD)Pt[CH₂C(CH₃)₂]₂ with triisopropylphosphine (Scheme I) or tricyclohexylphosphine (Scheme II) gave a complex reaction mixture containing *no* dineopentylbis(trialkylphosphine)platinum(II) compounds (as determined by ³¹P{¹H} NMR spectroscopy). Instead, a bis(trialkylphosphine)-3,3-dimethylplatinacyclobutane and bis(trialkylphosphine)platinum(0) complex were produced as the major platinum-containing compounds, along with neopentane and 2,2,5,5-tetramethylhexane. Oxidative addition of an allylic C–H bond of 1,5-cyclooctadiene to platinum(II) is undoubtedly the source of some of the hydrogen atoms which participate in the subsequent reductive elimination of a C–H bond to form neopentane; a 15–20% yield of 1,3,5-cyclooctatriene was obtained from the reaction of triisopropylphosphine with (1,5-cyclooctadiene)dineopentylplatinum(II).

The marked difference in the products of the reaction of (1,5-cyclooctadiene)dineopentylplatinum(II) and triisopropylphosphine, tricyclohexylphosphine, and triethylphosphine may be due to the larger steric bulk of the first two of these phosphines.¹⁸ A square-planar platinum(II) species composed of two neopentyl groups and two sterically demanding trialkylphosphine ligands is probably very unstable. The displacement of 1,5-cyclooctadiene by a bulky phosphine could produce a highly reactive, coordinatively unsaturated 14-electron platinum(II) species which could oxidatively add a C–H bond of a neopentyl group, cyclooctadiene, or phosphine alkyl group to yield the observed products.

Thermal Decomposition of Bis(trialkylphosphine)-3,3-dimethylplatinacyclobutanes. Products. The thermal decomposition of a cyclohexane solution of bis(triethylphosphine)-3,3-dimethylplatinacyclobutane (**1**) in a sealed tube at 157 °C produces





Figure 1. First-order plots of the thermal decompositions of **2** (■, □), **2**-(L^D)₂ (Δ), and **3** (●, ○) in cyclohexane at 116 °C. Solid symbols represent data obtained by ³¹P{¹H} NMR spectroscopy; open symbols are based on GLC data.

a heterogeneous reaction mixture containing neopentane, 1,1-dimethylcyclopropane, ethylene, and platinum metal.¹⁴ The formation of a platinum mirror and colloidal platinum was probably due to the disproportionation of bis(triethylphosphine)platinum(0), formed by the reductive elimination of 1,1-dimethylcyclopropane from **1**, to produce tris(triethylphosphine)platinum(0) and platinum metal. In fact, when the thermal decomposition of **1** was repeated at 178 °C in the presence of Hg(0), the reaction mixture remained homogeneous. 1,1-Dimethylcyclopropane and neopentane were produced in a ratio of 94:6 throughout the thermolysis. No other hydrocarbon products were observed by GLC when the reaction was repeated in isooctane or 2,2,4,4-tetramethylpentane. ³¹P{¹H} NMR spectroscopy revealed only the production of free triethylphosphine as **1** was consumed, and no (Et₃P)₃Pt⁰ was observed, indicating that platinum metal was amalgamated with mercury and rendered inert. The small percentage of neopentane produced in the presence of Hg(0) indicates that its formation in larger quantities in the absence of Hg(0) may be due to reactions catalyzed by the platinum metal which is formed.

Heating a sealed tube containing a 0.08 M solution of **2** or **3** in cyclohexane resulted in a homogeneous reaction which produced either bis(triisopropylphosphine)platinum(0) (**4**) or bis(tricyclohexylphosphine)platinum(0) (**5**), respectively, 1,1-dimethylcyclopropane, and a small quantity of neopentane (eq 1).¹⁹ The



homogeneous nature of the reaction and the high yields of 1,1-dimethylcyclopropane can be attributed to the stability of the bis(trialkylphosphine)platinum(0) complexes which are formed.²⁰ The source of the two hydrogen atoms which are required in the production of neopentane have not been positively identified, but they did not originate from the solvent, since the thermal decomposition of **2** in C₆D₁₂ produced only neopentane-d₀. When a cyclohexane solution of **2**-(L^D)₂ (L^D = [(CD₂)₂CD]₃P) was heated for 0.5 h at 178 °C, the neopentane produced was 80% d₀, 15% d₁, and 4% d₂, indicating some transfer of hydrogen atoms originally present in the phosphine ligand to the platinacyclobutane ring to form neopentane.

(19) The corresponding nickel analogue bis(triphenylphosphine)-3,3-dimethylnickelacyclobutane, is believed to be in equilibrium with an olefin-coordinated nickel-carbene complex and decomposes in toluene upon heating at 49 °C to give ethylene (15%), 2-methylpropene (26%), 1,1-dimethylcyclopropane (47%), 3-methyl-1-butene (6%), and 2,2-dimethylpropane (6%).¹⁶

(20) The preparation and reactivities of a variety of bis(tertiary phosphine)platinum(0) complexes have been reported: Otsuka, S.; Yoshida, T.; Matsumoto, M.; Nakatsu, K. *J. Am. Chem. Soc.* **1976**, *98*, 5850–5858. Yoshida, T.; Otsuka, S. *Ibid.* **1977**, *99*, 2134–2140.

(16) Miyashita, A.; Grubbs, R. H. *Tetrahedron Lett.* **1981**, *22*, 1255–1256.

(17) McDermott, J. X.; White, J. F.; Whitesides, G. M. *J. Am. Chem. Soc.* **1976**, *98*, 6521–6528. Young, G. B.; Whitesides, G. M. *Ibid.* **1978**, *100*, 5808–5815.

(18) The phosphine cone angles for triethylphosphine, triisopropylphosphine, and tricyclohexylphosphine are 132, 160, and 170°, respectively; Tolman, C. A. *Chem. Rev.* **1977**, *77*, 313–348.




Figure 2. Dependence of the rate of thermal decomposition of **2** on the concentration of added triisopropylphosphine. Rate constants were measured at 126.5 °C in cyclohexane solution by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy.


Kinetics. The thermal decomposition of **1** in cyclohexane (in the presence of $\text{Hg}(0)$) was followed by monitoring the appearance of 1,1-dimethylcyclopropane by GLC. This reaction followed first-order kinetics ($k^{\text{obsd}} = 1.2 \times 10^{-5} \text{ s}^{-1}$, 178 °C) through the first half-life of the reaction. Free triethylphosphine was also generated as the reaction progressed due to amalgamation of $\text{Pt}(0)$ by $\text{Hg}(0)$. Because added trialkylphosphine was found to inhibit the rate of decomposition of **2** or **3** (vide infra) and because the effect of $\text{Hg}(0)$ on the mechanism of thermal decomposition of **1** is unknown, further investigation of this reaction was deferred in favor of an examination of the simpler thermolysis of **2** and **3**, which proceeded homogeneously at much lower temperatures than those necessary for the decomposition of **1**.

Rates of decomposition of **2** and **3** were determined by following the disappearance of platinacyclobutane by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy and by monitoring the appearance of 1,1-dimethylcyclopropane by GLC (Figure 1). Rates determined by using these techniques were indistinguishable. Reaction rates were independent of the initial concentration of platinacyclobutane (from $2.4 \times 10^{-2} \text{ M}$ to $8.0 \times 10^{-2} \text{ M}$), and the disappearance of **2** or **3** and appearance of 1,1-dimethylcyclopropane were both first order in platinacyclobutane to greater than 90% decomposition.

Addition of triisopropylphosphine or tricyclohexylphosphine to cyclohexane solutions of the corresponding platinacyclobutane inhibited the reaction, but the yields of 1,1-dimethylcyclopropane and neopentane were unchanged. An examination of the dependence of the rates of decomposition of **2** (0.08 M) in the presence of added triisopropylphosphine (0.005–0.040 M) yielded rate constants which obey the empirical rate eq 2 (Figure 2).

$$-d[\mathbf{2}]/dt = k^{\text{obsd}}[\mathbf{2}][((\text{CH}_3)_2\text{CH})_3\text{P}]^{-0.97 \pm 0.12} \quad (2)$$

The inhibition in the decomposition produced by added trialkylphosphine is not due to an associative process (eq 3) since



the $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of the reaction mixtures containing various concentrations of added phosphine showed no compounds other than platinacyclobutane and free phosphine.

Phosphine dissociation was demonstrated to occur in a pre-equilibrium rather than rate-determining step by heating a cyclohexane solution of **2** (0.07 M) and $\mathbf{2}\text{-}(\text{L}^{\text{D}})_2$ (0.07 M, $\text{L}^{\text{D}} = [(\text{CD}_3)_2\text{CD}]_3\text{P}$) at 50 °C and following the rate of exchange of phosphines by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy. Complete equilibration of deuterated and nondeuterated phosphines among **2**, $\mathbf{2}\text{-L}^{\text{H}}\text{L}^{\text{D}}$, and $\mathbf{2}\text{-}(\text{L}^{\text{D}})_2$ was observed after 8 h, with no detectable decomposition of **2** to bis(triisopropylphosphine)platinum(0) (Figure 3).

Analysis of the temperature dependence of thermolyses of **2** and **3** over the range 90–120 °C yielded Arrhenius activation parameters of $E_a = 46 \pm 3 \text{ kcal mol}^{-1}$, $\log A = 23 \pm 2$ and E_a




Figure 3. $^{31}\text{P}\{^1\text{H}\}$ NMR spectra (101.3 MHz) of a cyclohexane solution of $\mathbf{2}\text{-}(\text{L}^{\text{H}})_2$ (0.07 M) and $\mathbf{2}\text{-}(\text{L}^{\text{D}})_2$ (0.07 M), A, and after heating at 50 °C for 8 h to produce an equilibrium mixture of $\mathbf{2}\text{-}(\text{L}^{\text{H}})_2$, $\mathbf{2}\text{-}(\text{L}^{\text{D}})_2$, and $\mathbf{2}\text{-L}^{\text{H}}\text{L}^{\text{D}}$, B. No decomposition to form 1,1-dimethylcyclopropane and L_2Pt^0 was observed. (^{195}Pt satellites are not shown; the underlined phosphine ($\text{L}^{\text{H(D)}}$) corresponds to the peak(s) indicated.)





Figure 4. Arrhenius plots for the thermal decompositions of **2** (●) and **3** (■) determined by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy.

$= 42 \pm 2 \text{ kcal mol}^{-1}$, $\log A = 21 \pm 1$, respectively (Figure 4). These activation parameters are similar in magnitude to those reported for the thermal decomposition of bis(2,2-dimethylakyl)bis(triethylphosphine)platinum(II) complexes, which produce bis(triethylphosphine)platinacycloalkanes and hydrocarbon.¹⁴ The rate-limiting step for the cyclometalation reactions is believed to be reductive elimination of alkane from a $\text{Pt}(\text{IV})$ intermediate, based on the large Arrhenius preexponential frequency factor and the observation of a kinetic deuterium isotope effect ($k_{\text{H}}/k_{\text{D}} \approx 3$) when deuterium is substituted for hydrogen in the alkyl groups. To examine the possibility that the thermal decomposition of **2** might proceed in a similar manner via a rate-limiting oxidative addition or reductive elimination of a triisopropylphosphine C-H bond, the rate of decomposition of **2** and $\mathbf{2}\text{-}(\text{L}^{\text{D}})_2$ at 116 °C were compared (Figure 1); a $k_{\text{H}}/k_{\text{D}} = 0.98 \pm 0.04$ was obtained, indicating the absence of a rate-limiting oxidative addition or reductive elimination of a C-H bond of the coordinated phosphine.


Discussion

The thermal decomposition of bis(trialkylphosphine)-3,3-dimethylplatinacyclobutane to bis(trialkylphosphine)platinum(0) and 1,1-dimethylcyclopropane can be explained by one of two possible mechanisms. Either mechanism is supported by the observation that dissociation of trialkylphosphine and formation of a coordinatively unsaturated 14-electron $\text{Pt}(\text{II})$ species is required for the reaction to occur. The large Arrhenius preexponential factor ($\log A \approx 21\text{--}23$) is also consistent with a mechanism

Scheme III. Proposed Mechanism for Thermal Decomposition of Bis(trialkylphosphine)-3,3-dimethylplatinacyclobutane in Which a C-C Bond Is Reductively Eliminated from a Pt(II) Intermediate



Scheme IV. Proposed Mechanism for Thermal Decomposition of Bis(trialkylphosphine)-3,3-dimethylplatinacyclobutane in Which a C-C Bond Is Reductively Eliminated from a Pt(IV) Intermediate



in which three particles (phosphine, 1,1-dimethylcyclopropane, and the remaining Pt moiety) are generated in the transition state from one particle in the ground state.

One mechanism of reaction would involve dissociation of trialkylphosphine, rate-limiting reductive elimination of a C-C bond to produce 1,1-dimethylcyclopropane and LPt^0 , and recombination of LPt^0 with dissociated phosphine (Scheme III). Dissociation of a phosphine ligand prior to reductive elimination of a C-C bond from a three-coordinate intermediate has been reported for thermal decompositions of $\text{LAu}(\text{CH}_3)_2\text{R}$, $\text{L}_2\text{Pd}(\text{CH}_3)_2$, and ArNiMeL_2 .²¹ In the absence of a polar, coordinating solvent, the L_2Pd^0 formed after reductive elimination of ethane disproportionates to metallic palladium and L_4Pd^0 , and for Au(III), the $\text{LAu}(\text{I})\text{R}$ species produced upon reductive elimination of R_2 undergoes further intermolecular decomposition to produce Au metal and free phosphine; in contrast, the thermal decomposition of ArNiMeL_2 in cyclohexane or benzene proceeds homogeneously to produce L_2Ni^0 as a stable product. Thus, the production of LPt^0 (an intermediate which is probably highly unstable) in a nonpolar solvent such as cyclohexane would have to be followed by rapid recombination with dissociated phosphine to prevent the formation of platinum metal.

An alternate mechanism could involve initial dissociation of trialkylphosphine, oxidative addition of a C-H bond of the remaining coordinated phosphine to produce a Pt(IV) intermediate, rate-limiting reductive elimination of 1,1-dimethylcyclopropane, and recombination of dissociated phosphine with reductive elimination of the phosphine C-H bond from the remaining Pt(II) intermediate to produce L_2Pt^0 (Scheme IV). The absence of a kinetic deuterium isotope effect upon substitution of $((\text{CD}_3)_2\text{CD})_3\text{P}$ for $((\text{CH}_3)_2\text{CH})_3\text{P}$ in the thermolysis of **2** ($2\text{-}(\text{L}^D)_2$) indicates only that the oxidative addition of a phosphine C-H bond, if it does occur, does not take place in the rate-limiting step. A number of reactions have been reported in which the oxidative addition of a trialkylphosphine C-H bond to Pt(II) has resulted either in the formation of a cyclometalated phosphine Pt(II) compound via a Pt(IV) intermediate²² or in the exchange of ^2H for ^1H in the alkyl groups of coordinated phosphines (in reactions performed in $\text{D}_2\text{O}/\text{CH}_3\text{COOD}$).²³ The observation of incorporation of ^2H into the ca. 1% neopentane produced from the thermal decom-

(21) For LAuR_3 , $\text{L} = \text{Ph}_3\text{P}$ and $\text{R} = \text{CH}_3$, C_2H_5 , $n\text{-C}_4\text{H}_9$, $i\text{-C}_3\text{H}_7$, and $t\text{-C}_4\text{H}_9$.¹⁰ For $\text{L}_2\text{Pd}(\text{CH}_3)_2$, $\text{L} = \text{PPh}_3$ and PPh_2CH_3 and $\text{L}_2 = \text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$.¹¹ For ArNiMeL_2 , $\text{L} = \text{Et}_3\text{P}$ and $\text{Ar} = o\text{-tolyl}$.¹³

(22) Goel, R. G.; Montemayor, R. G. *Inorg. Chem.* **1977**, *16*, 2183-2186. Cheney, A. J.; Mann, B. E.; Shaw, B. L.; Slade, R. M. *J. Chem. Soc. A* **1971**, 3833-3842. Al-Salem, N. A.; Empsall, H. D.; Markham, R.; Shaw, B. L.; Weeks, B. J. *Chem. Soc., Dalton Trans.* **1979**, 1972-1982.

(23) Kiffen, A. A.; Masters, C.; Raynand, L. *J. Chem. Soc., Dalton Trans.* **1975**, 853-857.





Figure 5. Gibbs free energy-reaction coordinate diagram illustrating the relative free energies of activation for reaction of **1** with neopentane or decomposition to L_2Pt^0 ($T\Delta S$ calculated for 0.1 M reactants and products).

position of $2\text{-}(\text{L}^D)_2$ might be due to a phosphine cyclometalation reaction in which the reductive elimination of a C-H bond instead of a C-C bond occurs a small percentage of the time. Since reductive elimination of a C-C bond from Pt(IV) is well documented^{9,24} and reports of this same reaction for Pt(II) are few,²⁵ the possibility of cyclometalation of a trialkylphosphine to produce a Pt(IV) intermediate from which reductive elimination of 1,1-dimethylcyclopropane could occur cannot be ignored, though no concrete evidence has been produced by this investigation to indicate such a reaction does take place.

Our study of the reactions of homogeneous Pt(II) compounds has been directed toward achieving the oxidative addition of an unactivated C-H bond of a hydrocarbon solvent to a homogeneous platinum species, so that comparisons of the mechanisms of the homogeneous and heterogeneous C-H bond activation reactions can be made. We have now completed the study of one particular system, that of the thermal decompositions of dineopentylbis(trialkylphosphine)platinum(II) and its product, bis(trialkylphosphine)-3,3-dimethylplatinacyclobutane (eq 4). One may ask



why the reaction proceeds only to the right, with no observable back reaction via oxidative addition of a C-H bond of neopentane or the solvent. The results of this investigation can be used to construct an argument which outlines the problems of trying to oxidatively add "unactivated" C-H bonds of hydrocarbon solvents to homogeneous platinum(II) species.

For the reaction $(\text{Et}_3\text{P})_2\text{Pt}[\text{CH}_2\text{C}(\text{CH}_3)_3]_2 \rightarrow \mathbf{1} + \text{neopentane}$ (178 °C), $k_{\text{calcd}} = 3.8 \times 10^{-2} \text{ s}^{-1}$,¹⁴ this rate corresponds to a $\Delta G^\ddagger \approx 30 \text{ kcal mol}^{-1}$. Two particles are produced from one in the reaction, resulting in a $T\Delta S$ (translation) = 15-17 kcal mol⁻¹ (178 °C, 0.1 M reactants and products).²⁶ The sums of bond energies in the reactant and products are the same to a first approximation (neglecting ring strain, vide infra), and one can make the sim-


(24) The reaction of cyclohexane solutions of **1** or **2** with bromine or iodine at 25 °C results in the immediate reductive elimination of 1,1-dimethylcyclopropane and formation of *trans*- L_2PtX_2 ($\text{X} = \text{Br}, \text{I}$); this reaction most likely proceeds through a Pt(IV) intermediate, from which reductive elimination of a C-C bond is rapid.

(25) The thermal decomposition of solid samples of *cis*- L_2PtR_2 ($\text{R} = \text{C}_6\text{H}_5$, $p\text{-CH}_3\text{C}_6\text{H}_4$; $\text{L} = \text{P}(\text{C}_6\text{H}_5)_3$, $\text{P}(p\text{-CH}_3\text{C}_6\text{H}_4)_3$; $\text{L}_2 = \text{dppe}$, dppm) in the absence of solvent resulted in the concerted, intramolecular reductive elimination of R_2 ; rates of decomposition were found to increase upon addition of added phosphine: Braterman, P. S.; Cross, R. J.; Young, G. B. *J. Chem. Soc., Dalton Trans.* **1976**, 1306-1314. Glockling, F.; McBride, T.; Pollock, R. J. *J. Chem. Soc., Chem. Commun.* **1973**, 650.

(26) Page and Jencks have calculated that for a reaction which produces two particles from one $T\Delta S \approx 10.5 \text{ kcal mol}^{-1}$ at 300 K: Page, M. I. *Angew. Chem., Int. Ed. Engl.* **1977**, *16*, 449-459. Page, M. I.; Jencks, W. P. *Proc. Natl. Acad. Sci. U.S.A.* **1971**, *68*, 1678-1683. Ibers, J. A.; DiCosimo, R.; Whitesides, G. M. *Organometallics* **1982**, *1*, 13.

plifying assumption that $\Delta H \approx 0$ and that ΔG for the reaction is primarily controlled by entropic effects. For the subsequent reaction of **1** \rightarrow $L_2Pt^0 + 1,1$ -dimethylcyclopropane (178 °C, with $Hg(0)$), $k_{obsd} = 1.2 \times 10^{-5} s^{-1}$. The actual reaction rate could be up to 10^2 faster than observed, since free triethylphosphine is produced during the reaction due to amalgamation of Pt(0) with $Hg(0)$ and may inhibit the rate of decomposition as was observed for thermolyses of **2** and **3**. This range of rate constants corresponds to a $\Delta G^\ddagger = 33$ –37 kcal mol $^{-1}$.²⁷ From this information one can construct a reaction coordinate diagram which illustrates the relative free energies of activation for the forward and back reactions of **1** (Figure 5); $\Delta G^\ddagger(back) - \Delta G^\ddagger(forward) \approx 8$ –14 kcal mol $^{-1}$.

The oxidative addition of a C-H bond of a solvent molecule to **1** should be more favorable than reaction with neopentane because of its greater relative concentration (eq 5); $\Delta S = R \ln$



[solvent]/[neopentane] $\approx R \ln 100 \approx 9.1$ eu. The hypothetical reaction of **1** with solvent should therefore have $T\Delta S = 4.1$ kcal mol $^{-1}$ (178 °C) more favorable than reaction of **1** with neopentane but still have a $\Delta G^\ddagger = 4$ –10 kcal mol $^{-1}$ greater than that for reductive elimination of 1,1-dimethylcyclopropane from **1**.

This entire argument is only qualitative in that it ignores changes in bond strengths and strain energies in proceeding from reactants to products and that it is based on very qualitative estimates of $T\Delta S$. Entropic factors alone are not necessarily the dominant contributors to ΔG for these reactions: steric crowding in $(Et_3P)_2Pt[CH_2C(CH_3)_2]$ is relieved upon loss of neopentane,²⁸ and ring strain in the platinacyclobutane which is produced is small but significant (<5 kcal mol $^{-1}$ greater than an analogous platinacyclopentane).^{14,29} Although the relative importance of entropic and enthalpic contributions in determining the free energy of these reactions have not been established, the above argument suggests that the highly favorable change in entropy in producing two particles from one, both for the production of **1** and for its subsequent decomposition, is significant in determining the direction of the reaction.

Experimental Section

General Data. All reactions involving organometallic compounds were performed under argon or prepurified dinitrogen by using standard anaerobic techniques.³⁰ The procedures followed for performing sealed tube reactions have been described previously.¹⁴ Boiling points are uncorrected. Melting points were determined on a Thomas-Hoover Apparatus and are uncorrected. $^{31}P\{^1H\}$ NMR spectra were recorded at 36.2 and 101.3 MHz on Jeol FX-90Q and Bruker WP-250 spectrometers, respectively. Spectra are reported at 36.2 MHz unless otherwise noted, and chemical shifts are reported relative to external 85% phosphoric acid

(27) At this same temperature (178 °C), the free energy of activation for decomposition of **2** or **3** is calculated to be $\Delta G^\ddagger \approx 26$ –27 kcal mol $^{-1}$. A comparison of the free energies of activation for addition of a C-H bond to **2** or **3** with their thermal decomposition to L_2Pt^0 and 1,1-dimethylcyclopropane is more difficult to construct than for **1**, since the corresponding dineopentylbis(trialkylphosphine)platinum(II) compounds could not be prepared in order to determine ΔG for formation of the platinacyclobutanes.

(28) Steric crowding about transition metal alkyls has been reported to result in a decrease of metal-carbon bond energies by 19–20 kcal mol $^{-1}$ for $[M(CH_2C(CH_3)_2)_4]$ ($M = Ti, Zr, \text{ or } Hf$) with respect to sterically less demanding alkyl substituents (Lappert, M. F.; Patil, D. S.; Padley, J. B. *J. Chem. Soc., Chem. Commun.* **1975**, 830–831). Differences in cobalt *n*-propyl and cobalt neopentyl bond energies of ~ 7 kcal mol $^{-1}$ have been observed in certain organocobalt complexes (Tsou, T.-T.; Loots, M.; Halpern, J. *J. Am. Chem. Soc.* **1982**, *104*, 623–624).

(29) The ring strain in the platinacyclobutane moiety of platinum(IV) complexes (e.g., $PtCl_2(C_3H_6)(bpy)$) has been estimated at ~ 12 kcal mol $^{-1}$ by comparisons of heats of decomposition measured by using differential scanning calorimetry (Puddephatt, R. J. *Coord. Chem. Rev.* **1980**, *33*, 149–194). Comparison of this estimate with that derived for similar platinum(II) complexes¹⁴ may not be appropriate, since steric crowding, bond strengths, and electronic structures of the two complexes may be different.

(30) Shriver, D. F. "The Manipulation of Air-Sensitive Compounds"; McGraw-Hill: New York, 1969.

(downfield shifts positive). 1H NMR spectra were recorded at 90 and 250 MHz, respectively, on these same spectrometers and are reported at 90 MHz unless otherwise specified. Chemical shifts are reported relative to internal tetramethylsilane. Mass spectra were recorded on a Hewlett-Packard 5990A GC/MS at an ionizing voltage of 70 eV. Diethyl ether and tetrahydrofuran were distilled from disodium benzophenone dianion under argon. Cyclohexane and cyclopentane were stirred over concentrated sulfuric acid for 48 h, washed with saturated sodium bicarbonate solution and then water, dried with phosphorus pentoxide, and distilled under argon. Triethyl phosphate was distilled and stored under argon. DCl (38% in D_2O , 99% D, Stohler), triisopropylphosphine (Strem), and tricyclohexylphosphine (Strem) were used as received. Dineopentyl(1,5-cyclooctadiene)platinum¹⁴ and bis(triethylphosphine)-3,3-dimethylplatinacyclobutane (**1**)¹⁴ were prepared as previously described. A published procedure was used to prepare 2-bromopropane-*d*₇ from 2-propanol-*d*₈ (Stohler, 99.5% D).³¹ Organoplatinum compounds were stored in tightly sealed vials in the dark at -15 °C and were freshly recrystallized before performing thermolyses.

Tris(isopropyl-*d*₇)phosphine. To a 500-mL round-bottomed flask equipped with stirring bar, addition funnel, and reflux condenser was added 7.0 g (0.29 mol) of magnesium metal and 200 mL of ether. The addition funnel was charged with a solution of 31 g (0.24 mol) of 2-bromopropane-*d*₇ in 30 mL of ether. This solution was added dropwise with stirring at a rate sufficient to maintain gentle reflux. The mixture was stirred for 3 h after the addition was complete and then filtered through a plug of glass wool to yield 230 mL of 0.97 M (isopropyl-*d*₇)-magnesium bromide (0.22 mol, 94% yield).

The deuterated phosphine was prepared according to a variation of a published procedure.³² To a flame-dried, argon-filled 500-mL round-bottomed flask equipped with stirring bar and addition funnel was added 6.1 mL (9.6 g, 0.070 mol) of freshly distilled phosphorous trichloride and 100 mL of ether. The addition funnel was charged with the Grignard solution described above, which was then added dropwise with stirring under argon at -10 °C. After the addition was complete, the mixture was refluxed for 2 h and cooled to room temperature. Degassed, distilled water (10 mL) was added slowly by syringe with stirring, followed by 50 mL of degassed 2 N HCl and 66 mL of degassed 2 N NH_4OH which dissolved the salts from the reaction and left two phases containing no solids. The organic layer was transferred by cannula to an argon-filled flask containing ca. 20 g of anhydrous sodium sulfate, the remaining aqueous layer was stirred with an additional 150 mL of ether, and the organic layers were combined and fractionally distilled under argon at atmospheric pressure. The fraction boiling at 170–175 °C (lit.³⁰ bp 175 °C) was collected and filtered under argon through a 0.5-cm \times 2.0-cm column of alumina (Woelm, Activity I) to yield 5.3 g (42% yield) of tris(isopropyl-*d*₇)phosphine: $^{31}P\{^1H\}$ NMR (C_6D_6) δ 17.3.

Reaction of Dineopentyl(1,5-cyclooctadiene)platinum(II) with Triisopropylphosphine and Tricyclohexylphosphine. Into a flame-dried, argon-filled 0.5-mL thermal decomposition tube was weighed 15 mg (0.034 mmol) of dineopentyl(1,5-cyclooctadiene)platinum(II); the tube was capped with a rubber septum and flushed with argon. To the tube was then added by syringe 0.10 mL (0.068 mmol) of a 0.68 M solution of triisopropylphosphine in cyclohexane or 0.25 mL (0.068 mmol) of a 0.27 M solution of tricyclohexylphosphine in cyclohexane, followed by 0.15 mL of a 0.102 M solution of cyclopentane (GLC internal standard) in cyclohexane. The contents of the tube were degassed by three freeze-pump-thaw cycles, and the tube was sealed under vacuum (0.01 torr). The tube was heated at 50 °C for 15 h, then frozen in liquid nitrogen, and opened under argon, and the reaction mixture examined by GLC for neopentane and 2,2,5,5-tetramethylhexane by using a 10-ft, $1/8$ -in. 20% SE-30 on Chromosorb P column and for 1,5-cyclooctadiene and 1,3,5-cyclooctatriene using a 10-ft, $1/8$ -in. 15% β,β' -oxydipropionitrile on Chromosorb P column. The yields of neopentane, 2,2,5,5-tetramethylhexane, 1,5-COD, and 1,3,5-cyclooctatriene were 80%, 36%, 80%, and 14%, respectively, for reaction with triisopropylphosphine and 85%, 15%, 30% and 3%, respectively, for reaction with tricyclohexylphosphine. Addition of 0.10 mL of concentrated HCl to these reaction mixtures produced an additional 53 and 83% yield of neopentane from the reactions with triisopropylphosphine and tricyclohexylphosphine, respectively, accounting for all the neopentyl groups originally on platinum.

Repeating the above procedure and heating the tube an additional 2 h at 126 °C after 15 h at 50 °C produced 1,1-dimethylcyclopropane and additional neopentane in 26% and 26% yields, respectively, for reaction

(31) Coulson, E. J.; Gerrard, W.; Hudson, H. R. *J. Chem. Soc.* **1965**, 2364–2369.

(32) Lorenz, H. J.; Zintl, A. R.; Volker, F. (Deutsche Advance Production G.m.b.H.) US 3 499 039 (Cl. 260-606.5; C 07F), Mar 3, 1970, Appl. Nov 10 1966 (CA 72 P132967s); Neth. Appl. 6,614,945 (Cl. C 07F), Apr 24, 1967; Ger. Appl. Oct 21, 1966 (CA 67 P108757z).

with triisopropylphosphine and in 44% and 41% yields for reaction with tricyclohexylphosphine. The yield of 1,1-dimethylcyclopropane corresponds to the fraction of the reaction of trialkylphosphine with dineopentyl(1,5-cyclooctadiene)platinum(II) which produces platinacyclobutane, determined by heating a reaction mixture at 50 °C for 15 h, removing all volatile components under vacuum, redissolving the remaining oil in dry cyclohexane, and adding 0.10 mL of 38% DCI in D₂O. The yields of (CH₃)₂C(CH₂D)₂ produced in this manner for reactions using triisopropylphosphine and tricyclohexylphosphine were 27% and 39%, respectively (Schemes I and II), as determined by GC/MS analysis on a 20-ft, 1/8-in. 3% UCW-98 on Chromosorb P column.

Bis(triisopropylphosphine)-3,3-dimethylplatinacyclobutane (2 and 2-(L^D)₂). Into each of three 12-in. × 0.5-in. medium-wall thermal decomposition tubes was weighed 0.89 g (2.0 mmol) of dineopentyl(1,5-cyclooctadiene)platinum(II); the tubes were capped with a rubber septum and flushed with argon. To each tube was then added by syringe 10 mL of dry, degassed cyclohexane and 0.83 mL (0.64 g, 4.0 mmol) of triisopropylphosphine; the tubes were degassed by three freeze-pump-thaw cycles and sealed under vacuum (0.01 torr). The tubes were heated at 50 °C for 18 h, then frozen in liquid nitrogen, opened, and capped with rubber septa under argon. The reaction mixtures were transferred by cannula to a flame-dried, argon-filled 100-mL round-bottomed flask equipped with stirring bar, and the solvent was removed under vacuum with stirring. The remaining yellow-brown solid was dissolved in 50 mL of degassed acetone and the solution cooled to -78 °C. The yellow precipitate was collected by suction filtration and redissolved in 25 mL of degassed acetone. The volume of the solution was reduced by rapidly flowing argon over the solution at -78 °C. The fine, white crystalline precipitate was collected by suction filtration and remaining acetone removed under vacuum to yield 0.68 g (19%) of **2**: mp 126–129 °C dec; ³¹P{¹H} NMR (C₆D₆) δ 33.0 (t, J_{Pt-P} = 1831 Hz); ¹H NMR (C₆D₆) δ 0.86 (4 H, t, J_{Pt-H} = 75.2 Hz, J_{P-H} = 10.3 Hz, PtCH₂), 1.05 (36 H, d of d, J_{P-H} = 11.7 Hz, J = 6.8 Hz, (CH₃)₂CHP), 1.49 (6 H, b s, C-(CH₃)₂), 2.17 (6 H, septet, J = 7.3 Hz, CHP); IR (KBr) 2990–2750 (vs br), 1450 (s), 1370 (s), 1350 (s), 1340 (m), 1235 (s), 1038 (s), 1019 (s), 915 (m), 872 (s), 757 (m), 632 (vs), 519 (c) cm⁻¹. Anal. (C₂₃H₅₂P₂Pt) C, H.

Compound **2-(L^D)₂** was prepared in the same manner as described for **2**, using 0.94 mL (0.72 g, 4.0 mmol) of tris(isopropyl-*d*₇)phosphine for each 0.89 g (2.0 mmol) of dineopentyl(1,5-cyclooctadiene)platinum(II) (21% yield): mp 129.0–130.5 °C dec; ³¹P{¹H} NMR (C₆D₆) δ 31.8 (t, J_{Pt-P} = 1834 Hz); ¹H NMR (C₆D₆) δ 0.85 (4 H, t, J_{Pt-H} = 76.2 Hz, J_{P-H} = 9.8 Hz, PtCH₂), 1.49 (6 H, b s, C(CH₃)₂); IR (KBr) 2920 (s), 2860 (s), 2820 (m), 2205 (m), 2190 (s), 2130 (m), 2100 (m), 2060 (m), 1450 (m), 1360 (m), 1310 (m), 1055 (m), 1018 (s), 805 (m), 785 (m), 595 cm⁻¹. Anal. (C₂₃H₁₀D₄₂P₂Pt) C, P.

Bis(tricyclohexylphosphine)-3,3-dimethylplatinacyclobutane (3). Into each of three 12-in. × 0.5-in. medium-wall thermal decomposition tubes was weighed 0.60 g (1.4 mmol) of dineopentyl(1,5-cyclooctadiene)platinum(II); the tubes were capped with a rubber septum and flushed with argon. To each tube was added 8.0 mL (2.8 mmol) of a 0.34 M solution of tricyclohexylphosphine in cyclohexane. The tube walls were washed down with an additional 2.0 mL of cyclohexane, and the resulting solutions were degassed by two freeze-pump-thaw cycles and sealed under vacuum (0.01 torr). The tubes were heated at 50 °C for 15 h and then opened under argon and the reaction mixtures transferred by cannula to an argon-filled 50-mL round-bottomed flask equipped with stirring bar. The volatile components of the reaction mixture were removed under vacuum while being heated at 40–50 °C, and the remaining red-brown oil was redissolved in 150 mL of degassed acetone and cooled to -10 °C for 12 h. The pale yellow solid which precipitated was collected by suction filtration and dissolved in ca. 100 mL of THF. The resulting solution was filtered through a 0.5-μm millipore filter and the filtrate reduced in volume to 10 mL by rotary evaporation at 25 °C. Acetone (50 mL) was added, and the resulting solution was cooled to -78 °C. The white, crystalline solid which precipitated was collected by suction filtration and remaining solvent removed under vacuum to yield 0.56 g (17%) of **3**: mp 155–156 °C (dec); ³¹P NMR (C₆D₆) δ 22.3 (t, J_{Pt-P} = 1797 Hz); ¹H NMR (C₆D₆) δ 0.98 (4 H, m, J_{Pt-H} = 75.1 Hz, PtCH₂), 1.27 (18 H, m, CH₂), 1.58 (6 H, s, C(CH₃)₂), 1.59–1.99 (30 H, b m), 2.14 (18 H, m); IR (KBr) 2990–2870 (vs br), 2845 (vs), 1443 (s), 1345 (m), 1295 (m), 1268 (m), 1172 (m), 1122 (m), 1002 (m), 845 (s), 729 (m) cm⁻¹. Anal. (C₄₁H₇₆P₂Pt) C, H, P.

Thermal Decomposition of Bis(triethylphosphine)-3,3-dimethylplatinacyclobutane (1) in the Presence of Hg(0). Into each of nine flame-dried, argon-filled 0.5-mL medium-wall thermal decomposition tubes was placed 0.40 g of triply distilled mercury. The tubes were capped with rubber septa and flushed with argon, and then 0.10 mL of a solution of 0.150 g (0.299 mmol) of freshly recrystallized **1** dissolved in 1.50 mL of a 0.102 M solution of cyclopentane in cyclohexane was

added to each tube by syringe. The constriction in each tube was rinsed with 0.03 mL of cyclohexane. The contents of the tubes were degassed and the tubes sealed under vacuum (0.01 torr). The tubes were heated for predetermined intervals at 178 °C, then frozen in liquid nitrogen, and opened and the reaction mixtures analyzed by GLC for production of 1,1-dimethylcyclopropane. The observed rate constant for the reaction was measured from data collected through 1 half-life of the reaction; *k*^{obsd} = 1.2 × 10⁻⁵ s⁻¹. At the completion of the reaction, the yields of 1,1-dimethylcyclopropane and neopentane were 92% and 9%, respectively.

Thermal Decomposition of 2 and 3. Into a 5-mm NMR tube was weighed 30 mg (0.052 mmol) of **2**; the tube was capped with a rubber septum and flushed with argon. To the tube was added by syringe 0.50 mL of a 0.108 M solution of cyclopentane in cyclohexane; the contents of the tube were degassed, and the tube was sealed under vacuum. The tube was then heated for 2 h at 126 °C and the pale yellow, homogeneous reaction mixture examined by ³¹P{¹H} NMR. No starting material remained, and the only product observed was bis(triisopropylphosphine)platinum(0)³³ δ 73.2 (t, 1:4:1, J_{Pt-P} = 4229 Hz) (lit.³⁴ (octane) δ 72.8 (J_{Pt-P} = 4202 Hz)). The tube was opened under argon and the reaction mixture examined by GLC on a 10-ft, 1/8-in. 20% SE-30 on Chromosorb P column, which indicated the formation of 1,1-dimethylcyclopropane (99%) and neopentane (2%).

Repeating the above procedure with 30 mg (0.036 mmol) of **3** produced bis(tricyclohexylphosphine)platinum(0) as the sole product by ³¹P{¹H} NMR: δ 62.5 (t, 1:4:1, J_{Pt-P} = 4229 Hz) (lit.³⁴ (heptane) δ 63.8 (J_{Pt-P} = 4227 Hz)). Analysis of the pale yellow, homogeneous mixture by GLC indicated the formation of 1,1-dimethylcyclopropane (94%) and neopentane (7%).

Kinetics of Thermal Decomposition of 2, 2-(L^D)₂, and 3. Into each of eight flame-dried, argon-filled 5-mm NMR tubes was weighed 23.0 mg (0.040 mmol) of freshly recrystallized **2**. The tubes were capped with rubber septa and flushed with argon. To each tube was added by syringe 0.50 mL of a 0.16 M solution of triethyl phosphate in cyclohexane; the tubes were degassed by three freeze-pump-thaw cycles and sealed under vacuum (0.01 torr). Duplicate thermolyses at four different temperatures were performed simultaneously, following the disappearance of starting material by ³¹P{¹H} NMR spectroscopy through at least 2 half-lives. The observed rate constants (average of two runs) for thermolyses of **2** at 98.5, 111.0, 116.5, and 125.0 °C were *k*^{obsd} = 5.70, 39.4, 84.6, and 401 × 10⁻⁵ s⁻¹, respectively.

Repeating the above procedure using 10.0 mg (0.0121 mmol) of **3** in each tube, and heating the tubes at 50 °C for 5 min to completely dissolve the platinum complex before beginning the kinetic runs, gave the following rate constants (average of two runs) at 90.5, 98.0, 109.5, and 116.0 °C: *k*^{obsd} = 2.41, 7.71, 44.8, and 105 × 10⁻⁵ s⁻¹, respectively.

The rate of production of 1,1-dimethylcyclopropane from the thermal decomposition of **2**, **2-(L^D)₂**, and **3** was measured by GLC. In a typical procedure, 93.6 mg (0.160 mmol) of **2** was dissolved in 1.0 mL of a 0.102 M solution of cyclopentane in cyclohexane and the resulting solution diluted to 2.0 mL with cyclohexane in a 2.0-mL volumetric flask. To each of nine 0.5-mL medium-wall thermal decomposition tubes was added 0.20 mL of this standard solution, and the walls of the tubes were washed down with an additional 0.10 mL of cyclohexane. The tubes were degassed three times and sealed under vacuum (0.01 torr), then heated at 116.5 °C for predetermined intervals, and opened, and the yield of 1,1-dimethylcyclopropane was determined. The observed rate constants for **2** and **2-(L^D)₂** at 116.5 °C were *k*^{obsd} = 8.20 and 8.39 × 10⁻⁴ s⁻¹, respectively, and the rate constant for **3** at 116.0 °C was *k*^{obsd} = 9.88 × 10⁻⁴ s⁻¹.

Thermal Decomposition of 2 and 3 in the Presence of Added Trialkylphosphine. Cyclohexane solutions 0.0075, 0.0125, 0.025, 0.050, and 0.10 M in triisopropylphosphine were prepared by diluting 0.075, 0.125, 0.25, 0.50, and 1.0 mL of a 0.20 M stock solution of the phosphine in cyclohexane to 2.0 mL in volumetric flasks under argon, and then 0.20 mL of each of these solutions was added by syringe to a 5-mm NMR tube containing 0.30 mL of a cyclohexane solution 0.133 M in **2** and 0.16 M in triethyl phosphate. The tubes were degassed and sealed under vacuum and then heated at 126.5 °C, and the rate of disappearance of **2** was measured relative to triethyl phosphate internal standard by ³¹P{¹H} NMR spectroscopy. The observed rate constants for reaction mixtures 80 mM in **2** and 3.0, 5.0, 10, 20, and 40 mM in triisopropylphosphine were *k*^{obsd} = 119, 67.2, 33.4, 21.8, and 8.79 × 10⁻⁵ s⁻¹, respectively.

(33) Reaction of the remaining bis(triisopropylphosphine)platinum(0) with 1 equiv of methyl iodide resulted in the immediate production of *trans*-(i-Pr₃P)₂Pt(I)CH₃: mp 178–180 °C dec; ³¹P{¹H} NMR (C₆D₆) δ 28 (t, J_{Pt-P} = 2801 Hz); ¹H NMR (C₆D₆) δ 0.77 (3 H, t, J_{Pt-H} = 88.9 Hz, PtCH₃), 1.23 (36 H, d of d, J_{P-H} = 13.7 Hz, J = 6.8 Hz, (CH₃)₂CHP), 2.95 (6 H, m, (CH₃)₂CHP). Anal. (C₁₉H₄₅IPt) C, H, I.

(34) Mann, B. E.; Musco, A. *J. Chem. Soc., Dalton Trans.* **1980**, 776–785.

Thermal decomposition of a cyclohexane solution of **3** (0.024 M) and tricyclohexylphosphine (0.028 M) at 116 °C as described above yielded $k^{\text{obsd}} = 5.93 \times 10^{-5} \text{ s}^{-1}$, compared to $k^{\text{obsd}} = 105 \times 10^{-5} \text{ s}^{-1}$ in the absence of added phosphine.

Phosphine Exchange of 2 with 2-(L^D)₂. A degassed cyclohexane solution of **2** (86 mM), 2-(L^D)₂ (86 mM), and triethyl phosphite (16 mM) was heated for 8 h at 50 °C in a 5-mm NMR tube. Examination of the resulting solution by ³¹P{¹H} NMR (C₆H₁₂, 101.28 MHz) revealed the presence of the following compounds: **2**, δ 33.45 (t, 1:4:1, $J_{\text{Pt-P}} = 1816$ Hz); 2-(L^D)₂, δ 32.16 (t, 1:4:1, $J_{\text{Pt-P}} = 1820$ Hz); 2-L^HL^D, δ 33.54 (d of t, $J_{\text{Pt-P}} = 1818$ Hz, $J_{\text{P-P}} = 4$ Hz). The doublet of triplets for L^D of 2-L^HL^D could not be resolved from the signal for 2-(L^D)₂ due to the

broadness of the peak. No bis(triisopropylphosphine)platinum(0) was observed, and heating the solution for an additional 16 h at 50 °C produced no change in the distribution of products.

Registry No. **1**, 70620-74-5; **2**, 81602-78-0; 2-(L^P)₂, 81602-79-1; **3**, 81602-80-4; tris(isopropyl)phosphine, 6476-36-4; 1,1-dimethylcyclopropane, 1630-94-0; neopentane, 463-83-1; 2,2,5,5-tetramethylhexane, 1071-81-4; 1,3,5-cyclooctatriene, 1871-52-9; tris(isopropyl-*d*₂₁)phosphine, 81583-58-6; isopropyl-*d*₇ bromide, 39091-63-9; 1,5-COD, 111-78-4; [(*i*-Pr)₃P]₂Pt, 60648-71-7; (1,5-COD)Pt[CH₂C(CH₃)₃]₂, 75101-19-8; (CH₃)₂C(CH₂D)₂, 72110-09-9; (CH₃)₃C(CH₂D), 4741-94-0; Cy₃P, 2622-14-2; (Cy₃P)₂Pt, 55664-33-0.

Synthesis and Characterization of Coordination Compounds of Chelating Ligands Containing Imidazole Groups. The Crystal and Molecular Structures of the Dinuclear Cu^I and Cu^{II} Compounds [*N,N,N',N'*-Tetrakis-(2-benzimidazolylmethyl)-1,2-ethanediamine]dicopper(I) Bis(perchlorate) and of μ -(Nitrato-*O,O'*)-bis(nitrato-*O*)-{*N,N,N',N'*-tetrakis[(1-methyl-2-benzimidazolyl)methyl]-1,2-ethanediamine}dicopper(II) Nitrate Tetrahydrate

Hugo M. J. Hendriks, Paul J. M. W. L. Birker, Jacobus van Rijn, Gerrit C. Verschoor, and Jan Reedijk*

Contribution from the Department of Chemistry, Gorlaeus Laboratories, State University of Leiden, 2300 RA Leiden, The Netherlands. Received July 27, 1981

Abstract: Copper(I) and copper(II) coordination compounds with general formulas Cu^{II}(L)(anion)₂, Cu^{II}(L)(anion)₄, and Cu^I₂(L)(anion)₂ have been prepared and characterized, where L stands for one of the sexadentate benzimidazole-containing ligands *N,N,N',N'*-tetrakis(2-benzimidazolylmethyl)-1,2-ethanediamine (abbreviated EDTB), *N,N,N',N'*-tetrakis[(1-methyl-2-benzimidazolyl)methyl]-1,2-ethanediamine (NMEDTB), *N,N,N',N'*-tetrakis[(1-benzyl-2-benzimidazolyl)methyl]-1,2-ethanediamine (NBEDTB), *N,N,N',N'*-tetrakis[(5-methyl-2-benzimidazolyl)methyl]-1,2-ethanediamine (5-MEDTB), or *N,N,N',N'*-tetrakis[(5,6-dimethyl-2-benzimidazolyl)methyl]-1,2-ethanediamine (5,6-DMEDTB). The electronic and ESR spectra of all compounds of composition Cu^{II}(L)(anion)₂ are identical with those for Cu(EDTB)(anion)₂ and suggest a structure similar to the one found for Cu(EDTB)(BF₄)(BF₃OC₂H₅), which has been described as highly distorted square pyramidal. ESR and electronic spectra of the compounds Cu₂(L)(anion)₄ indicate different coordination geometries in the presence of different anions. In compounds with Cl⁻ or Br⁻ the Cu^{II} atom appears to have trigonal-bipyramidal coordination geometry. Square-pyramidal geometry seems more likely for compounds Cu₂(L)X₄, where X = NO₃⁻, BF₄⁻, or ClO₄⁻. The crystal and molecular structures of [Cu^{II}₂(NMEDTB)(NO₃)₃](NO₃)₃·4H₂O (**1**) and of Cu^{II}₂(EDTB)(ClO₄)₂ (**2**) have been determined by single-crystal X-ray analysis. Both compounds crystallize in space group C2/c. For compound **1**, $a = 19.159$ (4) Å, $b = 15.907$ (2) Å, $c = 16.816$ (3) Å, $\beta = 109.57$ (2)°, $V = 4829.0$ Å³, $Z = 4$, $d_{\text{measd}} = 1.51$ (1) g cm⁻³, and $d_{\text{calcd}} = 1.49$ g cm⁻³. The intensities of 4241 independent reflections were measured on an automatic diffractometer, 2670 of which were considered as observed ($I > 2\sigma(I)$). For compound **2**, $a = 13.849$ (2) Å, $b = 18.913$ (2) Å, $c = 14.062$ (2) Å, $\beta = 102.10$ (1)°, $V = 3592.2$ Å³, $Z = 4$, $d_{\text{measd}} = 1.65$ (1) g cm⁻³, and $d_{\text{calcd}} = 1.68$ g cm⁻³. A total of 3522 independent reflections were measured, 2627 of which were considered as observed. Both structures were solved by heavy-atom methods and refined by using least-squares techniques to a residual R value of 0.046 for **1** and 0.039 for **2**. Structure **1** consists of dinuclear Cu₂(NMEDTB)(NO₃)₃⁺ cations having C₂ symmetry, a disordered nitrate ion, and disordered water molecules. The coordination geometry around each copper ion can be described as square pyramidal with two equatorial benzimidazole nitrogen atoms, one equatorial amine nitrogen atom, one equatorial oxygen atom of a monodentate coordinating nitrate ion, and one axial oxygen atom of a nitrate ion which is bridging to the axial position of the other copper ion in the cation. Structure **2** consists of dinuclear Cu₂(EDTB)²⁺ cations having C₂ symmetry and perchlorate ions. Each copper is linearly coordinated by two benzimidazole nitrogen atoms, with short Cu-N bonds of 1.869 (4) and 1.876 (4) Å, respectively. The Cu-Cu distance within one unit is 3.043 (1) Å. The structure of the Cu₂(EDTB)²⁺ ion shows a strong resemblance with a recently proposed model for the reduced state of hemocyanin. The compounds Cu₂(L)²⁺ do react with O₂ in a 1:1 molar ratio although no O₂ adducts could be isolated.

The oxygen-transporting copper protein hemocyanin has attracted much attention.¹ In a recently proposed² model for the

active site in hemocyanin, two Cu(I) ions separated by ~3.4 Å are coordinated by two (or three) imidazoles of histidine residues.³