

The reaction of $(\eta\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO})_2$ with sulfoxides

(Received December 15th, 1988)

** Authors to whom correspondence should be addressed.

Table 1


Reduction of sulfoxides with $\text{Cp}_2\text{Ti}(\text{CO})_2$ ^a

	Sulfoxide		Molar ratio 1/ $\text{Cp}_2\text{Ti}(\text{CO})_2$	Temperature	Time (h)	Sulfide (2) yield ^b
	R ¹	R ²				
1a	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	1/1	reflux THF	2	93
	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	2/1	reflux THF	2	91
1b	<i>p</i> -ClC ₆ H ₄	<i>p</i> -ClC ₆ H ₄	1/1	reflux THF	2	94
	<i>p</i> -ClC ₆ H ₄	<i>p</i> -ClC ₆ H ₄	2/1	reflux THF	2	91
1c	C ₆ H ₅	C ₆ H ₅	1/1	reflux THF	2	91
	C ₆ H ₅	C ₆ H ₅	2/1	reflux THF	4	96
	C ₆ H ₅	C ₆ H ₅	2/1	room temp.	40	96
1d	CH ₃	CH ₃	2/1	reflux THF	2	74 ^c
1e	<i>n</i> -C ₃ H ₇	<i>n</i> -C ₃ H ₇	1/1	reflux THF	2	81 ^c
1f	<i>n</i> -C ₄ H ₉	<i>n</i> -C ₄ H ₉	1/1	reflux THF	6	89
1g	C ₆ H ₅ CH ₂	C ₆ H ₅ CH ₂	1/1	reflux THF	6	93
1h	C ₆ H ₅	CH ₂ =CH-	2/1	40 ° C	5	80

^a All products were identified by comparison (IR and NMR) with authentic samples. ^b Yield of products isolated by column chromatography. ^c Yield determined by GC or NMR.

THF (13 ml) for 2 h, a stable yellow suspension formed and *p*-tolylsulfide was obtained (91%) after the same work-up procedure. The identities of the sulfide products were confirmed via spectroscopic comparison with an authentic sample.

Both aliphatic and aromatic sulfoxides are readily deoxygenated when treated with $\text{Cp}_2\text{Ti}(\text{CO})_2$ in THF under nitrogen in the temperature range 25° to refluxing THF. As shown in Table 1, the isolated yield of the sulfide product is generally high. The high yield and purity of the product is of interest with respect to the mechanism of this reaction. The reduction of sulfoxide to sulfide is not accompanied by Pummerer rearrangement product (3) which has sometimes been observed with the use of several other reagents [2,6,7]. This rearrangement is usually facilitated by the presence of a strong acid [2,6,7] hence the absence of 3 suggests that $\text{Cp}_2\text{Ti}(\text{CO})_2$ does not behave as a strong acid in this case. In addition benzyl



mercaptan, which is occasionally formed during reduction of benzyl sulfoxide (1g) under other conditions [13], could not be detected by TLC. While the most efficient reduction of sulfoxide is obtained using 1 mole equivalent of $\text{Cp}_2\text{Ti}(\text{CO})_2$, only 0.5 mole equivalent is required. This indicates greater reducing capacity than simple oxidation of Ti^{III} to Ti^{IV} would suggest. The formal similarity of these reactions to the reduction of amine oxides by metal carbonyls to give amine complexes and CO₂ is noted. Finally $\text{Cp}_2\text{Ti}(\text{CO})_2$ is a mild reductant; sulfones (4) and phosphine oxides (5) are inert under these reaction conditions.

The reducing agent, $\text{Cp}_2\text{Ti}(\text{CO})_2$, is a well characterized species [15,16] and the reduction is carried out under homogeneous reaction conditions. For many other low valent transition metal reagents [9–12], the exact nature of the starting material is not well characterized and the reaction is done under heterogeneous reaction

conditions. The reactivity of $\text{Cp}_2\text{Ti}(\text{CO})_2$ was followed by infrared spectroscopy. During the reaction carbon monoxide was released there being no evidence in the IR spectra of the presence of a monocarbonyl intermediate. In the absence of sulfoxide the reaction gave a dark blue precipitate after refluxing for 4 h suggesting the possibility that THF is also a possible oxidant for $\text{Cp}_2\text{Ti}(\text{CO})_2$. For all the sulfoxides studied here, when the reaction was carried out with a ratio of Ti reagent to sulfoxide of 1/1, the corresponding sulfide and an air-sensitive blue titanium-containing species was obtained; whereas with a ratio of Ti/sulfoxide of 1/2, the sulfide and a yellow titanium-containing polymer was obtained. The latter was similar to that which formed upon exposure of the 1/1 ratio reaction to air.

The IR spectrum of the yellow polymer displayed bands consistent with the presence of the C_5H_5 ligand and Ti–O–Ti linkages. Elemental analysis indicated that the polymer had not retained two cyclopentadienyl rings per titanium [17]. This was further confirmed by the isolation of cyclopentadiene in the volatile distillate from the reaction mixture. A blue colour is characteristic of reduced cyclopentadienyloxotitanium complexes which are easily oxidized to give yellow oxotitanium(IV) polymers [17–19]. Further investigation will be necessary before a reasonable mechanism can be suggested. Nevertheless, present results indicate that $\text{Cp}_2\text{Ti}(\text{CO})_2$ is an effective agent for the deoxygenation of sulfoxides. The reaction also suggests a possible route to early transition metal oxo complexes via the use of mild, non-protic [20] oxygen transfer reagents.

Acknowledgements. We thank NSERC of Canada of FCAC of Quebec for financial support. The award of a fellowship by Academia Sinica to TLC is gratefully acknowledged.

References

- 1 T.L. Chen, T.H. Chan and A. Shaver J. Organomet. Chem., 268 (1984) C1.
- 2 T.H. Chan, A. Melnyk and D.N. Harpp, Tetrahedron Lett., (1969) 201.
- 3 T.H. Chan and A. Melnyk, J. Amer. Chem. Soc., 92 (1970) 3718.
- 4 (a) K. Naumann, G. Zon and K. Mislow, J. Am. Chem. Soc., 91 (1969) 2788; (b) 91 (1969) 7012.
- 5 I.W.J. Still, S.K. Hasan and K. Turnbull, Synthesis, (1977) 468.
- 6 M. Wakisaka, M. Hatanaka, H. Nitta, M. Hatamura and T. Ishimaru, Synthesis, (1980) 67.
- 7 J. Drabowicz and S. Oae, Synthesis, (1977) 404.
- 8 Y. Akita, M. Inaba, H. Uchida and A. Ohta, Synthesis, (1977) 792.
- 9 J. Drabowicz and M. Mikolajczyk, Synthesis, (1978) 138.
- 10 D. Zhemilev, Izv. Akad. Nauk. SSSR Ser. Khim., (1980) 734.
- 11 S. Kano, Y. Takana, E. Sugino and S. Hibino, Synthesis, (1980) 695.
- 12 G.A. Olah, G.K.S. Prakash and T.L. Ho, Synthesis, (1976) 810.
- 13 S.K. Chung and G. Han, Synth. Comm., (1982) 903.
- 14 R.G. Nazzo, H.J. Simon and J. San Filipo Jr., J. Org. Chem., 42 (1977) 568.
- 15 B. Demerseman, G. Bouquet and M. Bigorgne, J. Organomet. Chem., 101 (1975) C24.
- 16 G. Fachinetti, C. Floriani, F. Marchetti and M. Mellini, J. Chem. Soc. Dalton, (1978) 1398.
- 17 S.A. Giddings, Inorg. Chem., 3 (1964) 684.
- 18 (a) F. Bottomley and H.H. Brintzinger, J. Chem. Soc. Chem. Commun., (1978) 234; (b) F. Bottomley, I.J.B. Lin and M. Mukaida, J. Am. Chem. Soc., 102 (1980) 5238.
- 19 J.C. Huffman, J.G. Stone, W.C. Krusell and K.G. Caulton, J. Am. Chem. Soc., 99 (1977) 5829.
- 20 F. Bottomley, D.F. Drummond, G.O. Egharevba and P.S. While, Organometallics, 5 (1986) 1620.